

**MEDIQUE BAN-ACID MAXIMUM STRENGTH 750- calcium carbonate tablet, chewable**

**Unifirst First Aid Corporation**

-----

**Medique Ban-Acid**

***Drug Facts***

***Active ingredient (in each tablet)***

Calcium Carbonate USP 750 mg

***Purpose***

Antacid

***Uses***

relieves

- acid indigestion
- sour stomach
- heartburn
- upset stomach associated with these symptoms

***Warnings***

***Warnings***

**Do not use**

■ the maximum dosage of this product for more than 2 weeks except under the advice and

supervision of a doctor

**Ask a doctor or pharmacist before use if you**

- are taking a prescription drug. Antacids may interact with certain prescription drugs.
- have kidney disease

**When using this product**

- do not take more than 10 tablets in a 24 hours
- if pregnant, do not take more than 6 tablets in 24 hours

**Keep out of the reach of children.**

***Directions***

■ do not take for symptoms that persist for more than 2 weeks unless advised by a doctor

**Adults and children: (12 years and over):** Chew 2 to 4 tablets as symptoms occur, or as directed by a doctor.

**Children under 12 years:** Ask a doctor

***Other information***

- **calcium content per tablet:** 300 mg
- 5 calories per tablet
- store at room temperature 59-86°F (15-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

### Inactive ingredients

assorted flavors, dextrose, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, magnesium stearate, maltodextrin

**Questions or comments?** Call 1-800-634-7680

## Medique Ban-acid Label

Medique®

Ban-acid

Maximum Strength 750

Calcium Rich Extra Strength Antacid

Antiácido extra fuerte y /ricas en calcio

Pull To Open

Tire Para Abrir

✓ Heartburn, Indigestion, •Calcium Carbonate 750mg

las agruras, la indigestion •Carbonato de Calcio 750mg

Assorted Berry

Bayas surtidas

100

Tablets

(50 x 2)

Tamper Evident Unit Dose Packets

Empaquetado con Sellado Evidente en Dosis Unitarias

<table border="1"> <tr> <td> <b>Drug Facts</b>  <b>Active ingredient (in each tablet) Purpose</b>            Calcium Carbonate USP 750 mg Antacid   <b>Uses</b>            relieves            ■ acid indigestion            ■ sour stomach            ■ heartburn            ■ upset stomach associated with these symptoms   <b>Warnings</b>            Do not use            ■ the maximum dosage of this product for more than 2 weeks except under the advice and supervision of a doctor.            Ask a doctor or pharmacist before use if you            ■ are taking a prescription drug. Antacids may interact with certain prescription drugs.            ■ have kidney disease   <b>When using this product</b>            ■ do not take more than 10 tablets in 24 hours            ■ if pregnant, do not take more than 6 tablets in 24 hours            Keep out of the reach of children.         </td> <td> <b>Drug Facts (continued)</b>   <b>Directions</b>            ■ do not take for symptoms that persist for more than 2 weeks unless advised by a doctor   <b>Adults and children:</b> Chew 2 to 4 tablets as symptoms occur, or as directed by a doctor   <b>Children under 12 years:</b> Ask a doctor   <b>Other information</b>            ■ calcium content per tablet: 300 mg            ■ 5 calories per tablet            ■ store at room temperature 59-86°F (15-30°C)            ■ tamper-evident sealed packets            ■ do not use any opened or torn packets   <b>Inactive ingredients</b>            assorted flavors, dextrose, FD&amp;C blue #1 aluminum lake, FD&amp;C red #40 aluminum lake, magnesium stearate, maltodextrin   <b>Questions or comments</b>            Call 1-800-634-7680         </td> </tr> </table>	<b>Drug Facts</b> <b>Active ingredient (in each tablet) Purpose</b> Calcium Carbonate USP 750 mg Antacid  <b>Uses</b> relieves ■ acid indigestion ■ sour stomach ■ heartburn ■ upset stomach associated with these symptoms  <b>Warnings</b> Do not use ■ the maximum dosage of this product for more than 2 weeks except under the advice and supervision of a doctor. Ask a doctor or pharmacist before use if you ■ are taking a prescription drug. Antacids may interact with certain prescription drugs. ■ have kidney disease  <b>When using this product</b> ■ do not take more than 10 tablets in 24 hours ■ if pregnant, do not take more than 6 tablets in 24 hours Keep out of the reach of children.	<b>Drug Facts (continued)</b>  <b>Directions</b> ■ do not take for symptoms that persist for more than 2 weeks unless advised by a doctor  <b>Adults and children:</b> Chew 2 to 4 tablets as symptoms occur, or as directed by a doctor  <b>Children under 12 years:</b> Ask a doctor  <b>Other information</b> ■ calcium content per tablet: 300 mg ■ 5 calories per tablet ■ store at room temperature 59-86°F (15-30°C) ■ tamper-evident sealed packets ■ do not use any opened or torn packets  <b>Inactive ingredients</b> assorted flavors, dextrose, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, magnesium stearate, maltodextrin  <b>Questions or comments</b> Call 1-800-634-7680	 <p><b>Medique®</b> <b>Ban-acid®</b> Maximum Strength 750 Calcium Rich Extra Strength Antacid Antiácido extra fuerte y /ricas en calcio</p> <p>Pull To Open Tire Para Abrir</p> <p>✓ Heartburn, Indigestion • Calcium Carbonate 750 mg las agruras, la indigestion • Carbonato de Calcio 750 mg</p> <p>Assorted Berry Bayas surtidas</p> <p>100 Tablets (50 x 2)</p> <p>Tamper Evident Unit Dose Packets Empaquetado con Sellado Evidente en Dosis Unitarias</p>	<table border="1"> <tr> <td> <b>Información del Medicamento</b>  <b>Ingrediente activo (en cada tableta)</b>            Carbonato de Calcio 750 mg   <b>Usos</b>            alivia            ■ indigestión ácida ■ acidez estomacal ■ agruras            ■ malestar estomacal (debido a estos síntomas)   <b>Advertencias</b>            No consuma            ■ la dosis máxima de este producto por más de 2 semanas, excepto bajo indicaciones y supervisión médica.   <b>Consulte con el doctor o un farmacéutico antes de consumir este medicamento si usted está</b>            ■ actualmente tomando un medicamento recetado. Los antiácidos pueden interactuar con ciertos medicamentos recetados.            ■ tiene una enfermedad renal   <b>Cuando administre este producto</b>            ■ no tomar más de 10 tabletas en 24 horas            ■ si está embarazada, no tomar más de 6 tabletas en 24 horas            Mantenga este medicamento fuera de alcance de los niños.         </td> <td> <b>Propiedad</b>            Antiácido   <b>Información del medicamento (continuación)</b>  <b>Instrucciones</b>            ■ no tomar para síntomas que persisten por más de dos semanas a menos que lo recomiende un doctor   <b>Adultos y niños:</b> Masticar de 2 a 4 tabletas a medida que aparecen los síntomas o como lo indique un médico.   <b>Niños menores de 12 años:</b> consultar al médico   <b>Otra información</b>            ■ contenido de calcio por comprimido: 300 mg            ■ 5 calorías por tableta            ■ almacenamiento: consérvase a temperatura ambiente 59-86°F (15-30°C) en un área seca            ■ empaquetado con sellado evidente            ■ no use ningún paquete abierto o roto   <b>Ingredientes inactivos</b>            saborizantes surtidos, dextrosa, azul FD y C #1 lacas de aluminio, rojo FD y C #40 lacas de aluminio, estearato de magnesio, maltodextrina   <b>¿Tiene alguna pregunta o comentario?</b>            Llame al 1-800-634-7680         </td> </tr> </table>	<b>Información del Medicamento</b> <b>Ingrediente activo (en cada tableta)</b> Carbonato de Calcio 750 mg  <b>Usos</b> alivia ■ indigestión ácida ■ acidez estomacal ■ agruras ■ malestar estomacal (debido a estos síntomas)  <b>Advertencias</b> No consuma ■ la dosis máxima de este producto por más de 2 semanas, excepto bajo indicaciones y supervisión médica.  <b>Consulte con el doctor o un farmacéutico antes de consumir este medicamento si usted está</b> ■ actualmente tomando un medicamento recetado. Los antiácidos pueden interactuar con ciertos medicamentos recetados. ■ tiene una enfermedad renal  <b>Cuando administre este producto</b> ■ no tomar más de 10 tabletas en 24 horas ■ si está embarazada, no tomar más de 6 tabletas en 24 horas Mantenga este medicamento fuera de alcance de los niños.	<b>Propiedad</b> Antiácido  <b>Información del medicamento (continuación)</b> <b>Instrucciones</b> ■ no tomar para síntomas que persisten por más de dos semanas a menos que lo recomiende un doctor  <b>Adultos y niños:</b> Masticar de 2 a 4 tabletas a medida que aparecen los síntomas o como lo indique un médico.  <b>Niños menores de 12 años:</b> consultar al médico  <b>Otra información</b> ■ contenido de calcio por comprimido: 300 mg ■ 5 calorías por tableta ■ almacenamiento: consérvase a temperatura ambiente 59-86°F (15-30°C) en un área seca ■ empaquetado con sellado evidente ■ no use ningún paquete abierto o roto  <b>Ingredientes inactivos</b> saborizantes surtidos, dextrosa, azul FD y C #1 lacas de aluminio, rojo FD y C #40 lacas de aluminio, estearato de magnesio, maltodextrina  <b>¿Tiene alguna pregunta o comentario?</b> Llame al 1-800-634-7680	 <p><b>Medique®</b> <b>Ban-acid®</b> Maximum Strength 750 Calcium Rich Extra Strength Antacid Antiácido extra fuerte y /ricas en calcio</p> <p>Reorder #28533 Manufactured for Medique Products Fort Myers, FL 33967 1-800-634-7680 www.mediqueproducts.com</p> <p>47682 28533 3</p>
<b>Drug Facts</b> <b>Active ingredient (in each tablet) Purpose</b> Calcium Carbonate USP 750 mg Antacid  <b>Uses</b> relieves ■ acid indigestion ■ sour stomach ■ heartburn ■ upset stomach associated with these symptoms  <b>Warnings</b> Do not use ■ the maximum dosage of this product for more than 2 weeks except under the advice and supervision of a doctor. Ask a doctor or pharmacist before use if you ■ are taking a prescription drug. Antacids may interact with certain prescription drugs. ■ have kidney disease  <b>When using this product</b> ■ do not take more than 10 tablets in 24 hours ■ if pregnant, do not take more than 6 tablets in 24 hours Keep out of the reach of children.	<b>Drug Facts (continued)</b>  <b>Directions</b> ■ do not take for symptoms that persist for more than 2 weeks unless advised by a doctor  <b>Adults and children:</b> Chew 2 to 4 tablets as symptoms occur, or as directed by a doctor  <b>Children under 12 years:</b> Ask a doctor  <b>Other information</b> ■ calcium content per tablet: 300 mg ■ 5 calories per tablet ■ store at room temperature 59-86°F (15-30°C) ■ tamper-evident sealed packets ■ do not use any opened or torn packets  <b>Inactive ingredients</b> assorted flavors, dextrose, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, magnesium stearate, maltodextrin  <b>Questions or comments</b> Call 1-800-634-7680						
<b>Información del Medicamento</b> <b>Ingrediente activo (en cada tableta)</b> Carbonato de Calcio 750 mg  <b>Usos</b> alivia ■ indigestión ácida ■ acidez estomacal ■ agruras ■ malestar estomacal (debido a estos síntomas)  <b>Advertencias</b> No consuma ■ la dosis máxima de este producto por más de 2 semanas, excepto bajo indicaciones y supervisión médica.  <b>Consulte con el doctor o un farmacéutico antes de consumir este medicamento si usted está</b> ■ actualmente tomando un medicamento recetado. Los antiácidos pueden interactuar con ciertos medicamentos recetados. ■ tiene una enfermedad renal  <b>Cuando administre este producto</b> ■ no tomar más de 10 tabletas en 24 horas ■ si está embarazada, no tomar más de 6 tabletas en 24 horas Mantenga este medicamento fuera de alcance de los niños.	<b>Propiedad</b> Antiácido  <b>Información del medicamento (continuación)</b> <b>Instrucciones</b> ■ no tomar para síntomas que persisten por más de dos semanas a menos que lo recomiende un doctor  <b>Adultos y niños:</b> Masticar de 2 a 4 tabletas a medida que aparecen los síntomas o como lo indique un médico.  <b>Niños menores de 12 años:</b> consultar al médico  <b>Otra información</b> ■ contenido de calcio por comprimido: 300 mg ■ 5 calorías por tableta ■ almacenamiento: consérvase a temperatura ambiente 59-86°F (15-30°C) en un área seca ■ empaquetado con sellado evidente ■ no use ningún paquete abierto o roto  <b>Ingredientes inactivos</b> saborizantes surtidos, dextrosa, azul FD y C #1 lacas de aluminio, rojo FD y C #40 lacas de aluminio, estearato de magnesio, maltodextrina  <b>¿Tiene alguna pregunta o comentario?</b> Llame al 1-800-634-7680						
<p>Retain carton for complete product information</p>	<p>Conserve la caja para obtener la información completa del producto</p>						

# MEDIQUE BAN-ACID MAXIMUM STRENGTH 750

calcium carbonate tablet, chewable

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-285
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK) (CALCIUM CATION - UNII:2M83C4R6ZB)	CALCIUM CARBONATE	750 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>DEXTROSE</b> (UNII: IY9XDZ35W2)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	

## Product Characteristics

<b>Color</b>	red (RED) , purple (PURPLE) , gray (GRAY)	<b>Score</b>	no score
<b>Shape</b>	ROUND (ROUND)	<b>Size</b>	15mm
<b>Flavor</b>	STRAWBERRY (Blueberry Blackberry)	<b>Imprint Code</b>	RH;031
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-285-33	50 in 1 BOX	09/29/2017	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-285-99	2 in 1 PACKET; Type 0: Not a Combination Product	07/01/2014	
3	NDC:47682-285-36	75 in 1 BOX	04/01/2019	
3		2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC Monograph Drug M001

07/01/2014

**Labeler** - Unifirst First Aid Corporation (832947092)

Revised: 6/2024

Unifirst First Aid Corporation