

**ACETAMINOPHEN- acetaminophen tablet**  
**Rite Aid Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Rite Aid Corporation Acetaminophen Drug Facts**

**Active ingredient (in each caplet)**

Acetaminophen 500 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

liver disease

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"> <li>• take 2 caplets every 6 hours while symptoms last</li> <li>• do not take more than 6 caplets in 24 hours, unless directed by a doctor</li> <li>• do not use for more than 10 days unless directed by a doctor</li> </ul>
children under 12 years	ask a doctor

**Other information**

- store at 20-25°C (68-77°F)

## Inactive ingredients

carnauba wax, corn starch\*, croscarmellose sodium\*, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate\*, stearic acid

\*may contain one or more of these ingredients

## Questions or comments?

**1-800-719-9260**

## Principal Display Panel

Compare to the active ingredient of Extra Strength Tylenol® Caplets

FREE FROM

GLUTEN FREE

CAFFEINE FREE

EXTRA STRENGTH

ACETAMINOPHEN

ACETAMINOPHEN CAPLETS, 500 mg

ACTUAL SIZE

PAIN RELIEVER/FEVER REDUCER

For adults

225 CAPLETS





**Drug Facts**

**Active ingredient (In each caplet)**      **Purpose**  
Acetaminophen 500 mg.....Pain reliever/fever reducer

**Uses** ■ temporarily relieves minor aches and pains due to:  
■ the common cold    ■ headache    ■ backache  
■ minor pain of arthritis    ■ toothache    ■ muscular aches  
■ premenstrual and menstrual cramps  
■ temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - new symptoms occur    ■ redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

■ do not take more than directed (see overdose warning)

adults and children 12 years and over	■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

ACTUAL SIZE





# ACETAMINOPHEN

acetaminophen tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-4484
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

## Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3S)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

## Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	L484
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-4484-1	1 in 1 CARTON	09/08/2020	
1		225 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11822-4484-2	1000 in 1 BOTTLE; Type 0: Not a Combination Product	09/04/2020	
3	NDC:11822-4484-3	1 in 1 CARTON	07/01/2022	
3		24 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:11822-4484-4	1 in 1 CARTON	07/11/2022	
4		50 in 1 BOTTLE; Type 0: Not a Combination		

<b>4</b>		Product		
<b>5</b>	NDC:11822-4484-5	1 in 1 CARTON	07/11/2022	
<b>5</b>		100 in 1 BOTTLE; Type 0: Not a Combination Product		
<b>6</b>	NDC:11822-4484-6	500 in 1 BOTTLE; Type 0: Not a Combination Product	07/11/2022	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	09/04/2020	

**Labeler** - Rite Aid Corporation (014578892)

Revised: 7/2022

Rite Aid Corporation