

**BACKPRIN- acetaminophen, caffeine, salicylamide tablet**  
**HART Health**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**BACKPRIN**

**Active Ingredients (in each tablet):**

Acetaminophen 250 mg

Caffeine 50 mg

Magnesium Salicylate (NSAID\*) 290 mg (\*nonsteroidal anti-inflammatory drug)

**Purpose:**

Pain Reliever/Fever Reducer

Pain Reliever Aid

Pain Reliever/Fever Reducer

**Uses:**

Temporarily relieves minor back pain

**Warnings:**

**Reye's syndrome:** Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with salicylates.

**Allergy alert:** Acetaminophen and Magnesium Salicylate may cause severe allergic reaction which may include hives, skin reddening, blisters, rash, facial swelling, asthma (wheezing), or shock.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- take more or for a longer time than directed
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- take other drugs containing prescription or non-prescription NSAID (aspirin, ibuprofen, naproxen, others)

**Do not use:**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer

- with any other drug containing acetaminophen (prescription or non-prescription)

**Ask a doctor before use if**

- you have asthma, high blood pressure, heart disease, liver cirrhosis, or kidney disease
- stomach bleeding warning applies to you
- you are taking a diuretic
- you have a history of stomach problems, such as heartburn

**Ask a doctor or pharmacist before use if you**

- take a blood thinning (anticoagulant) or steroid drug
- take any drug for diabetes, gout or arthritis
- do not know if other drugs you are taking contain acetaminophen
- take any other drug containing an NSAID

**Stop use and ask a doctor if:**

- an allergic reaction occurs, seek medical help right away
- fever worsens or lasts for more than 3 days
- redness or swelling is present
- new symptoms occur
- pain worsens or lasts for more than 10 days
- ringing in the ears or a loss of hearing occurs
- you have any of the following signs of stomach bleeding: feel faint, vomit blood, have bloods or black stools, or stomach pain that does not get better

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use salicylates during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. In case of overdose, get medical help or contact a Poison Control Center right away. 1-800-222-1222

**Directions:**

**Do not take more than directed**

Adults and children 12 years of age and over:

- take 2 tablets with water every 4 hours while symptoms persist.
- do not take more than 8 tablets in 24 hours unless directed by a doctor

Children under 12 years of age: ask a doctor

**Inactive Ingredients:**

Hypromellose, Maltodextrin, Microcrystalline Cellulose, Povidone, Polyethylene Glycol, Starch,

# Stearic Acid, Titanium Dioxide

50 count  
 PMS 259C purple  
 PMS 7404C yellow  
 small medication box dieline  
 10.8cm x 6cm x 9cm

NDC 50332-0120-4      Item 5526

**COMPARE TO**  
**Doans®**  
 Active Ingredient

**Backprin®**  
**Back Pain Reliever**

Acetaminophen  
 Caffeine  
 Magnesium Salicylate (NSAID)

Reduces Back Pain  
 50 packets of 2 tablets  
 100 total tablets

HART-Health

Manufactured for:  
**HART-Health**  
 6000  
 50332-0120-4  
 9448-19

\*This product is not manufactured or distributed by Foster-Milburn Company, owner of the registered trademark Doans® products.

Item 5526

**COMPARE TO**  
**Doans®**  
 Active Ingredient

**Backprin®**  
**Back Pain Reliever**  
*Analgesico de Espalda*

Acetaminophen  
 Caffeine  
 Magnesium Salicylate (NSAID)

Reduces Back Pain  
*Reduce el Dolor de Espalda*  
 50 packets of 2 tablets  
 100 total tablets

HART-Health

Item 5526

**Backprin®**  
**Back Pain Reliever**  
*Analgesico de Espalda*

Acetaminophen  
 Caffeine  
 Magnesium Salicylate (NSAID)

Reduces Back Pain  
*Reduce el Dolor de Espalda*  
 50 packets of 2 tablets  
 100 total tablets

HART-Health

**Drug Facts** (continued)

**Backprin®**

**Active Ingredients** (in each tablet): Acetaminophen 325 mg, Caffeine 50 mg, Magnesium Salicylate (NSAID) 250 mg (Nonsteroidal anti-inflammatory drug).

**Purpose:** Pain Reliever/Fever Reducer

**Warnings:** Temporarily relieves minor back pain.

**Directions:** Do not take more than directed. Adults and children 12 years of age and over: take 2 tablets with water every 4 hours with symptoms present. Do not take more than 8 tablets in 24 hours unless directed by a doctor. Children under 12 years of age: ask a doctor.

**Other Information:** Tablets enclosed in a sealed packet stored at 20° to 25° (68° to 77°). Do not use if packet is torn, out, or open or if not sealed and tampered.

**Inactive Ingredients:** Hypromellose, Methylcellulose, Microcrystalline Cellulose, Povidone, Polyethylene Glycol, Starch, Stearic Acid, Titanium Dioxide.


Manufactured for:  
**HART-Health**  
 Seattle, WA 98124    www.harthealth.com

**KEY**

- cut line —————
- inward fold line - - - - -
- outward fold line - . . . . .
- perf —————
- slit —————
- outside glue
- inside glue



50 count  
PMS 259C purple  
PMS 7404C yellow  
small medication box dieline  
10.8cm x 6cm x 9cm



- KEY**
- cut line ————
  - inward fold line - - - - -
  - outward fold line ·····
  - perf ————
  - slit ————
  - outside glue
  - inside glue

## BACKPRIN

acetaminophen, caffeine, salicylamide tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50 332-0 116
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	50 mg
MAGNESIUM SALICYLATE (UNII: 41728CY7UX) (SALICYLIC ACID - UNII:O414PZ4LPZ)	MAGNESIUM SALICYLATE	290 mg

### Inactive Ingredients

Ingredient Name	Strength
MALTO DEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	

<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	

<b>Product Characteristics</b>			
<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50332-0116-4	50 in 1 BOX, UNIT-DOSE	07/14/2000	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:50332-0116-2	125 in 1 BOX, UNIT-DOSE	07/14/2000	
2		2 in 1 PACKET; Type 0: Not a Combination Product		

<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	07/14/2000	

**Labeler** - HART Health (069560969)