

**ANTACID AND GASRELIEF- aluminum hydroxide, magnesium hydroxide,
dimethicone suspension
GOODSENSE**

GS ANTACID ORIGINAL

Active ingredients (in each 10 mL dose)

Aluminum hydroxide 400 mg (equivalent to dried gel, USP)
Magnesium hydroxide 400 mg
Simethicone 40 mg

Purposes

Antacid
Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years of age and older: take 10 mL to 20 mL (1 to 2 doses) between meals as needed, at bedtime, or as directed by a doctor
- do not exceed 120 mL (12 doses) in a 24 hour period or use the maximum dosage for more than 2 weeks

- children under 12 years of age: ask a doctor

Other information

- each 10 mL dose contains: **magnesium 165 mg, sodium 6 mg**
- store at room temperature tightly closed protect from freezing
- keep tightly closed

Inactive ingredients

benzyl alcohol, butylparaben, flavor (contains alcohol), hydroxyethylcellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments?

1-800-540-3765

package label

GOODSENSE®

NDC 50804-629-12

Regular Strength

Antacid & Gas Relief

Alumina, Magnesia, and Simethicone
Oral Suspension USP

Fast Relief of:

Acid Indigestion

Heartburn

Sour Stomach

Pressure & Bloating

Original
Flavored Liquid

Compare to active ingredients of
MYLANTA Regular Strength*

12 FL OZ (355mL)

274-05112-1 REV GC1023

Drug Facts

TAMPER-EVIDENT: Do not use if breakaway band on bottle cap is missing or broken.

Active ingredients (in each 10 mL dose)	Purpose
Aluminum hydroxide (equivalent to dried gel, USP) 400 mg	Antacid
Magnesium hydroxide 400 mg	Antacid
Simethicone 40 mg	Antigas

Uses relieves • heartburn • sour stomach
• acid indigestion • the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years of age and older: take 10 mL to 20 mL (1 to 2 doses) between meals as needed, at bedtime, or as directed by a doctor
- do not exceed 120 mL (12 doses) in a 24 hour period or use the maximum dosage for more than 2 weeks
- children under 12 years of age: ask a doctor

Other information

- each 10 mL dose contains: magnesium 165 mg, sodium 6 mg
- store at room temperature tightly closed • protect from freezing

Inactive ingredients benzyl alcohol, butylparaben, flavor (contains alcohol), hydroxyethylcellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

*This product is not manufactured or distributed by the owner of the registered trademark MYLANTA®. 974-05112-1 REV GC0324

Distributed by:
Perrigo Direct
Peachtree City, GA 30269
www.PerrigoDirect.com
1-800-540-3765
GoodSense® is a registered trademark of L. Perrigo Company



ANTACID AND GASRELIEF

aluminum hydroxide, magnesium hydroxide, dimethicone suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50804-629
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength

ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	400 mg in 10 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	400 mg in 10 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	40 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50804-629-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	07/01/2020	

Labeler - GOODSENSE (076059836)

Registrant - GCP Laboratories (965480861)

Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(50804-629)