

AIRMIT ACE- dimenhydrinate tablet
Sato Pharmaceutical Co., Ltd.

Airmit Ace

Active ingredient (in each tablet)
Dimenhydrinate 25 mg

Purpose
Antiemetic

Uses
For the prevention and treatment of the following symptoms associated with motion sickness
■nausea ■vomiting ■dizziness

Warnings

Do not use

- for children under 2 years of age unless directed by a doctor
- this product if you are taking sedatives or tranquilizers, without first consulting your doctor

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

When using this product

- may cause marked drowsiness: alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use.

Keep our of reach of children. In case of over dose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 12 years of age and over - oral dosage is 2 to 4 tablets every 4 to 6 hours, not to exceed 16 tablets in 24 hours, or as directed by a doctor

children 6 to under 12 years of age - oral dosage is 1 to 2 tablets every 6 to 8 hours, not to exceed 6 tablets in 24 hours, or as directed by a doctor

children 2 to under 6 years of age - oral dosage is 1 tablet every 6 to 8 hours, not to

exceed 3 tablets in 24 hours, or as directed by a doctor

children under 2 years of age - ask a doctor

Other information

■each tablet contains: calcium 5mg

Inactive ingredients Dibasic calcium phosphate, FD and C Yellow No. 6, hydroxypropyl cellulose, hydroxypropyl starch, lactose, magnesium stearate, mannitol, microcrystalline cellulose, povidone, sugar.



Drug Facts

Active ingredient (in each tablet) **Purpose**
Dimenhydrinate 25 mg Antiemetic

Uses For the prevention and treatment of the following symptoms associated with motion sickness.
■ nausea ■ vomiting ■ dizziness

Warnings

Do not use ■ for children under 2 years of age unless directed by a doctor ■ this product if you are taking sedatives or tranquilizers, without first consulting your doctor

Ask a doctor before use if you have

■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma
■ difficulty in urination due to enlargement of the prostate gland

When using this product

■ may cause marked drowsiness: alcohol, sedatives, and tranquilizers may increase the drowsiness effect
■ avoid alcoholic beverages ■ use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 12 years of age and over	oral dosage is 2 to 4 tablets every 4 to 6 hours, not to exceed 16 tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years of age	oral dosage is 1 to 2 tablets every 6 or 8 hours, not to exceed 6 tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years of age	oral dosage is 1 tablet every 6 to 8 hours, not to exceed 3 tablets in 24 hours, or as directed by a doctor
children under 2 years of age	ask a doctor

Other information

■ each tablet contains : calcium 5mg

Inactive ingredients Dbasic calcium phosphate, FD&C Yellow No.6, hydroxypropyl cellulose, hydroxypropyl starch, lactose, magnesium stearate, mannitol, microcrystalline cellulose, povidone, sugar.

EXPDATE 使用期限
LOT 製造ロット
USC

PUSH

USC

用法・用量 成人及び12才以上の小児 1回2～4錠を4～6時間おきに服用してください。 1回1～2錠を6～8時間おきに服用してください。 2歳未満の小児は、医師の指示に従ってください。	2才未満の小児 医師に相談ください。 また、医師に相談ください。	2～5才の小児 1回1錠を6～8時間おきに服用してください。24時間以内3錠をすぎず服用しないでください。	6～11才の小児 1回1～2錠を6～8時間おきに服用してください。24時間以内3錠をすぎず服用しないでください。
その他の成分 ■ 1錠中：カルシウム 5mg相当	その他の成分 ■ 1錠中：カルシウム 5mg相当		

ANTHEMATIC
(CAUTION: SICKNESS REMEDY)
airmit Ace

2才のお子様からのめます
アイミットエース
嘔吐剤 (薬物酔いに)
May be given to children 2 years of age or older
ANTHEMATIC (MOTION SICKNESS REMEDY)
airmit Ace
SATO
NDC 49873-802-01
6 Tablets (6錠)
Icons: airplane, car, boat

Manufactured by
SATO PHARMACEUTICAL CO., LTD.
1-5-27 MOTOKAKASAKA MINATO-KU TOKYO, JAPAN
Reports of serious side effects associated with use of this product can be sent to:
SATO PHARMACEUTICAL INC.
20501 East Street, Suite 3, Torrance, CA 90503



* Tablets have been individually blister sealed and are enclosed in this carton sealed by adhesive. If seals are broken or missing, do not use.

* 錠剤は個々にブリスター包装されているので、カートンに入られては密封されています。もしシールが破損していたり、無い場合は服用しないでください。

医薬品表示 アイミットエース 25mg 剤名 有効成分・分量 効能 乗船酔いによる下腹部の予防及び緩和 効心 乗船酔い 副作用 嘔吐 使用上の注意 次の人は本剤を服用しないでください。医師の指示に従って服用してください。 ■ 本剤を服用して3～5分後に嘔吐が繰り返す場合、本剤を服用しないでください。	用法・用量 成人及び12才以上の小児 1回2～4錠を4～6時間おきに服用してください。 1回1～2錠を6～8時間おきに服用してください。 2歳未満の小児は、医師の指示に従ってください。	2才未満の小児 医師に相談ください。 また、医師に相談ください。	2～5才の小児 1回1錠を6～8時間おきに服用してください。24時間以内3錠をすぎず服用しないでください。	6～11才の小児 1回1～2錠を6～8時間おきに服用してください。24時間以内3錠をすぎず服用しないでください。
成分 ■ 1錠中：カルシウム 5mg相当	その他の成分 ■ 1錠中：カルシウム 5mg相当			

AIRMIT ACE
dimenhydrinate tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49873-802

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIMENHYDRINATE (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIMENHYDRINATE	25 mg

Inactive Ingredients

Ingredient Name	Strength
CALCIUM PHOSPHATE, DIBASIC, ANHYDROUS (UNII: L11K75P92J)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYDROXYPROPYL CORN STARCH (5% SUBSTITUTION BY WEIGHT) (UNII: 9M44R3409A)	
HYDROXYPROPYL CELLULOSE (UNII: RFW2ET671P)	
LACTOSE (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POVIDONE (UNII: FZ989GH94E)	
SUCROSE (UNII: C151H8M554)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	SATO;7
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49873-802-01	1 in 1 CARTON	03/11/2005	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	03/11/2005	

Labeler - Sato Pharmaceutical Co., Ltd. (690575642)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

Sato Pharmaceutical Co., Ltd.	715699133	manufacture(49873-802) , label(49873-802) , pack(49873-802)
-------------------------------	-----------	---

Revised: 12/2023

Sato Pharmaceutical Co., Ltd.