

**TUSSIN MULTI SYMPTOM COLD CF ADULT- dextromethorphan hbr,
guaifenesin, phenylephrine liquid
Major Pharmaceuticals**

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr 20 mg

Guaifenesin 200 mg

Phenylephrine HCl 10 mg

Purposes

Cough suppressant

Expectorant

Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
 - nasal congestion
 - cough due to minor throat and bronchial irritation

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- diabetes
- thyroid
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are

taking any other oral nasal decongestant or stimulant.

**When using this product,
do not use more than directed.**

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age
- adult and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use

Other information

- store between 20-25°C (68°-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

compare to the active ingredient in Robitussin® Multi-Symptom Cold CF†

robafen

CF multi-symptom cold

Dextromethorphan HBr, 20 mg

cough suppressant

Guaifenesin, 200 mg

expectorant

Phenylephrine HCl, 10 mg

nasal decongestant

Relieves:

- Cough
- Mucus
- Nasal Congestion

for ages 12 & over

non-drowsy

alcohol-free

FOR ADULTS

For Ages 12 Years and Over

Alcohol-Free

FL OZ (mL)

Dosing Cup Included

†This product is not manufactured or distributed by Haleon group of companies, distributors of Robitussin® Multi-Symptom Cold CF.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Distributed by:

MAJOR® PHARMACEUTICALS

Indianapolis, IN 46268

Livonia, MI 48152

Package Label

Drug Facts (continued)

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age
- adults and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use

Other information

- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

Questions or comments?
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by Haleon group of companies, distributor of Robitussin® Multi-Symptom Cold CF.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

PARENTS:
Learn about teen medicine abuse.
www.StopMedicineAbuse.org



Dosing Cup Included

Distributed by:
MAJOR® PHARMACEUTICALS
Indianapolis, IN 46268
(800) 616-2471
www.major-rugby.com

Rev. 09/24 M-58 Re-order No. 700529

MAJOR® NDC 0904-6537-20

compare to the active ingredients in Robitussin® Multi-Symptom Cold CF†

robafen CF multi-symptom cold

dextromethorphan HBr 20 mg
cough suppressant
guaifenesin 200 mg
expectorant
phenylephrine HCl 10 mg
nasal decongestant

relieves:

- cough
- mucus
- nasal congestion

for ages 12 & over
non-drowsy
alcohol-free

4 fl oz (118 mL)

NDC 0904-6537-20

compare to the active ingredients in Robitussin® Multi-Symptom Cold CF†

robafen CF multi-symptom cold

dextromethorphan HBr 20 mg
cough suppressant
guaifenesin 200 mg
expectorant
phenylephrine HCl 10 mg
nasal decongestant

relieves:

- cough
- mucus
- nasal congestion

for ages 12 & over
non-drowsy
alcohol-free

4 fl oz (118 mL)

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr 20 mg.....Cough suppressant
Guaifenesin 200 mg.....Expectorant
Phenylephrine HCl 10 mg.....Nasal decongestant

Purposes

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold
- nasal congestion
- cough due to minor throat and bronchial irritation

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.



3 09046 53720 5

PLD-E322C F0009119

Lot No.:
Exp. Date:

MAJOR Multi-Symptom Cold

TUSSIN MULTI SYMPTOM COLD CF ADULT

dextromethorphan hbr, guaifenesin, phenylephrine liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-6537
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE -	PHENYLEPHRINE	10 mg

UNII:1WS297W6MV)

HYDROCHLORIDE

in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
LACTIC ACID (UNII: 33X04XA5AT)	
MENTHOL (UNII: L7T10EIP3A)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-6537-20	1 in 1 BOX	11/30/2015	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/30/2015	

Labeler - Major Pharmaceuticals (191427277)

Revised: 10/2024

Major Pharmaceuticals