

**TUSSIN MULTI SYMPTOM COLD CF ADULT- dextromethorphan hbr,
guaifenesin, phenylephrine liquid
Major Pharmaceuticals**

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr 20 mg

Guaifenesin 200 mg

Phenylephrine HCl 10 mg

Purposes

Cough suppressant

Expectorant

Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
 - nasal congestion
 - cough due to minor throat and bronchial irritation

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- diabetes
- thyroid
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are

taking any other oral nasal decongestant or stimulant.

**When using this product,
do not use more than directed.**

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age
- adult and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use

Other information

- store between 20-25°C (68°-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal Display Panel

ROBAFEN

CF MULTI-SYMPTOM COLD

Dextromethorphan HBr, 20 mg / COUGH SUPPRESSANT

Guaifenesin, 200 mg / EXPECTORANT

Phenylephrine HCl, 10 mg / NASAL DECONGESTANT

PEAK COLD

Relieves:

- Cough
- Mucus
- Nasal Congestion

Non-Drowsy

COMPARE TO the active ingredients in ROBITUSSIN® PEAK COLD MULTI-SYMPTOM COLD CF*

FOR ADULTS

For Ages 12 Years and Over

Alcohol-Free

FL OZ (mL)

Dosing Cup Included

*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributors of Robitussin® Peak Cold Multi-Symptom Cold CF.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Distributed by:

MAJOR PHARMACEUTICALS

17177 N Laurel Park Drive, Suite 233

Livonia, MI 48152

Package Label

Drug Facts (continued)

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age
- adults and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use

Other information

- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

Questions or comments?

Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Robitussin® Peak Cold Multi-Symptom Cold CF.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

PARENTS:
Learn about how medicine abuse
www.StopMedicineAbuse.org



Dosing Cup Included

Distributed by:
Major® Pharmaceuticals
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152
Re-Order No. 700529
Rev. 01/17 M-58

NDC 0904-6537-20



ROBAFEN

CF MULTI-SYMPOM COLD

Dextromethorphan HBr, 20 mg /
COUGH SUPPRESSANT
Guaifenesin, 200 mg / EXPECTORANT
Phenylephrine HCl, 10 mg /
NASAL DECONGESTANT

PEAK COLD

Relieves:

- Cough
- Mucus
- Nasal Congestion

Non-Drowsy

COMPARE TO the active ingredients
in **ROBITUSSIN® PEAK COLD
MULTI-SYMPOM COLD CF***

FOR ADULTS
For Ages 12 Years and Over
Alcohol-Free

4 FL OZ (118 mL)

NDC 0904-6537-20



ROBAFEN

CF MULTI-SYMPOM COLD

Dextromethorphan HBr, 20 mg /
COUGH SUPPRESSANT
Guaifenesin, 200 mg / EXPECTORANT
Phenylephrine HCl, 10 mg /
NASAL DECONGESTANT

PEAK COLD

Relieves:

- Cough
- Mucus
- Nasal Congestion

Non-Drowsy

COMPARE TO the active ingredients
in **ROBITUSSIN® PEAK COLD
MULTI-SYMPOM COLD CF***

FOR ADULTS
For Ages 12 Years and Over
Alcohol-Free

4 FL OZ (118 mL)

Drug Facts

Active ingredients (in each 10 mL)

Active ingredients (in each 10 mL)	Purposes
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold
 - nasal congestion
 - cough due to minor throat and bronchial irritation

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.



3 09046 53720 5

PLD-0322C FC004098

Lot No:
Exp. Date:

MAJOR Multi-Symptom Cold

TUSSIN MULTI SYMPTOM COLD CF ADULT

dextromethorphan hbr, guaifenesin, phenylephrine liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-6537
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL

PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)		PHENYLEPHRINE HYDROCHLORIDE	10 mg in 10 mL	
Inactive Ingredients				
Ingredient Name			Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
GLYCERIN (UNII: PDC6A3C0OX)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
LACTIC ACID (UNII: 33X04XA5AT)				
MENTHOL (UNII: L7T10EIP3A)				
SORBITOL (UNII: 506T60A25R)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-6537-20	1 in 1 BOX	11/30/2015	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M012	11/30/2015		

Labeler - Major Pharmaceuticals (191427277)