

## **ATROPINE SULFATE- atropine sulfate injection, solution**

**Henry Schein, Inc.**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).*

-----

### **HIGHLIGHTS OF PRESCRIBING INFORMATION**

**Atropine Sulfate 0.1 mg/mL Injection, USP 10 mL PreFilled Syringe**

### **HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use ATROPINE SULFATE INJECTION safely and effectively. See full prescribing information for ATROPINE SULFATE INJECTION. ATROPINE SULFATE INJECTION, for intravenous use**  
**Initial U.S. Approval: 1960**

### **----- INDICATIONS AND USAGE -----**

(1)

### **----- DOSAGE AND ADMINISTRATION -----**

(2)

### **----- DOSAGE FORMS AND STRENGTHS -----**

0.1 mg/mL injection in LifeShield™ Abboject™ Glass Syringe (3) (3)

### **----- CONTRAINDICATIONS -----**

None. (4) (4)

### **----- WARNINGS AND PRECAUTIONS -----**

Tachycardia (5.1) (5) (5)

Glaucoma (5.2) (5) (5)

Pyloric obstruction (5.3) (5) (5)

Worsening urinary retention (5.4) (5) (5)

Viscid bronchial plugs (5.5) (5) (5)

### **----- DRUG INTERACTIONS -----**

Mexiletine: Decreases rate of mexiletine absorption. (7.1) (7) (6)

Revised: 3/2018 (6)

**Revised: 10/2023**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

**Indications and Usage**

**Dosage and Administration**

**Dosage Forms and Strengths**

**Contraindications**

**Warnings and Precautions**

**Drug Interactions**

**Use In Specific Populations**

**Overdosage**

**Description**

**Clinical Pharmacology**

**Nonclinical Toxicology**

**How Supplied**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## FULL PRESCRIBING INFORMATION

### Indications and Usage

Atropine Sulfate Injection, USP, is indicated for temporary blockade of severe or life threatening muscarinic effects, e.g., as an antisialagogue, an antivagal agent, an antidote for organophosphorus or muscarinic mushroom poisoning, and to treat bradycardiac arrest.

### Dosage and Administration

#### 2.1 General Administration

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Do not administer unless solution is clear and seal is intact. Discard unused portion.

For intravenous administration.

Titrate based on heart rate, PR interval, blood pressure and symptoms

#### 2.2 Adult Dosage

**Table 1: Recommended Dosage**

Use	Dose (adults)	Repeat
Antisialagogue or other antivagal	0.5 to 1 mg	1–2 hours
Organophosphorus or muscarinic mushroom poisoning	2 to 3 mg	20–30 minutes
Bradycardiac arrest	1 mg	3–5 minutes; 3 mg maximum total dose

#### 2.3 Pediatric Dosage

Dosing in pediatric populations has not been well studied. Usual initial dose is 0.01 to 0.03 mg/kg.

### Dosage Forms and Strengths

Injection: 0.1 mg/mL in LifeShield™ Abboject™ Glass Syringes

## **Contraindications**

None.

## **Warnings and Precautions**

### **5.1 Tachycardia**

When the recurrent use of atropine is essential in patients with coronary artery disease, the total dose should be restricted to 2 to 3 mg (maximum 0.03 to 0.04 mg/kg) to avoid the detrimental effects of atropine-induced tachycardia on myocardial oxygen demand.

### **5.2 Acute Glaucoma**

Atropine may precipitate acute glaucoma.

### **5.3 Pyloric Obstruction**

Atropine may convert partial organic pyloric stenosis into complete obstruction.

### **5.4 Complete Urinary Retention**

Atropine may lead to complete urinary retention in patients with prostatic hypertrophy.

### **5.5 Viscid Plugs**

Atropine may cause inspissation of bronchial secretions and formation of viscid plugs in patients with chronic lung disease.

## **Drug Interactions**

Atropine Sulfate Injection decreased the rate of mexiletine absorption without altering the relative oral bioavailability; this delay in mexiletine absorption was reversed by the combination of atropine and intravenous metoclopramide during pretreatment for anesthesia.

## **Use In Specific Populations**

### **8.1 Pregnancy**

Animal reproduction studies have not been conducted with atropine. It also is not known whether atropine can cause fetal harm when given to a pregnant woman or can affect reproduction capacity.

### **8.3 Nursing Mothers**

Trace amounts of atropine was found in breast milk. The clinical impact of this is not known.

### **8.4 Pediatric Use**

Recommendations for use in pediatric patients are not based on clinical trials.

## 8.5 Geriatric Use

An evaluation of current literature revealed no clinical experience identifying differences in response between elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

## Overdosage

Excessive dosing may cause palpitation, dilated pupils, difficulty in swallowing, hot dry skin, thirst, dizziness, restlessness, tremor, fatigue and ataxia. Toxic doses lead to restlessness and excitement, hallucinations, delirium and coma. Depression and circulatory collapse occur only with severe intoxication. In such cases, blood pressure declines and death due to respiratory failure may ensue following paralysis and coma.

The fatal adult dose of atropine is not known. In pediatric populations, 10 mg or less may be fatal.

In the event of toxic overdosage, a short acting barbiturate or diazepam may be given as needed to control marked excitement and convulsions. Large doses for sedation should be avoided because central depressant action may coincide with the depression occurring late in atropine poisoning. Central stimulants are not recommended.

Physostigmine, given as an atropine antidote by slow intravenous injection of 1 to 4 mg (0.5 to 1 mg in pediatric populations), rapidly abolishes delirium and coma caused by large doses of atropine. Since physostigmine is rapidly destroyed, the patient may again lapse into coma after one to two hours, and repeated doses may be required.

Artificial respiration with oxygen may be necessary. Ice bags and alcohol sponges help to reduce fever, especially in pediatric populations.

Atropine is not removed by dialysis.

## Description

Atropine Sulfate Injection, USP is a sterile, nonpyrogenic isotonic solution of atropine sulfate monohydrate in water for injection with sodium chloride sufficient to render the solution isotonic. It is administered parenterally by intravenous injection.


Each milliliter (mL) contains 0.1 mg of atropine sulfate monohydrate equivalent to 0.083 mg of atropine, and sodium chloride, 9 mg. May contain sodium hydroxide and/or sulfuric acid for pH adjustment 0.308 mOsmol/mL (calc.) pH 3.0 to 6.5.

Sodium chloride added to render the solution isotonic for injection of the active ingredient is present in amounts insufficient to affect serum electrolyte balance of sodium (Na<sup>+</sup>) and chloride (Cl<sup>-</sup>) ions.

The solution contains no bacteriostat, antimicrobial agent or added buffer (except for pH adjustment) and is intended for use only as a single-dose injection. When smaller doses are required the unused portion should be discarded.

Atropine Sulfate, USP is chemically designed 1aH, 5aH-Tropan-3-a-ol(±)-tropate (ester), sulfate (2:1) (salt) monohydrate, (C<sub>17</sub>H<sub>23</sub>NO<sub>3</sub>)<sub>2</sub>•H<sub>2</sub>SO<sub>4</sub>•H<sub>2</sub>O, colorless crystals or

white crystalline powder very soluble in water. It has the following structural formula:



Atropine, a naturally occurring belladonna alkaloid, is a racemic mixture of equal parts of d- and l-hyocyamine, whose activity is due almost entirely to the levo isomer of the drug.

Sodium Chloride, USP is chemically design NaCl, a white crystalline powder freely soluble in water.

## Clinical Pharmacology

### 12.1 Mechanism of Action

Atropine is an antimuscarinic agent since it antagonizes the muscarine-like actions of acetylcholine and other choline esters.

Atropine inhibits the muscarinic actions of acetylcholine on structures innervated by postganglionic cholinergic nerves, and on smooth muscles which respond to endogenous acetylcholine but are not so innervated. As with other antimuscarinic agents, the major action of atropine is a competitive or surmountable antagonism which can be overcome by increasing the concentration of acetylcholine at receptor sites of the effector organ (e.g., by using anticholinesterase agents which inhibit the enzymatic destruction of acetylcholine). The receptors antagonized by atropine are the peripheral structures that are stimulated or inhibited by muscarine (i.e., exocrine glands and smooth and cardiac muscle). Responses to postganglionic cholinergic nerve stimulation also may be inhibited by atropine but this occurs less readily than with responses to injected (exogenous) choline esters.

### 12.2 Pharmacodynamics

Atropine-induced parasympathetic inhibition may be preceded by a transient phase of stimulation, especially on the heart where small doses first slow the rate before characteristic tachycardia develops due to paralysis of vagal control. Atropine exerts a more potent and prolonged effect on heart, intestine and bronchial muscle than scopolamine, but its action on the iris, ciliary body and certain secretory glands is weaker than that of scopolamine. Unlike the latter, atropine in clinical doses does not depress the central nervous system but may stimulate the medulla and higher cerebral centers. Although mild vagal excitation occurs, the increased respiratory rate and (sometimes) increase depth of respiration produced by atropine are more probably the

result of bronchiolar dilatation. Accordingly, atropine is an unreliable respiratory stimulant and large or repeated doses may depress respiration.

Adequate doses of atropine abolish various types of reflex vagal cardiac slowing or asystole. The drug also prevents and abolishes bradycardia or asystole produced by injection of choline esters, anticholinesterase agents or other parasympathomimetic drugs, and cardiac arrest produced by stimulation of the vagus. Atropine also may lessen the degree of partial heart block when vagal activity is an etiologic factor. In some patients with complete heart block, the idioventricular rate may be accelerated by atropine; in others, the rate is stabilized. Occasionally a large dose may cause atrioventricular (A-V) block and nodal rhythm.

Atropine Sulfate Injection, USP in clinical doses counteracts the peripheral dilatation and abrupt decrease in blood pressure produced by choline esters. However, when given by itself, atropine does not exert a striking or uniform effect on blood vessels or blood pressure. Systemic doses slightly raise systolic and lower diastolic pressures and can produce significant postural hypotension. Such doses also slightly increase cardiac output and decrease central venous pressure. Occasionally, therapeutic doses dilate cutaneous blood vessels, particularly in the “blush” area (atropine flush), and may cause atropine “fever” due to suppression of sweat gland activity in infants and small children.

The effects of intravenous atropine on heart rate (maximum heart rate) and saliva flow (minimum flow) after intravenous administration (rapid, constant infusion over 3 min.) are delayed by 7 to 8 minutes after drug administration and both effects are non-linearly related to the amount of drug in the peripheral compartment. Changes in plasma atropine levels following intramuscular administration (0.5 to 4 mg doses) and heart rate are closely overlapped but the time course of the changes in atropine levels and behavioral impairment indicates that pharmacokinetics is not the primary rate-limiting mechanism for the central nervous system effect of atropine.

### 12.3 Pharmacokinetics

Atropine disappears rapidly from the blood following injection and is distributed throughout the body. Exercise, both prior to and immediately following intramuscular administration of atropine, significantly increases the absorption of atropine due to increased perfusion in the muscle and significantly decreases the clearance of atropine. The pharmacokinetics of atropine is nonlinear after intravenous administration of 0.5 to 4 mg. Atropine's plasma protein binding is about 44% and saturable in the 2-20 µg/mL concentration range. Atropine readily crosses the placental barrier and enters the fetal circulation, but is not found in amniotic fluid. Much of the drug is destroyed by enzymatic hydrolysis, particularly in the liver; from 13 to 50% is excreted unchanged in the urine. Traces are found in various secretions, including milk. The major metabolites of atropine are noratropine, atropin-n-oxide, tropine, and tropic acid. The metabolism of atropine is inhibited by organophosphate pesticides.

#### Specific Populations

The elimination half-life of atropine is more than doubled in children under two years and the elderly (>65 years old) compared to other age groups. There is no gender effect on the pharmacokinetics and pharmacodynamics (heart rate changes) of atropine.

### Nonclinical Toxicology

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Studies have not been performed to evaluate the carcinogenic or mutagenic potential of atropine or its potential to affect fertility adversely.

#### How Supplied

Atropine Sulfate Injection, USP, 0.1 mg/mL, is a clear and colorless solution available in 5 mL and 10 mL single-dose glass vials. Each vial is co-packaged with an injector, which together make a LifeShield Abboject Glass Syringe. It is supplied in the following presentations:

Container	Total Content (Concentration)	Volume	NDC# (Unit of Sale)
LifeShield™ Abboject™ Glass Syringe	0.5 mg/5 mL (0.1 mg/mL)	5 mL	0409-4910-34 (Bundle of 10)
LifeShield™ Abboject™ Glass Syringe	1 mg/10 mL (0.1 mg/mL)	10 mL	0409-4911-34 (Bundle of 10)

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F). [See USP Controlled Room Temperature.]

#### Product repackaged by: Henry Schein, Inc., Bastian, VA 24314

From Original Manufacturer/Distributor's NDC and Unit of Sale	To Henry Schein Repackaged Product NDC and Unit of Sale	Total Strength/Total Volume (Concentration) per unit
NDC 0409-4910-34 Bundle of 10 LifeShield™ Abboject™ Glass Syringe	NDC 0404-9822-05 1 Lifeshield™ Abboject™ Glass Syringe in a bag (Syringe bears NDC 0409- 4910-11)	Concentration: 0.1 mg/mL Fill Volume: 5 mL Total Atropine Content: 0.5 mg
NDC 0409-4911-34 Bundle of 10 LifeShield™ Abboject™ Glass Syringe	NDC 0404-9821-10 1 Lifeshield™ Abboject™ Glass Syringe in a bag (Syringe bears NDC 0409- 4911-11)	Concentration: 0.1 mg/mL Fill Volume: 10 mL Total Atropine Content: 1 mg



Distributed by Hospira, Inc., Lake Forest, IL 60045 USA

Abboject™ is a trademark of the Abbott group of companies.

LifeShield™ is the trademark of ICU Medical, Inc. and is used under license.

LAB-0879-4.0

### Sample Package Label

<b>ATROPINE SULFATE</b> <b>1mg/10ml</b>	
<b>0.1 mg/ml</b> <b>10 ml</b>	<b>INJECTION, USP</b> <b>Abboject Syringe</b>
<b>FOR I.V. USE.</b> <b>USE ASEPTIC TECHNIQUE. DO NOT ASSEMBLE UNTIL</b> <b>READY TO USE. DISCARD UNUSED PORTION. LIQUID</b> <b>IN GLASS. HANDLE WITH CARE. SINGLE-DOSE SYRINGE.</b>	
<b>Keep out of</b> <b>children's reach.</b>	
<b>Store at controlled room</b> <b>temperature 68F to 77F. Excursions</b> <b>permitted between 59F and 86F.</b>	
<b>MANUFACTURER INFORMATION</b> <b>Mfr: Hospira</b> <b>ORIG MFG LOT: XX-XXX-XX</b> <b>NDC: 0409-4911-34</b>	

RX ONLY

NDC:   
0404-9821-10

ITEM# :2480979  
LOT# XXXXXXXXX  
EXP: mm-dd-yy

SEE MANUFACTURER'S INSERT  
FOR COMPLETE PRODUCT AND  
PRESCRIBING INFORMATION

Packaged By  
Henry Schein, Inc.  
80 Summit View Lane  
Bastian, VA 24314



GTIN:(01)XXXXXXXXXXXXXXXXX  
SER:(21)XXXXXXXXXXXXXXXXX  
LOT:(10)XXXXXX  
EXP:(17)XXXXXX



# ATROPINE SULFATE

atropine sulfate injection, solution

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0404-9821(NDC:0409-4911)
<b>Route of Administration</b>	INTRAVENOUS, INTRAMUSCULAR, SUBCUTANEOUS		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>Atropine Sulfate</b> (UNII: 03J5ZE7KA5) (Atropine - UNII:7C0697DR9I)	Atropine Sulfate	0.1 mg in 1 mL

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>WATER</b> (UNII: 059QF0KO0R)	

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:0404-9821-10	1 in 1 BAG	01/09/2022	
1		10 mL in 1 SYRINGE, GLASS; Type 1: Convenience Kit of Co-Package		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
unapproved drug other		01/09/2022	

**Labeler** - Henry Schein, Inc. (012430880)

Revised: 10/2023

Henry Schein, Inc.