

**KETAMINE HYDROCHLORIDE- ketamine hydrochloride injection, solution**  
**General Injectables and Vaccines, Inc.**

-----  
**Ketamine Hydrochloride**

200 mg/20mL (10 mg/mL), 500 mg/10mL (50 mg/mL) and 1,000 mg/10 mL (100 mg/mL)

Rx Only

For intravenous and intramuscular use

**SPECIAL NOTE:** EMERGENCE REACTIONS HAVE OCCURRED IN APPROXIMATELY 12 PERCENT OF PATIENTS.

THE PSYCHOLOGICAL MANIFESTATIONS VARY IN SEVERITY BETWEEN PLEASANT DREAM-LIKE STATES, VIVID IMAGERY, HALLUCINATIONS, AND EMERGENCE DELIRIUM. IN SOME CASES THESE STATES HAVE BEEN ACCOMPANIED BY CONFUSION, EXCITEMENT, AND IRRATIONAL BEHAVIOR WHICH A FEW PATIENTS RECALL AS AN UNPLEASANT EXPERIENCE. THE DURATION ORDINARILY IS NO MORE THAN A FEW HOURS; IN A FEW CASES, HOWEVER, RECURRENCES HAVE TAKEN PLACE UP TO 24 HOURS POSTOPERATIVELY. NO RESIDUAL PSYCHOLOGICAL EFFECTS ARE KNOWN TO HAVE RESULTED FROM USE OF KETAMINE HYDROCHLORIDE INJECTION.

THE INCIDENCE OF THESE EMERGENCE PHENOMENA IS LEAST IN THE ELDERLY (OVER 65 YEARS OF AGE) PATIENT. ALSO, THEY ARE LESS FREQUENT WHEN THE DRUG IS GIVEN INTRAMUSCULARLY AND THE INCIDENCE IS REDUCED AS EXPERIENCE WITH THE DRUG IS GAINED.


THE INCIDENCE OF PSYCHOLOGICAL MANIFESTATIONS DURING EMERGENCE, PARTICULARLY DREAM-LIKE OBSERVATIONS AND EMERGENCE DELIRIUM, MAY BE REDUCED BY USING LOWER RECOMMENDED DOSAGES OF KETAMINE HYDROCHLORIDE INJECTION IN CONJUNCTION WITH INTRAVENOUS DIAZEPAM DURING INDUCTION AND MAINTENANCE OF ANESTHESIA. (See DOSAGE AND ADMINISTRATION section.) ALSO, THESE REACTIONS MAY BE REDUCED IF VERBAL, TACTILE, AND VISUAL STIMULATION OF THE PATIENT IS MINIMIZED DURING THE RECOVERY PERIOD. THIS DOES NOT PRECLUDE THE MONITORING OF VITAL SIGNS.

IN ORDER TO TERMINATE A SEVERE EMERGENCE REACTION, THE USE OF A SMALL HYPNOTIC DOSE OF A SHORT-ACTING OR ULTRA SHORT-ACTING BARBITURATE MAY BE REQUIRED.

WHEN KETAMINE HYDROCHLORIDE INJECTION IS USED ON AN OUTPATIENT BASIS, THE PATIENT SHOULD NOT BE RELEASED UNTIL RECOVERY FROM ANESTHESIA IS COMPLETE AND THEN SHOULD BE ACCOMPANIED BY A RESPONSIBLE ADULT.

**DESCRIPTION:**

Ketamine hydrochloride injection, USP is a nonbarbiturate general anesthetic chemically designated *dl* 2-(*o*-chlorophenyl)-2-(methylamino) cyclohexanone hydrochloride. It is formulated as a slightly acid (pH 3.5-5.5) sterile solution for intravenous or intramuscular injection in concentrations containing the equivalent of either 10, 50 or 100 mg ketamine base per milliliter and contains not more than 0.1 mg/mL benzethonium chloride added as a preservative. The 10 mg/mL solution has been made isotonic with sodium chloride.



Molecular Weight: 274.19  
 $C_{13}H_{16}ClNO \cdot HCl$

### CLINICAL PHARMACOLOGY:

Ketamine hydrochloride injection is a rapid-acting general anesthetic producing an anesthetic state characterized by profound analgesia, normal pharyngeal-laryngeal reflexes, normal or slightly enhanced skeletal muscle tone, cardiovascular and respiratory stimulation, and occasionally a transient and minimal respiratory depression. The mechanism of action is primarily due to antagonism of N-methyl-D-aspartate (NMDA receptors) in the central nervous system.

A patent airway is maintained partly by virtue of unimpaired pharyngeal and laryngeal reflexes. (See WARNINGS and PRECAUTIONS sections.)

The biotransformation of ketamine hydrochloride injection includes N-dealkylation (metabolite I), hydroxylation of the cyclohexone ring (metabolites III and IV), conjugation with glucuronic acid and dehydration of the hydroxylated metabolites to form the cyclohexene derivative (metabolite II).

Following intravenous administration, the ketamine concentration has an initial slope (alpha phase) lasting about 45 minutes with a half-life of 10 to 15 minutes. This first phase corresponds clinically to the anesthetic effect of the drug. The anesthetic action is terminated by a combination of redistribution from the CNS to slower equilibrating peripheral tissues and by hepatic biotransformation to metabolite I. This metabolite is about 1/3 as active as ketamine in reducing halothane requirements (MAC) of the rat. The later half-life of ketamine (beta phase) is 2.5 hours.

The anesthetic state produced by ketamine hydrochloride injection has been termed "dissociative anesthesia" in that it appears to selectively interrupt association pathways of the brain before producing somesthetic sensory blockade. It may selectively depress the thalamoneocortical system before significantly obtunding the more ancient cerebral centers and pathways (reticular-activating and limbic systems).

Elevation of blood pressure begins shortly after injection, reaches a maximum within a few minutes and usually returns to preanesthetic values within 15 minutes after injection. In the majority of cases, the systolic and diastolic blood pressure peaks from 10% to 50% above preanesthetic levels shortly after induction of anesthesia, but the elevation can be higher or longer in individual cases (see CONTRAINDICATIONS section).

Ketamine has a wide margin of safety; several instances of unintentional administration of overdoses of ketamine hydrochloride injection (up to ten times that usually required) have been followed by prolonged but complete recovery.

Ketamine hydrochloride injection has been studied in over 12,000 operative and diagnostic procedures, involving over 10,000 patients from 105 separate studies. During the course of these studies ketamine hydrochloride injection was administered as the sole agent, as induction for other general agents, or to supplement low-potency agents.

Specific areas of application have included the following:

1. debridement, painful dressings, and skin grafting in burn patients, as well as other superficial surgical procedures.
2. neurodiagnostic procedures, such as pneumoencephalograms, ventriculograms, myelograms, and lumbar punctures. See also Precaution concerning increased intracranial pressure.
3. diagnostic and operative procedures of the eye, ear, nose, and mouth, including dental extractions.
4. diagnostic and operative procedures of the pharynx, larynx, or bronchial tree. NOTE: Muscle relaxants, with proper attention to respiration, may be required (see PRECAUTIONS section).
5. sigmoidoscopy and minor surgery of the anus and rectum, and circumcision.
6. extraperitoneal procedures used in gynecology such as dilatation and curettage.
7. orthopedic procedures such as closed reductions, manipulations, femoral pinning, amputations, and biopsies.
8. as an anesthetic in poor-risk patients with depression of vital functions.
9. in procedures where the intramuscular route of administration is preferred.
10. in cardiac catheterization procedures.

In these studies, the anesthesia was rated either “excellent” or “good” by the anesthesiologist and the surgeon at 90% and 93%, respectively; rated “fair” at 6% and 4%, respectively; and rated “poor” at 4% and 3%, respectively. In a second method of evaluation, the anesthesia was rated “adequate” in at least 90%, and “inadequate” in 10% or less of the procedures.

#### **INDICATIONS AND USAGE:**

Ketamine hydrochloride injection is indicated as the sole anesthetic agent for diagnostic and surgical procedures that do not require skeletal muscle relaxation. Ketamine hydrochloride injection is best suited for short procedures but it can be used, with additional doses, for longer procedures.

Ketamine hydrochloride injection is indicated for the induction of anesthesia prior to the administration of other general anesthetic agents.

Ketamine hydrochloride injection is indicated to supplement low-potency agents, such as nitrous oxide.

Specific areas of application are described in the CLINICAL PHARMACOLOGY section.

#### **CONTRAINDICATIONS:**

Ketamine hydrochloride is contraindicated in those in whom a significant elevation of blood pressure would constitute a serious hazard and in those who have shown hypersensitivity to the drug.

#### **WARNINGS:**

Cardiac function should be continually monitored during the procedure in patients found to have hypertension or cardiac decompensation.

Postoperative confusional states may occur during the recovery period. (See SPECIAL NOTE.)

Respiratory depression may occur with overdose or too rapid a rate of administration of ketamine hydrochloride injection, in which case supportive ventilation should be employed. Mechanical support of respiration is preferred to administration of analeptics.

**Pediatric Neurotoxicity:** Published animal studies demonstrate that the administration of anesthetic and

sedation drugs that block NMDA receptors and/or potentiate GABA activity increase neuronal apoptosis in the developing brain and result in long-term cognitive deficits when used for longer than 3 hours. The clinical significance of these findings is not clear. However, based on the available data, the window of vulnerability to these changes is believed to correlate with exposures in the third trimester of gestation through the first several months of life, but may extend out to approximately three years of age in humans. (See PRECAUTIONS: Pregnancy.)

Some published studies in children suggest that similar deficits may occur after repeated or prolonged exposures to anesthetic agents early in life and may result in adverse cognitive or behavioral effects. These studies have substantial limitations, and it is not clear if the observed effects are due to the anesthetic/sedation drug administration or other factors such as the surgery or underlying illness.

Anesthetic and sedation drugs are a necessary part of the care of children needing surgery, other procedures, or tests that cannot be delayed, and no specific medications have been shown to be safer than any other. Decisions regarding the timing of any elective procedures requiring anesthesia should take into consideration the benefits of the procedure weighed against the potential risks.

## **PRECAUTIONS:**

**General:** Ketamine hydrochloride injection should be used by or under the direction of physicians experienced in administering general anesthetics and in maintenance of an airway and in the control of respiration.

Because pharyngeal and laryngeal reflexes are usually active, ketamine hydrochloride injection should not be used alone in surgery or diagnostic procedures of the pharynx, larynx, or bronchial tree. Mechanical stimulation of the pharynx should be avoided, whenever possible, if ketamine hydrochloride injection is used alone. Muscle relaxants, with proper attention to respiration, may be required in both of these instances.

Resuscitative equipment should be ready for use.

The *incidence of emergence reactions may be reduced* if verbal and tactile stimulation of the patient is minimized during the recovery period. This does not preclude the monitoring of vital signs (see SPECIAL NOTE).

The intravenous dose should be administered over a period of 60 seconds. More rapid administration may result in respiratory depression or apnea and enhanced pressor response.

In surgical procedures involving visceral pain pathways, ketamine hydrochloride injection should be supplemented with an agent which obtunds visceral pain.

Use with caution in the chronic alcoholic and the acutely alcohol-intoxicated patient.

An increase in cerebrospinal fluid pressure has been reported following administration of ketamine hydrochloride. Use with extreme caution in patients with preanesthetic elevated cerebrospinal fluid pressure.

**Carcinogenesis, Mutagenesis, Impairment of Fertility: Carcinogenesis:** Long-term animal studies have not been conducted to evaluate the carcinogenic potential of ketamine.

**Mutagenesis:** In a published report, ketamine was clastogenic in the *in vitro* chromosomal aberration assay.

**Impairment of Fertility:** Adequate studies to evaluate the impact of ketamine on male or female fertility have not been conducted. Male and female rats were treated with 10 mg/kg ketamine IV (0.8 times the average human induction dose of 2 mg/kg IV based on body surface area) on Days 11, 10, and 9 prior to mating. No impact on fertility was noted; however, this study design does not adequately characterize the impact of a drug on fertility endpoints.

**Pregnancy: Risk Summary:** There are no adequate and well-controlled studies of ketamine

hydrochloride injection in pregnant women. In animal reproduction studies in rats developmental delays (hypoplasia of skeletal tissues) were noted at 0.3 times the human intramuscular dose of 10 mg/kg. In rabbits, developmental delays and increased fetal resorptions were noted at 0.6 times the human dose. Published studies in pregnant primates demonstrate that the administration of anesthetic and sedation drugs that block NMDA receptors and/or potentiate GABA activity during the period of peak brain development increases neuronal apoptosis in the developing brain of the offspring when used for longer than 3 hours. There are no data on pregnancy exposures in primates corresponding to periods prior to the third trimester in humans.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

**Clinical Considerations:** Since the safe use in pregnancy, including obstetrics (either vaginal or abdominal delivery), has not been established, such use is not recommended (see ANIMAL PHARMACOLOGY AND TOXICOLOGY).

**Data: Animal Data:** Pregnant rats were treated intramuscularly with 20 mg/kg ketamine (0.3 times the human dose of 10 mg/kg IM based on body surface area) on either Gestation Days 6 to 10 or Gestation Days 11 to 15. Ketamine treatment produced an increased incidence of hypoplastic skull, phalanges, and sternebrae in the pups.

Pregnant rabbits were treated intramuscularly with 20 mg/kg ketamine (0.6 times the human dose of 10 mg/kg IM based on body surface area) on either Gestation Days 6 to 10 or Gestation Days 11 to 15. An increase in resorptions and skeletal hypoplasia of the fetuses were noted. Additional pregnant rabbits were treated intramuscularly with a single dose 60 mg/kg (1.9 times the human dose of 10 mg/kg IM based on body surface area) on Gestation Day 6 only. Skeletal hypoplasia was reported in the fetuses.

In a study where pregnant rats were treated intramuscularly with 20 mg/kg ketamine (0.3 times the human dose of 10 mg/kg IM based on body surface area) from Gestation Day 18 to 21. There was a slight increase in incidence of delayed parturition by one day in treated dams of this group. No adverse effects on the litters or pups were noted; however, learning and memory assessments were not completed.

Three pregnant beagle dogs were treated intramuscularly with 25 mg/kg ketamine (1.3 times the human dose of 10 mg/kg IM based on body surface area) twice weekly for the three weeks of the first, second and third trimesters of pregnancy, respectively, without the development of adverse effects in the pups.

In a published study in primates, administration of an anesthetic dose of ketamine for 24 hours on Gestation Day 122 increased neuronal apoptosis in the developing brain of the fetus. In other published studies, administration of either isoflurane or propofol for 5 hours on Gestation Day 120 resulted in increased neuronal and oligodendrocyte apoptosis in the developing brain of the offspring. With respect to brain development, this time period corresponds to the third trimester of gestation in the human. The clinical significance of these findings is not clear; however, studies in juvenile animals suggest neuroapoptosis correlates with long-term cognitive deficits. (See WARNINGS: Pediatric Neurotoxicity, PRECAUTIONS: Pediatric Use, and ANIMAL TOXICOLOGY AND PHARMACOLOGY.)

**Information for Patients: Risk of Drowsiness:** As appropriate, especially in cases where early discharge is possible, the duration of ketamine hydrochloride injection and other drugs employed during the conduct of anesthesia should be considered. The patients should be cautioned that driving an automobile, operating hazardous machinery or engaging in hazardous activities should not be undertaken for 24 hours or more (depending upon the dosage of ketamine hydrochloride injection and consideration of other drugs employed) after anesthesia.

**Effect of Anesthetic and Sedation Drugs on Early Brain Development:** Studies conducted in young animals and children suggest repeated or prolonged use of general anesthetic or sedation drugs in children younger than 3 years may have negative effects on their developing brains. Discuss with parents and caregivers the benefits, risks, and timing and duration of surgery or procedures requiring anesthetic and sedation drugs (See WARNINGS: Pediatric Neurotoxicity).

**Drug Interactions:** Prolonged recovery time may occur if barbiturates and/or narcotics are used concurrently with ketamine hydrochloride injection. Ketamine hydrochloride injection is clinically compatible with the commonly used general and local anesthetic agents when an adequate respiratory exchange is maintained.

**Geriatric Use:** Clinical studies of ketamine hydrochloride did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

**Pediatric Use:** Safety and effectiveness in pediatric patients below the age of 16 have not been established.

Published juvenile animal studies demonstrate that the administration of anesthetic and sedation drugs, such as ketamine hydrochloride injection, that either block NMDA receptors or potentiate the activity of GABA during the period of rapid brain growth or synaptogenesis, results in widespread neuronal and oligodendrocyte cell loss in the developing brain and alterations in synaptic morphology and neurogenesis. Based on comparisons across species, the window of vulnerability to these changes is believed to correlate with exposures in the third trimester of gestation through the first several months of life, but may extend out to approximately 3 years of age in humans.

In primates, exposure to 3 hours of ketamine that produced a light surgical plane of anesthesia did not increase neuronal cell loss, however, treatment regimens of 5 hours or longer of isoflurane increased neuronal cell loss. Data from isoflurane-treated rodents and ketamine-treated primates suggest that the neuronal and oligodendrocyte cell losses are associated with prolonged cognitive deficits in learning and memory. The clinical significance of these nonclinical findings is not known, and healthcare providers should balance the benefits of appropriate anesthesia in pregnant women, neonates, and young children who require procedures with the potential risks suggested by the nonclinical data. (See WARNINGS: Pediatric Neurotoxicity and PRECAUTIONS: Pregnancy.)

## **ADVERSE REACTIONS:**

**Cardiovascular:** Blood pressure and pulse rate are frequently elevated following administration of ketamine hydrochloride injection alone. However, hypotension and bradycardia have been observed. Arrhythmia has also occurred.

**Respiration:** Although respiration is frequently stimulated, severe depression of respiration or apnea may occur following rapid intravenous administration of high doses of ketamine hydrochloride injection. Laryngospasms and other forms of airway obstruction have occurred during ketamine hydrochloride injection anesthesia.

**Eye:** Diplopia and nystagmus have been noted following ketamine hydrochloride injection administration. It also may cause a slight elevation in intraocular pressure measurement.

**Genitourinary:** In individuals with history of chronic ketamine use or abuse, lower urinary tract and bladder symptoms including dysuria, increased urinary frequency, urgency, urge incontinence, and hematuria have been reported (see DOSAGE AND ADMINISTRATION section). In addition, diagnostic studies performed to assess the cause of these symptoms have reported cystitis (including cystitis non-infective, cystitis interstitial, cystitis ulcerative, cystitis erosive and cystitis hemorrhagic)

as well as hydronephrosis and reduced bladder capacity.

**Psychological:** (See SPECIAL NOTE.)

**Neurological:** In some patients, enhanced skeletal muscle tone may be manifested by tonic and clonic movements sometimes resembling seizures (see DOSAGE AND ADMINISTRATION section).

**Gastrointestinal:** Anorexia, nausea and vomiting have been observed; however, this is not usually severe and allows the great majority of patients to take liquids by mouth shortly after regaining consciousness (see DOSAGE AND ADMINISTRATION section). **General:** Anaphylaxis. Local pain and exanthema at the injection site have infrequently been reported. Transient erythema and/or morbilliform rash have also been reported.

**or medical advice about adverse reactions contact your medical professional. To report SUSPECTED ADVERSE REACTIONS, contact Mylan at 1-877-446-3679 (1-877-4-INFO-RX) or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### **DRUG ABUSE AND DEPENDENCE:**

Ketamine has been reported being used as a drug of abuse.

Reports suggest that ketamine produces a variety of symptoms including, but not limited to anxiety, dysphoria, disorientation, insomnia, flashbacks, hallucinations, and psychotic episodes.

Ketamine dependence and tolerance are possible following prolonged administration. A withdrawal syndrome with psychotic features has been described following discontinuation of long-term ketamine use. Therefore, ketamine should be prescribed and administered with caution.

### **OVERDOSAGE:**

Respiratory depression may occur with overdose or too rapid a rate of administration of ketamine hydrochloride injection, in which case supportive ventilation should be employed. Mechanical support of respiration is preferred to administration of analeptics.

### **DOSAGE AND ADMINISTRATION:**

**Note:** Barbiturates and ketamine hydrochloride injection, being chemically incompatible because of precipitate formation, *should not* be injected from the same syringe.

If the ketamine hydrochloride injection dose is augmented with diazepam, the two drugs must be given separately. Do not mix ketamine hydrochloride injection and diazepam in the syringe or infusion flask. For additional information on the use of diazepam, refer to the WARNINGS and DOSAGE AND ADMINISTRATION sections of the diazepam insert.

### **Preoperative Preparations:**

1. While vomiting has been reported following ketamine hydrochloride injection administration, some airway protection may be afforded because of active laryngeal-pharyngeal reflexes. However, since aspiration may occur with ketamine hydrochloride injection and since protective reflexes may also be diminished by supplementary anesthetics and muscle relaxants, the possibility of aspiration must be considered. Ketamine hydrochloride injection is recommended for use in the patient whose stomach is not empty when, in the judgment of the practitioner, the benefits of the drug outweigh the possible risks.
2. Atropine, scopolamine, or other drying agent should be given at an appropriate interval prior to induction.

**Onset and Duration:** Because of rapid induction following the initial intravenous injection, the patient should be in a supported position during administration.

The onset of action of ketamine hydrochloride injection is rapid; an intravenous dose of 2 mg/kg of body weight usually produces surgical anesthesia within 30 seconds after injection, with the anesthetic effect usually lasting five to ten minutes. If a longer effect is desired, additional increments can be administered intravenously or intramuscularly to maintain anesthesia without producing significant cumulative effects.

Intramuscular doses, in a range of 9 mg/kg to 13 mg/kg usually produce surgical anesthesia within 3 to 4 minutes following injection, with the anesthetic effect usually lasting 12 to 25 minutes.

**Dosage:** As with other general anesthetic agents, the individual response to ketamine hydrochloride injection is somewhat varied depending on the dose, route of administration, and age of patient, so that dosage recommendation cannot be absolutely fixed. The drug should be titrated against the patient's requirements.

In individuals with a history of chronic ketamine use for off-label indications, there have been case reports of genitourinary pain that may be related to the ketamine treatment, not the underlying condition (see ADVERSE REACTIONS section). Consider cessation of ketamine if genitourinary pain continues in the setting of other genitourinary symptoms.

**Induction: Intravenous Route:** The initial dose of ketamine hydrochloride injection administered intravenously may range from 1 mg/kg to 4.5 mg/kg. The average amount required to produce five to ten minutes of surgical anesthesia has been 2 mg/kg.

Alternatively, in adult patients an induction dose of 1 mg to 2 mg/kg intravenous ketamine at a rate of 0.5 mg/kg/min may be used for induction of anesthesia. In addition, diazepam in 2 mg to 5 mg doses, administered in a separate syringe over 60 seconds, may be used. In most cases, 15 mg of intravenous diazepam *or less* will suffice. The incidence of psychological manifestations during emergence, particularly dream-like observations and emergence delirium, may be reduced by this induction dosage program.

**Note:** The 100 mg/mL concentration of ketamine hydrochloride injection should not be injected intravenously without proper dilution. It is recommended the drug be diluted with an equal volume of either Sterile Water for Injection, USP, Normal Saline, or 5% Dextrose in Water.

**Rate of Administration:** It is recommended that ketamine hydrochloride injection be administered slowly (over a period of 60 seconds). More rapid administration may result in respiratory depression and enhanced pressor response.

**Intramuscular Route:** The initial dose of ketamine hydrochloride injection administered intramuscularly may range from 6.5 to 13 mg/kg. A dose of 10 mg/kg will usually produce 12 to 25 minutes of surgical anesthesia.

**Maintenance of Anesthesia:** The maintenance dose should be adjusted according to the patient's anesthetic needs and whether an additional anesthetic agent is employed.

Increments of one-half to the full induction dose may be repeated as needed for maintenance of anesthesia. However, it should be noted that purposeless and tonic-clonic movements of extremities may occur during the course of anesthesia. These movements do not imply a light plane and are not indicative of the need for additional doses of the anesthetic.

It should be recognized that the larger the total dose of ketamine hydrochloride injection administered, the longer will be the time to complete recovery.

Adult patients induced with ketamine hydrochloride injection augmented with intravenous diazepam may be maintained on ketamine hydrochloride injection given by slow microdrip infusion technique at a dose of 0.1 to 0.5 mg/minute, augmented with diazepam 2 to 5 mg administered intravenously as needed. In many cases 20 mg or *less* of intravenous diazepam total for combined induction and maintenance will suffice. However, slightly more diazepam may be required depending on the nature and duration of the operation, physical status of the patient, and other factors. The incidence of psychological

manifestations during emergence, particularly dream-like observations and emergence delirium, may be reduced by this maintenance dosage program.

**Dilution:** To prepare a dilute solution containing 1 mg of ketamine per mL, aseptically transfer 10 mL from a 50 mg per mL vial or 5 mL from a 100 mg per mL vial to 500 mL of 5% Dextrose Injection, USP or Sodium Chloride (0.9%) Injection, USP (Normal Saline) and mix well. The resultant solution will contain 1 mg of ketamine per mL.

The fluid requirements of the patient and duration of anesthesia must be considered when selecting the appropriate dilution of ketamine hydrochloride injection. If fluid restriction is required, ketamine hydrochloride injection can be added to a 250 mL infusion as described above to provide a ketamine hydrochloride injection concentration of 2 mg/mL.

Ketamine hydrochloride injection 10 mg/mL vials are not recommended for dilution.

**Supplementary Agents:** Ketamine hydrochloride injection is clinically compatible with the commonly used general and local anesthetic agents when an adequate respiratory exchange is maintained.

The regimen of a reduced dose of ketamine hydrochloride injection supplemented with diazepam can be used to produce balanced anesthesia by combination with other agents such as nitrous oxide and oxygen.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

## **HOW SUPPLIED:**

Ketamine Hydrochloride Injection, USP is supplied as the hydrochloride in concentrations equivalent to ketamine base.

NDC 67457-181-20	carton containing 10, 20 mL multi-dose vials with 10 mg/mL
NDC 67457-001-10	carton containing 10, 10 mL multi-dose vials with 50 mg/mL
NDC 67457-108-10	carton containing 10, 10 mL multi-dose vials with 100 mg/mL

**Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]**

**Protect from light. Retain in carton until time of use.**

## **ANIMAL PHARMACOLOGY AND TOXICOLOGY:**

Published studies in animals demonstrate that the use of anesthetic agents during the period of rapid brain growth or synaptogenesis results in widespread neuronal and oligodendrocyte cell loss in the developing brain and alterations in synaptic morphology and neurogenesis. Based on comparisons across species, the window of vulnerability to these changes is believed to correlate with exposures in the third trimester through the first several months of life, but may extend out to approximately 3 years of age in humans.

In primates, exposure to 3 hours of an anesthetic regimen that produced a light surgical plane of anesthesia did not increase neuronal cell loss, however, treatment regimens of 5 hours or longer increased neuronal cell loss. Data in rodents and in primates suggest that the neuronal and oligodendrocyte cell losses are associated with subtle but prolonged cognitive deficits in learning and memory. The clinical significance of these nonclinical findings is not known, and healthcare providers should balance the benefits of appropriate anesthesia in neonates and young children who require procedures against the potential risks suggested by the nonclinical data. (See WARNINGS: Pediatric Neurotoxicity and PRECAUTIONS: Pregnancy, Pediatric Use.)

In published studies, intraperitoneal administration of ketamine at doses greater than 40 mg/kg induced vacuolation in neuronal cells of the posterior cingulate and retrosplenial cortices in adult rats, similar to what has been reported in rodents administered other NMDA receptor antagonists. These

vacuoles were demonstrated to be reversible and did not progress to degeneration or neuronal death up to doses of 80 mg/kg (1.2 times the human dose of 10 mg/kg based on body surface area). A no-effect level for neuronal vacuolation was 20 mg/kg intraperitoneal (0.3 times a human dose of 10 mg/kg on a body surface area basis). The window of vulnerability to these changes is believed to correlate with exposures in humans from the onset of puberty through adulthood. The relevance of this finding to humans is unknown.



Manufactured for:  
**Mylan Institutional LLC**  
Rockford, IL 61103 U.S.A.

Manufactured by:  
**Mylan Institutional**  
Galway, Ireland

0942L101

Revised: 8/2018  
MI:KETAIJ:R6

**SAMPLE PACKAGE LABEL**

NDC # 52584 - 108 - 00  
ITEM# :2480700  
LOT # XXXXXXXXX  
EXP : mm - yy

Packaged By  
General Injectables and Vaccines, Inc  
80 Summit View Lane  
Bastian, VA 24314



**KETAMINE HYDROCHLORIDE  
1,000MG/10ML**

100 MG/ML

INJECTION, USP

10 ML

MULTI DOSE VIAL

SEE MANUFACTURER'S INSERT  
FOR COMPLETE PRODUCT AND  
PRESCRIBING INFORMATION

PROTECT FROM LIGHT. RETAIN IN BAG UNTIL TIME OF  
USE. FOR IM OR SLOW IV USE. MUST BE DILUTED BEFORE  
IV USE. STERILE. DO NOT USE IF PRECIPITATE  
APPEARS.

Keep out of  
children's reach.

Store at controlled room  
temperature 68F to 77F.



**MANUFACTURER INFORMATION**

Mfr: Mylan Institutional  
ORIG MFG LOT: XX - XXX - XX  
NDC:67457 - 108 - 10

(10)XXXXXX  
(17)XXXXXXXX  
(01)XXXXXXXXXXXX  
(21)XXXXXXXXXXXX  
RX ONLY

NDC # 52584 - 108 - 00  
 ITEM# :2480700  
 LOT # XXXXXXXXX  
 EXP : mm - yy

Packaged By  
 General Injectables and Vaccines, Inc  
 80 Summit View Lane  
 Bastian, VA 24314



**KETAMINE HYDROCHLORIDE**  
**1,000MG/10ML**  
 100 MG/ML INJECTION, USP  
 10 ML MULTI DOSE VIAL

SEE MANUFACTURER'S INSERT  
 FOR COMPLETE PRODUCT AND  
 PRESCRIBING INFORMATION

PROTECT FROM LIGHT. RETAIN IN BAG UNTIL TIME OF  
 USE. FOR IM OR SLOW IV USE. MUST BE DILUTED BEFORE  
 IV USE. STERILE. DO NOT USE IF PRECIPITATE  
 APPEARS.

Keep out of  
 children's reach.

Store at controlled room  
 temperature 68F to 77F.



**MANUFACTURER INFORMATION**  
 Mfr: Mylan Institutional  
 ORIG MFG LOT: XX - XXX - XX  
 NDC:67457 - 108 - 10

(10)XXXXXX  
 (17)XXXXXXXX  
 (01)XXXXXXXXXXXX  
 (21)XXXXXXXXXXXX  
 RX ONLY

## KETAMINE HYDROCHLORIDE

ketamine hydrochloride injection, solution

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:52584-108(NDC:67457-108)
Route of Administration	INTRAVENOUS, INTRAMUSCULAR	DEA Schedule	CIII

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
KETAMINE HYDRO CHLORIDE (UNII: O18YU00J83) (KETAMINE - UNII:690G0D6V8H)	KETAMINE	100 mg in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744)	0.1 mg in 1 mL

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52584-108-00	1 in 1 BAG	02/28/2018	
1		10 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076092	02/28/2018	

**Labeler** - General Injectables and Vaccines, Inc. (108250663)

Revised: 7/2019

General Injectables and Vaccines, Inc.