

# HYDRALAZINE HYDROCHLORIDE- hydralazine hydrochloride injection, solution Henry Schein, Inc.

## Hydralazine Hydrochloride

### Hydralazine Hydrochloride Injection, USP

Preservative-Free

Rx only

#### DESCRIPTION

Hydralazine Hydrochloride Injection, USP is an antihypertensive available in a 1 mL vial for intravenous and intramuscular administration. Hydralazine Hydrochloride Injection, USP is a sterile, nonpyrogenic colorless solution.


#### Each mL contains:

**Active:** Hydralazine Hydrochloride USP, 20 mg


#### Preservative Free Formulation

**Inactives:** Propylene Glycol USP, 103.6 mg; Sodium Hydroxide and/or Hydrochloric Acid USP to adjust pH (3.4 to 4.4) and Water for Injection.

Hydralazine Hydrochloride USP is 1-hydrazinophthalazine monohydrochloride, and its structural formula is:



Molecular Formula: C<sub>8</sub>H<sub>8</sub>N<sub>4</sub>•HCl



Molecular Formula: C<sub>8</sub>H<sub>8</sub>N<sub>4</sub>•HCl

Hydralazine Hydrochloride USP is a white to off-white, odorless crystalline powder. It is soluble in water, slightly soluble in alcohol, and very slightly soluble in ether. It melts at about 275°C, with decomposition, and has a molecular weight of 196.64.

## **CLINICAL PHARMACOLOGY**

Although the precise mechanism of action of hydralazine is not fully understood, the major effects are on the cardiovascular system. Hydralazine apparently lowers blood pressure by exerting a peripheral vasodilating effect through a direct relaxation of vascular smooth muscle. Hydralazine, by altering cellular calcium metabolism, interferes with the calcium movements within the vascular smooth muscle that are responsible for initiating or maintaining the contractile state.

The peripheral vasodilating effect of hydralazine results in decreased arterial blood pressure (diastolic more than systolic), decreased peripheral vascular resistance, and an increased heart rate, stroke volume, and cardiac output. The preferential dilatation of arterioles, as compared to veins, minimizes postural hypotension and promotes the increase in cardiac output. Hydralazine usually increases renin activity in plasma, presumably as a result of increased secretion of renin by the renal juxtaglomerular cells in response to reflex sympathetic discharge. This increase in renin activity leads to the production of angiotensin II, which then causes stimulation of aldosterone and consequent sodium reabsorption.

Hydralazine also maintains or increases renal and cerebral blood flow.

The average maximal decrease in blood pressure usually occurs 10 to 80 minutes after administration of hydralazine hydrochloride injection. No other pharmacokinetic data on hydralazine hydrochloride injection are available.

## **INDICATIONS AND USAGE**

Severe essential hypertension when the drug cannot be given orally or when there is an urgent need to lower blood pressure.

## **CONTRAINDICATIONS**

Hypersensitivity to hydralazine; coronary artery disease; mitral valvular rheumatic heart disease.

## **WARNINGS**

In a few patients hydralazine may produce a clinical picture simulating systemic lupus erythematosus including glomerulonephritis. In such patients hydralazine should be discontinued unless the benefit-to-risk determination requires continued antihypertensive therapy with this drug. Symptoms and signs usually regress when the drug is discontinued but residua have been detected many years later. Long-term treatment with steroids may be necessary (see **PRECAUTIONS, LABORATORY TESTS**).

## **PRECAUTIONS**

### **General**

Myocardial stimulation produced by hydralazine can cause anginal attacks and ECG changes of myocardial ischemia. The drug has been implicated in the production of

myocardial infarction. It must, therefore, be used with caution in patients with suspected coronary artery disease.

The “hyperdynamic” circulation caused by hydralazine may accentuate specific cardiovascular inadequacies. For example, hydralazine may increase pulmonary artery pressure in patients with mitral valvular disease. The drug may reduce the pressor responses to epinephrine. Postural hypotension may result from hydralazine but is less common than with ganglionic blocking agents. It should be used with caution in patients with cerebral vascular accidents.

In hypertensive patients with normal kidneys who are treated with hydralazine, there is evidence of increased renal blood flow and a maintenance of glomerular filtration rate. In some instances where control values were below normal, improved renal function has been noted after administration of hydralazine. However, as with any antihypertensive agent, hydralazine should be used with caution in patients with advanced renal damage.

Peripheral neuritis, evidenced by paresthesia, numbness, and tingling, has been observed. Published evidence suggests an antipyridoxine effect, and that pyridoxine should be added to the regimen if symptoms develop.

### **Laboratory Tests**

Complete blood counts and antinuclear antibody titer determinations are indicated before and periodically during prolonged therapy with hydralazine even though the patient is asymptomatic. These studies are also indicated if the patient develops arthralgia, fever, chest pain, continued malaise, or other unexplained signs or symptoms.

A positive antinuclear antibody titer requires that the physician carefully weigh the implications of the test results against the benefits to be derived from antihypertensive therapy with hydralazine hydrochloride.

Blood dyscrasias, consisting of reduction in hemoglobin and red cell count, leukopenia, agranulocytosis, and purpura, have been reported. If such abnormalities develop, therapy should be discontinued.

### **Drug Interactions**

MAO inhibitors should be used with caution in patients receiving hydralazine.

When other potent parenteral antihypertensive drugs, such as diazoxide, are used in combination with hydralazine, patients should be continuously observed for several hours for any excessive fall in blood pressure. Profound hypotensive episodes may occur when diazoxide injection and hydralazine injection are used concomitantly.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

In a lifetime study in Swiss albino mice, there was a statistically significant increase in the incidence of lung tumors (adenomas and adenocarcinomas) of both male and female mice given hydralazine continuously in their drinking water at a dosage of about 250 mg/kg per day (about 80 times the maximum recommended human dose). In a 2-year carcinogenicity study of rats given hydralazine by gavage at dose levels of 15, 30, and 60 mg/kg/day (approximately 5 to 20 times the recommended human daily dosage), microscopic examination of the liver revealed a small, but statistically significant, increase in benign neoplastic nodules in male and female rats from the high-dose group and in female rats from the intermediate-dose group. Benign interstitial cell tumors of the testes were also significantly increased in male rats from the high-dose group. The tumors observed are common in aged rats and a significantly increased incidence was

not observed until 18 months of treatment. Hydralazine was shown to be mutagenic in bacterial systems (Gene Mutation and DNA Repair) and in one of two rats and one rabbit hepatocyte in vitro DNA repair studies. Additional in vivo and in vitro studies using lymphoma cells, germinal cells, and fibroblasts from mice, bone marrow cells from chinese hamsters and fibroblasts from human cell lines did not demonstrate any mutagenic potential for hydralazine.

The extent to which these findings indicate a risk to man is uncertain. While long-term clinical observation has not suggested that human cancer is associated with hydralazine use, epidemiologic studies have so far been insufficient to arrive at any conclusions.

### **Pregnancy**

Teratogenic effects: Animal studies indicate that hydralazine is teratogenic in mice at 20 to 30 times the maximum daily human dose of 200 to 300 mg and possibly in rabbits at 10 to 15 times the maximum daily human dose, but that it is nonteratogenic in rats. Teratogenic effects observed were cleft palate and malformations of facial and cranial bones.

There are no adequate and well-controlled studies in pregnant women. Although clinical experience does not include any positive evidence of adverse effects on the human fetus, hydralazine should be used during pregnancy only if the expected benefit justifies the potential risk to the fetus.

### **Nursing Mothers**

Hydralazine has been shown to be excreted in breast milk. Because many drugs are excreted in human milk, caution should be exercised when hydralazine injection is administered to a nursing woman.

### **Pediatric Use**

Safety and effectiveness in pediatric patients have not been established in controlled clinical trials, although there is experience with the use of hydralazine hydrochloride in children. The usual recommended parenteral dosage, administered intramuscularly or intravenously, is 1.7 to 3.5 mg/kg of body weight daily, divided into four to six doses.

## **ADVERSE REACTIONS**

Adverse reactions with hydralazine hydrochloride are usually reversible when dosage is reduced. However, in some cases it may be necessary to discontinue the drug.

The following adverse reactions have been observed, but there has not been enough systematic collection of data to support an estimate of their frequency.

**Common:** Headache, anorexia, nausea, vomiting, diarrhea, palpitations, tachycardia, angina pectoris.

### **Less Frequent:**

Digestive: constipation, paralytic ileus.

Cardiovascular: hypotension, paradoxical pressor response, edema.

Respiratory: dyspnea.

Neurologic: peripheral neuritis, evidenced by paresthesia, numbness, and tingling; dizziness; tremors; muscle cramps; psychotic reactions characterized by depression, disorientation, or anxiety.

Genitourinary: difficulty in urination.

Hematologic: blood dyscrasias, consisting of reduction in hemoglobin and red cell count, leukopenia, agranulocytosis, purpura; lymphadenopathy; splenomegaly.

Hypersensitive Reactions: rash, urticaria, pruritus, fever, chills, arthralgia, eosinophilia, and, rarely, hepatitis.

Other: nasal congestion, flushing, lacrimation, conjunctivitis.

## **OVERDOSAGE**

### **Acute Toxicity:**

No deaths due to acute poisoning have been reported.

Highest known dose survived: adults, 10 g orally. Oral LD50 in rats: 173 and 187 mg/kg.

**Signs and Symptoms:** Signs and symptoms of overdosage include hypotension, tachycardia, headache, and generalized skin flushing.

Complications can include myocardial ischemia and subsequent myocardial infarction, cardiac arrhythmia, and profound shock.

**Treatment:** There is no specific antidote.

Support of the cardiovascular system is of primary importance. Shock should be treated with plasma expanders. If possible, vasopressors should not be given, but if a vasopressor is required, care should be taken not to precipitate or aggravate cardiac arrhythmia. Tachycardia responds to beta blockers.

Digitalization may be necessary, and renal function should be monitored and supported as required.

No experience has been reported with extracorporeal or peritoneal dialysis.

## **DOSAGE AND ADMINISTRATION**

When there is urgent need, therapy in the hospitalized patient may be initiated intramuscularly or as a rapid intravenous bolus injection directly into the vein. Hydralazine hydrochloride injection should be used only when the drug cannot be given orally. The usual dose is 20 to 40 mg, repeated as necessary.

Certain patients (especially those with marked renal damage) may require a lower dose. Blood pressure should be checked frequently. It may begin to fall within a few minutes after injection, with the average maximal decrease occurring in 10 to 80 minutes. In cases where there has been increased intracranial pressure, lowering the blood pressure may increase cerebral ischemia. Most patients can be transferred to oral hydralazine hydrochloride within 24 to 48 hours.

The product should be used immediately after the vial is opened. It should not be added to infusion solutions. Hydralazine hydrochloride injection may discolor upon contact with metal; discolored solutions should be discarded.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

## HOW SUPPLIED

Hydralazine Hydrochloride Injection USP, 20 mg/mL

NDC 0641-6231-25	1 mL Single-dose Vial	Packaged in boxes of 25
------------------	-----------------------	-------------------------

### Product repackaged by: Henry Schein, Inc., Bastian, VA 24314

From Original Manufacturer/Distributor's NDC and Unit of Sale	To Henry Schein Repackaged Product NDC and Unit of Sale	Total Strength/Total Volume (Concentration) per unit
NDC 0641-6231-25 Packaged in boxes of 25	NDC 0404-9994-01 1 1 mL Single-dose Vial in a bag (Vial bears NDC 0641-6231- 01)	20 mg/mL

**Storage:** Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

### Manufactured by:

Hikma Pharmaceuticals USA Inc.

Berkeley Heights, NJ 07922

December 2020

### Sample Package Label

# HYDRALAZINE HYDROCHLORIDE

20 mg/mL  
1 mL

Injection, USP  
Single Dose Vial

For Intramuscular or Intravenous Use.  
Preservative – Free.  
Warning: Discard Unused Portion.  
Sterile, Nonpyrogenic.  
For emergency use only in patients unable to take oral medications.

Keep out of children's reach.

Store at 20 to 25C (68 to 77F) (see USP Controlled Room Temperature).

NDC:



0404-9994-01

ITEM# :2580455  
LOT# XXXXXXXXX  
EXP: mm – yy

SEE MANUFACTURER'S INSERT  
FOR COMPLETE PRODUCT AND  
PRESCRIBING INFORMATION

Packaged By  
Henry Schein, Inc.  
80 Summit View Lane  
Bastian, VA 24314

MANUFACTURER INFORMATION

Mfr: Hikma

ORIG MFG LOT: XX – XXX – XX  
NDC: 0641 – 6231 – 25



GTIN:(01)XXXXXXXXXXXXXXXXXX  
SER:(21)XXXXXXXXXXXXXXXXXX  
LOT:(10)XXXXXX  
EXP:(17)XXXXXX

RX ONLY

## HYDRALAZINE HYDROCHLORIDE

hydralazine hydrochloride injection, solution

### Product Information

Product Type

HUMAN PRESCRIPTION DRUG

Item Code  
(Source)

NDC:0404-9994(NDC:0641-6231)

Route of Administration

INTRAMUSCULAR,  
INTRAVENOUS

### Active Ingredient/Active Moiety

Ingredient Name

Basis of Strength

Strength

<b>HYDRALAZINE HYDROCHLORIDE</b> (UNII: FD171B778Y) (HYDRALAZINE - UNII:26NAK24LS8)	HYDRALAZINE HYDROCHLORIDE	20 mg in 1 mL
---	---------------------------	---------------

<b>Inactive Ingredients</b>	
<b>Ingredient Name</b>	<b>Strength</b>
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	103.6 mg in 1 mL
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>HYDROCHLORIC ACID</b> (UNII: QTT17582CB)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	

<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:0404-9994-01	1 in 1 BAG	07/17/2023	
1		1 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

<b>Marketing Information</b>			
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA213667	07/17/2023	

**Labeler** - Henry Schein, Inc. (012430880)

Revised: 7/2023

Henry Schein, Inc.