

OXICONAZOLE NITRATE- oxiconazole nitrate cream
E. Fougera & Co. a division of Fougera Pharmaceuticals Inc.

OXICONAZOLE NITRATE CREAM, 1%*

***Potency expressed as oxiconazole**


Rx only

FOR TOPICAL DERMATOLOGIC USE ONLY – NOT FOR OPHTHALMIC OR INTRAVAGINAL USE.

DESCRIPTION

Oxiconazole nitrate cream, 1% contains the antifungal active compound oxiconazole nitrate. This formulation is for topical dermatologic use only.

Chemically, oxiconazole nitrate is 2',4'-dichloro-2-imidazol-1-ylacetophenone (Z)-[O-(2,4-dichlorobenzyl)oxime], mononitrate. The compound has the molecular formula $C_{18}H_{13}ON_3Cl_4 \cdot HNO_3$, a molecular weight of 492.15, and the following structural formula:



Oxiconazole nitrate is a nearly white crystalline powder, soluble in methanol; sparingly soluble in ethanol, chloroform, and acetone; and very slightly soluble in water.

Oxiconazole nitrate cream contains 10 mg of oxiconazole per gram of cream in a white to off-white, opaque cream base of purified water USP, white petrolatum USP, stearyl alcohol NF, propylene glycol USP, polysorbate 60 NF, cetyl alcohol NF, and benzoic acid USP, 0.2% as a preservative.

CLINICAL PHARMACOLOGY

Pharmacokinetics: The penetration of oxiconazole nitrate into different layers of the skin was assessed using an *in vitro* permeation technique with human skin. Five hours after application of 2.5 mg/cm² of oxiconazole nitrate cream onto human skin, the concentration of oxiconazole nitrate was demonstrated to be 16.2 μmol in the epidermis, 3.64 μmol in the upper corium, and 1.29 μmol in the deeper corium. Systemic absorption of oxiconazole nitrate is low. Using radiolabeled drug, less than 0.3% of the applied dose of oxiconazole nitrate was recovered in the urine of volunteer subjects up to 5 days after application of the cream formulation.

Neither *in vitro* nor *in vivo* studies have been conducted to establish relative activity between the lotion and cream formulations.

Microbiology: Oxiconazole nitrate is an imidazole derivative whose antifungal activity is derived

primarily from the inhibition of ergosterol biosynthesis, which is critical for cellular membrane integrity. It has *in vitro* activity against a wide range of pathogenic fungi.

Oxiconazole has been shown to be active against most strains of the following organisms both *in vitro* and in clinical infections at indicated body sites (see **INDICATIONS AND USAGE**):

Epidermophyton floccosum
Trichophyton mentagrophytes
Trichophyton rubrum
Malassezia furfur

The following *in vitro* data are available; **however, their clinical significance is unknown.** Oxiconazole exhibits satisfactory *in vitro* minimum inhibitory concentrations (MICs) against most strains of the following organisms; however, the safety and efficacy of oxiconazole in treating clinical infections due to these organisms have not been established in adequate and well-controlled clinical trials:

Candida albicans
Microsporum audouini
Microsporum canis
Microsporum gypseum
Trichophyton tonsurans
Trichophyton violaceum

INDICATIONS AND USAGE

Oxiconazole nitrate cream is indicated for the topical treatment of the following dermal infections: tinea pedis, tinea cruris, and tinea corporis due to *Trichophyton rubrum*, *Trichophyton mentagrophytes*, or *Epidermophyton floccosum*. Oxiconazole nitrate cream is indicated for the topical treatment of tinea (pityriasis) versicolor due to *Malassezia furfur* (see **DOSAGE AND ADMINISTRATION** and **CLINICAL STUDIES**).

Oxiconazole nitrate cream may be used in pediatric patients for tinea corporis, tinea cruris, tinea pedis, and tinea (pityriasis) versicolor; however, these indications for which oxiconazole nitrate cream has been shown to be effective rarely occur in children below the age of 12.

CONTRAINDICATIONS

Oxiconazole nitrate cream is contraindicated in individuals who have shown hypersensitivity to any of their components.

WARNINGS

Oxiconazole nitrate cream, 1% is not for ophthalmic or intravaginal use.

PRECAUTIONS

General: Oxiconazole nitrate cream is for external dermal use only. Avoid introduction of oxiconazole nitrate cream into the eyes or vagina. If a reaction suggesting sensitivity or chemical irritation should occur with the use of oxiconazole nitrate cream, treatment should be discontinued and appropriate therapy instituted. If signs of epidermal irritation should occur, the drug should be discontinued.

Information for Patients: The patient should be instructed to:

1. Use oxiconazole nitrate as directed by the physician. The hands should be washed after applying the medication to the affected area(s). Avoid contact with the eyes, nose, mouth, and other mucous membranes. Oxiconazole nitrate is for external use only.
2. Use the medication for the **full** treatment time recommended by the physician, even though symptoms may have improved. Notify the physician if there is no improvement after 2 to 4 weeks, or sooner if the condition worsens (see below).
3. Inform the physician if the area of application shows signs of increased irritation, itching, burning, blistering, swelling, or oozing.
4. Avoid the use of occlusive dressings unless otherwise directed by the physician.
5. Do not use this medication for any disorder other than that for which it was prescribed.

Drug Interactions: Potential drug interactions between oxiconazole nitrate and other drugs have not been systematically evaluated.

Carcinogenesis, Mutagenesis, Impairment of Fertility: Although no long-term studies in animals have been performed to evaluate carcinogenic potential, no evidence of mutagenic effect was found in 2 mutation assays (Ames test and Chinese hamster V79 *in vitro* cell mutation assay) or in 2 cytogenetic assays (human peripheral blood lymphocyte *in vitro* chromosome aberration assay and *in vivo* micronucleus assay in mice).

Reproductive studies revealed no impairment of fertility in rats at oral doses of 3 mg/kg/day in females (1 time the human dose based on mg/m²) and 15 mg/kg/day in males (4 times the human dose based on mg/m²). However, at doses above this level, the following effects were observed: a reduction in the fertility parameters of males and females, a reduction in the number of sperm in vaginal smears, extended estrous cycle, and a decrease in mating frequency.

Pregnancy: Teratogenic Effects: Pregnancy Category B.

Reproduction studies have been performed in rabbits, rats, and mice at oral doses up to 100, 150, and 200 mg/kg/day (57, 40, and 27 times the human dose based on mg/m²), respectively, and revealed no evidence of harm to the fetus due to oxiconazole nitrate. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers: Because oxiconazole is excreted in human milk, caution should be exercised when the drug is administered to a nursing woman.

Pediatric Use: Oxiconazole nitrate cream may be used in pediatric patients for tinea corporis, tinea cruris, tinea pedis, and tinea (pityriasis) versicolor; however, these indications for which oxiconazole nitrate cream has been shown to be effective rarely occur in children below the age of 12.

Geriatric Use: A limited number of patients at or above 60 years of age (n ~ 396) have been treated with oxiconazole nitrate cream in US and non-US clinical trials. The number of patients is too small to permit separate analysis of efficacy and safety. The adverse reactions reported with oxiconazole nitrate cream in this population were similar to those reported by younger patients. Based on available data, no adjustment of dosage of oxiconazole nitrate cream in geriatric patients is warranted.

ADVERSE REACTIONS

During clinical trials, of 955 patients treated with oxiconazole nitrate cream, 1%, 41 (4.3%) reported adverse reactions thought to be related to drug therapy. These reactions included pruritus (1.6%); burning (1.4%); irritation and allergic contact dermatitis (0.4% each); folliculitis (0.3%); erythema (0.2%); and papules, fissure, maceration, rash, stinging, and nodules (0.1% each).

OVERDOSAGE

When 5% oxiconazole cream (5 times the concentration of the marketed product) was applied at a rate of 1 g/kg to approximately 10% of body surface area of a group of 40 male and female rats for 35 days, 3 deaths and severe dermal inflammation were reported. No overdoses in humans have been reported with use of oxiconazole nitrate cream.

DOSAGE AND ADMINISTRATION

Oxiconazole nitrate cream should be applied to affected and immediately surrounding areas once to twice daily in patients with tinea pedis, tinea corporis, or tinea cruris. Oxiconazole nitrate cream should be applied once daily in the treatment of tinea (pityriasis) versicolor. Tinea corporis, tinea cruris, and tinea (pityriasis) versicolor should be treated for 2 weeks and tinea pedis for 1 month to reduce the possibility of recurrence. If a patient shows no clinical improvement after the treatment period, the diagnosis should be reviewed.

Note: Tinea (pityriasis) versicolor may give rise to hyperpigmented or hypopigmented patches on the trunk that may extend to the neck, arms, and upper thighs. Treatment of the infection may not immediately result in restoration of pigment to the affected sites. Normalization of pigment following successful therapy is variable and may take months, depending on individual skin type and incidental sun exposure. Although tinea (pityriasis) versicolor is not contagious, it may recur because the organism that causes the disease is part of the normal skin flora.

CLINICAL STUDIES

The following definitions were applied to the clinical and microbiological outcomes in patients enrolled in the clinical trials that form the basis for the approval of oxiconazole nitrate cream.

Definitions:

1. Mycological Cure: No evidence (culture and KOH preparation) of the baseline (original) pathogen in a specimen from the affected area taken at the 2-week post-treatment visit (for tinea [pityriasis] versicolor, mycological cure was limited to KOH only).
2. Treatment Success: Both a global evaluation of 90% clinical improvement and a microbiologic eradication (see above) at the 2-week post-treatment visit.

Tinea Pedis: THERE ARE NO HEAD-TO-HEAD COMPARISON TRIALS OF THE OXICONAZOLE NITRATE CREAM AND LOTION FORMULATIONS IN THE TREATMENT OF TINEA PEDIS.

The two pivotal trials for the oxiconazole nitrate cream formulation involved 281 evaluable patients (total from both trials) with clinically and microbiologically established tinea pedis.

The combined results of these 2 clinical trials at the 2-week post-treatment follow-up visit are shown in the following table:

Patient Outcome	Oxiconazole Nitrate Cream		Vehicle
	b.i.d.	q.d.	
Mycological cure	77%	79%	33%
Treatment success	52%	43%	14%

All the improvement and cure rates of the b.i.d.- and q.d.- treated groups did not differ significantly (95% confidence interval) from each other but were statistically (95% confidence interval) superior to the vehicle-treated group.

In addition, pediatric data (95 children ages 10 and under) available with the cream formulation indicate that it is safe and effective for use in children when used as directed. Adverse events were reported in 2

children; 1 child was reported to have reddening of the skin and 1 child was reported to have eczema-like skin alterations.

Tinea (pityriasis) Versicolor: Two pivotal clinical trials of oxiconazole nitrate cream in tinea (pityriasis) versicolor involved 219 evaluable patients in the q day oxiconazole nitrate and vehicle arms of the trial with clinical and mycological evidence of tinea (pityriasis) versicolor. Patients were treated for 2 weeks with oxiconazole nitrate cream once daily, or with cream vehicle. The combined results of these clinical trials at the 2-week post-treatment follow-up visit are shown in the following table. These results are based on 207 patients (110 in the oxiconazole nitrate group and 97 in the vehicle group) with efficacy evaluations at this visit.

Patient Outcome	Oxiconazole Nitrate Cream	Vehicle
	q.d.	
Mycological cure	88%	67%
Treatment success	83%	62%

Only once a day was shown in both studies to be statistically superior to vehicle for all efficacy parameters at 2 weeks and follow-up.

HOW SUPPLIED

Oxiconazole nitrate cream, 1% is supplied in:

30 g tubes NDC 0168-0358-30,
60 g tubes NDC 0168-0358-60,
90 g tubes NDC 0168-0358-90.

Store between 15° and 30°C (59° and 86°F).

E. FOUGERA & CO.

A division of

Fougera

PHARMACEUTICALS INC.

Melville, New York 11747

46144436A

R09/14

#316

PACKAGE LABEL – PRINCIPAL DISPLAY PANEL – 30 G CONTAINER

NDC 0168-0358-30

OXICONAZOLE

NITRATE CREAM, 1% *

*Potency expressed as oxiconazole.

**For Topical Dermatologic Use ONLY -
Not for Ophthalmic or Intravaginal Use.**

Rx Only

NET WT 30 grams

NDC 0168-0358-30

**OXICONAZOLE
NITRATE CREAM, 1%***

*Potency expressed as oxiconazole.

**For Topical Dermatologic Use ONLY –
Not for Ophthalmic or Intravaginal Use.**

fougera®

**WARNING: Keep out of reach
of children.**

See package insert for full
prescribing information.

Important: Do not use if seal has
been punctured or is not visible.

To Open: Use cap to
puncture seal.

E. FOUGERA & CO.
A division of
Fougera Pharmaceuticals Inc.
Melville, New York 11747

R only

Contains: Oxiconazole 1% as
oxiconazole nitrate in a cream
base composed of purified water
USP, white petrolatum USP,
stearyl alcohol NF, propylene
glycol USP, polysorbate 60 NF,
cetyl alcohol NF, and benzoic
acid 0.2% as a preservative.

NET WT 30 grams

Store between 15° and 30°C
(59° and 86°F).

See crimp of tube for
lot no. and expiration date.

C5405B R09/14



PACKAGE LABEL – PRINCIPAL DISPLAY PANEL – 30 G CARTON

NDC 0168-0358-30


Rx Only

OXICONAZOLE NITRATE CREAM, 1%*

*Potency expressed as oxiconazole.

**For Topical Dermatologic Use ONLY -
Not for Ophthalmic or Intravaginal Use.**

NET WT 30 grams



OXICONAZOLE NITRATE

oxiconazole nitrate cream

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0168-0358
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OXICONAZOLE NITRATE (UNII: RQ8UL4C17S) (OXICONAZOLE - UNII:C668Q9I33J)	OXICONAZOLE	10 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
PETROLATUM (UNII: 4T6H12BN9U)	
STEARYL ALCOHOL (UNII: 2KR89I4HIY)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POLYSORBATE 60 (UNII: CAL22UVI4M)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
BENZOIC ACID (UNII: 8SKN0B0MIM)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0168-0358-30	30 g in 1 TUBE; Type 0: Not a Combination Product	03/09/2016	
2	NDC:0168-0358-60	60 g in 1 TUBE; Type 0: Not a Combination Product	03/09/2016	
3	NDC:0168-0358-90	90 g in 1 TUBE; Type 0: Not a Combination Product	03/10/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA authorized generic	NDA019828	03/09/2016	

Labeler - E. Fougera & Co. a division of Fougera Pharmaceuticals Inc. (043838424)

Revised: 9/2020

E. Fougera & Co. a division of Fougera Pharmaceuticals Inc.