

**MEIJER MAX. STRENGTH ANTI-ITCH CREAM INTENSIVE HEALING FORMULA-
hydrocortisone cream
MEIJER, INC.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Meijer Max. Strength Anti-Itch Cream Intensive Healing Formula 2oz ZDP

Active Ingredient Purpose

Hydrocortisone 1%.....Anti-itch

Uses

temporarily relieves itching associated with minor skin irritations, inflammation, and rashes due to:

- eczema
- psoriasis
- jewelry
- insect bites
- soaps
- cosmetics
- detergents
- seborrheic dermatitis
- poison ivy, oak sumac

temporarily relieves external anal and genital itching

other uses of this product should only be under the advice and supervision of a doctor

Warnings

For external use only

Do not use

- in the genital area if you have a vaginal discharge. Consult a doctor.
- for the treatment of diaper rash. Consult a doctor.
- more than directed unless directed by a doctor

When using this product

- avoid contact with eyes
- do not put directly into the rectum by using fingers or any mechanical device or applicator

Stop use and ask a doctor if

- condition worsens, symptoms persist for more than 7 days or clear up and occur again within a few days, and do not begin use of any other hydrocortisone product unless you have asked a doctor
- rectal bleeding occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

for itching of skin irritation, inflammation, and rashes:

- adults and children 2 years of age and older: apply to affected area not more than 3 to 4 times daily
- children under 2 years of age: ask a doctor

for external anal and genital itching, adults:

- when practical, clean the affected area with mild soap and warm water and rinse thoroughly
- gently dry by patting or blotting with toilet tissue or a soft cloth before applying
- apply to affected area not more than 3 to 4 times daily
- children under 12 years of age: ask a doctor

Other Information

- store at room temperature 20-25°C (68-77°F)

Inactive ingredients aloe vera leaf, .alpha.-tocopherol acetate, benzyl alcohol, cetyl alcohol, dimethicone 350, edetate disodium, gelatin hydrolysate, glycerin, glyceryl stearate/peg-100 stearate, magnesium ascorbyl phosphate, matricaria chamomilla flowering top, menthyl lactate, methyl gluceth-20, methylparaben, pretrolatum, polysorbate 60, propylene glycol, propylparaben, purified water, steareth-2, steareth-21, stearyl alcohol, vitamin a palmitate

MAXIMUM STRENGTH

anti-itch cream

Hydrocortisone 1% (Anti-itch) | Vitamins A, C and E

Intensive Healing Formula with Chamomile and Antioxidants

ECZEMA | PSORIASIS | RASHES | ITCHY SKIN

Ultra creamy formula helps heal problem skin.
Calms, nourishes, hydrates.
Safe for sensitive skin.

meijer

Compare to Cortizone-10® Intensive Healing Anti-Itch Creme active ingredient*

MAXIMUM STRENGTH

anti-itch cream

Hydrocortisone 1% (Anti-itch) | Vitamins A, C and E

Intensive Healing Formula with Chamomile and Antioxidants

ECZEMA | PSORIASIS | RASHES | ITCHY SKIN

NET WT 2 OZ (57 g)

MAXIMUM STRENGTH

anti-itch cream

Hydrocortisone 1% (Anti-itch) | Vitamins A, C and E



DIST. BY MEIJER DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com

Made in China

LOT & EXP.

40312-Z0P-8X-81
0M-ND-Z0P1-20-81

*This product is not manufactured or distributed by Chatham, Inc., owner of the registered trademark Cortizone-10®.

Questions or comments?
Call 1-888-645-8244

Inactive ingredients
store at room temperature 20-25°C (68-77°F)
water, benzyl alcohol, cetyl alcohol, dimethicone, SS1, oleate, disodium, glycerin, glyceryl stearate, PEG-100, stearate, magnesium ascorbyl phosphate, methoxy glactin-20, flower, top, menthyl lactate, polyisobutyl acrylate, propylene glycol, propylparaben, perubalium, polysorbate 60, propylene glycol, alcohol, vitamin E palmitate

Other information
■ store at room temperature 20-25°C (68-77°F)
■ children under 12 years of age: ask a doctor

Directions
■ for itching of skin irritation, inflammation, and rashes.
■ adults and children 2 years of age and older: apply to affected area not more than 3 to 4 times daily
■ children under 2 years of age: ask a doctor
■ for external anal and genital itching, adults:
■ when practical, clean the affected area with mild soap and gently dry by patting or blotting with toilet tissue or a soft cloth before applying
■ apply to affected area not more than 3 to 4 times daily

Warnings
For external use only
Do not use
■ in the genital area if you have a vaginal discharge. Consult a doctor.
■ for the treatment of diaper rash. Consult a doctor.
■ more than directed unless directed by a doctor
When using this product
■ avoid contact with eyes
■ do not put directly into the rectum by using fingers or any mechanical device or applicator
Stop use and ask a doctor if
■ condition worsens, symptoms persist for more than 7 days or clear up and occur again within a few days, and do not begin use of any other hydrocortisone product unless you have asked a doctor
■ rectal bleeding occurs
If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)
Purpose
Anti-itch
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Drug Facts
Temporarily relieves itching associated with minor skin irritations, inflammations, and rashes due to:
■ eczema ■ insect bites
■ allergic reactions ■ jewelry
■ cosmetics ■ deodorants
■ soaps ■ poisons (ivy, oak, sumac)
■ atrophic dermatitis
■ other uses of this product should only be under the advice and supervision of a doctor
Warnings
■ temporarily relieves external anal and genital itching
■ other uses of this product should only be under the advice and supervision of a doctor
Do not use
■ in the genital area if you have a vaginal discharge. Consult a doctor.
■ for the treatment of diaper rash. Consult a doctor.
■ more than directed unless directed by a doctor
When using this product
■ avoid contact with eyes
■ do not put directly into the rectum by using fingers or any mechanical device or applicator
Stop use and ask a doctor if
■ condition worsens, symptoms persist for more than 7 days or clear up and occur again within a few days, and do not begin use of any other hydrocortisone product unless you have asked a doctor
■ rectal bleeding occurs
If pregnant or breast-feeding, ask a health professional before use.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

TAMPERS EVIDENT: DO NOT USE IF CARTON IS OPEN OR SEAL UNDER CAP IS BROKEN OR MISSING.

MEIJER MAX. STRENGTH ANTI-ITCH CREAM INTENSIVE HEALING FORMULA
hydrocortisone cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-259
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROCORTISONE (UNII: W4X0X7BPJ) (HYDROCORTISONE - UNII:W4X0X7BPJ)	HYDROCORTISONE	10 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
ALOE VERA LEAF (UNII: ZY81Z83H0X)	
MAGNESIUM ASCORBYL PHOSPHATE (UNII: 0R822556M5)	
MENTHYL LACTATE, (-)- (UNII: 2BF9E65L7I)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POLYSORBATE 60 (UNII: CAL22UVI4M)	
STEARETH-21 (UNII: 53J3F32P58)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
VITAMIN A PALMITATE (UNII: 1D1K0N0VVC)	
DIMETHICONE 350 (UNII: 2Y53S6ATLU)	
GELATIN HYDROLYSATE (PORCINE SKIN, MW 3000) (UNII: 0K9R94573C)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
GLYCERYL STEARATE/PEG-100 STEARATE (UNII: RD25J5V947)	
METHYL GLUCETH-20 (UNII: J3QD0LD11P)	
.ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)	
GLYCERIN (UNII: PDC6A3C0OX)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	
PETROLATUM (UNII: 4T6H12BN9U)	
STEARETH-2 (UNII: V56DFE46J5)	
STEARYL ALCOHOL (UNII: 2KR89I4H1Y)	
MATRICARIA CHAMOMILLA FLOWERING TOP (UNII: 3VNC7T6Z02)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-259-57	1 in 1 CARTON	07/16/2020	
1		57 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
---------------------------	---	-----------------------------	---------------------------

OTC monograph not final	part348	07/16/2020	
-------------------------	---------	------------	--

Labeler - MEIJER, INC. (006959555)

Revised: 3/2022

MEIJER, INC.