

**PAIN RELIEF PLUS MULTI SYMPTOM COLD CHILDRENS- acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, phenylephrine hcl liquid P & L Development, LLC**

-----

**Drug Facts**

**Active ingredients (in each 5 mL)**

**Acetaminophen 160 mg**

Chlorpheniramine maleate 1 mg

Dextromethorphan HBr 5 mg

Phenylephrine HCl 2.5 mg

**Purpose**

**Pain reliever/fever reducer**

Antihistamine

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves the following cold/flu symptoms:
  - minor aches and pains
  - sore throat
  - cough
  - sneezing and runny nose
  - headache
  - nasal congestion
  - stuffy nose
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child take:

- more than 5 doses (10 mL) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- to make a child sleepy
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

### **Ask a doctor before use if the child has**

- liver disease
- glaucoma
- thyroid disease
- diabetes
- heart disease
- high blood pressure
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

### **Ask a doctor or pharmacist before use if the child is taking**

- sedatives or tranquilizers
- the blood thinning drug warfarin

### **When using this product**

- **do not exceed recommended dose (see overdose warning)**
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- nervousness, dizziness or sleeplessness occurs
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

### Directions

- **this product does not contain directions or complete warnings for adult use**
- mL=milliliter
- shake well before using
- do not give more than 5 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- if needed, repeat dose every 4 hours while symptoms last
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age

Weight (lb)	Age (yr)	Dose (mL)
48-95	6-11	10 mL
36-47	4-5	do not use unless directed by a doctor
under 36	under 4	do not use

### Other information

- store between 20-25°C (68-77°F). Do not refrigerate

### Inactive ingredients

carboxymethylcellulose sodium, microcrystalline cellulose, anhydrous citric acid, D&C red 33, FD&C blue 1, FD&C red 40, flavors, glycerin, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum

### Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

### Principal Display Panel

Compare to the active ingredient in **Children's Tylenol® Plus Multi-Symptom Cold\***

#### Children's

pain relief plus multi-symptom cold

**Acetaminophen 160 mg**

Chlorpheniramine Maleate 1 mg

Dextromethorphan HBr 5 mg

Phenylephrine HCl 2.5 mg

for ages 6-11 years

oral suspension

Relieves:

- Fever
- Sore Throat
- Runny Nose
- Sneezing
- Cough
- Stuffy Nose
- Nasal Congestion

grape flavor

FL OZ (mL)

\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Plus Multi-Symptom Cold.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

Manufactured by:

**PL Developments**

11865 S. Alameda St

Lynwood, CA 90262

**Package Label**

**Drug Facts (continued)**

- nervousness, dizziness or sleeplessness occurs
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than 5 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

Weight (lb)	Age (yr)	Dose (mL)
48-95	6-11	10 mL
36-47	4-5	do not use unless directed by a doctor
under 36	under 4	do not use

**Other information**

- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** carboxymethylcellulose sodium, citric acid, D&C red #33, FD&C blue #1, FD&C red #40, flavor, glycerin, microcrystalline cellulose, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum

**Questions or comments?**  
Call 1-877-753-3838 Monday-Friday 9AM-5PM EST

\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Plus Multi-Symptom Cold.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**PARENTS:**  
Learn about this medicine online  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**TRUST QUALITY EXCELLENCE**

Manufactured by:  
**PL Developments**  
11865 S. Alameda St  
Lynwood, CA 90262

Compare to the active ingredients in  
**Children's Tylenol® Plus Multi-Symptom Cold\***  
NDC 49580-0304-4

**children's**  
**pain relief plus**  
**multi-symptom**  
**cold**

**Acetaminophen** 160 mg  
chlorpheniramine maleate 1 mg  
dextromethorphan HBr 5 mg  
phenylephrine HCl 2.5 mg

for ages 6-11 years  
oral suspension

relieves:

- fever
- runny nose
- cough
- nasal congestion

grape flavor

**4 fl oz (118 mL)**

**children's**  
**pain relief plus**  
**multi-symptom**  
**cold**

**Acetaminophen** 160 mg  
chlorpheniramine maleate 1 mg  
dextromethorphan HBr 5 mg  
phenylephrine HCl 2.5 mg

for ages 6-11 years  
oral suspension

relieves:

- fever
- runny nose
- cough
- nasal congestion

grape flavor

**4 fl oz (118 mL)**

3 49580 30404 8

PLD-A253C FC007674

Lot No.:

Exp. Date:

**Drug Facts**

**Active ingredients (in each 5 mL)**

Active ingredients (in each 5 mL)	Purposes
Acetaminophen 160 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 1 mg	Antihistamine
Dextromethorphan HBr 5 mg	Cough suppressant
Phenylephrine HCl 2.5 mg	Nasal decongestant

**Uses**

- temporarily relieves the following cold/flu symptoms
- minor aches and pains
- sore throat
- sneezing and runny nose
- headache
- nasal congestion
- cough
- stuffy nose
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses (10 mL) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- to make a child sleepy
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if your child has**

- liver disease
- glaucoma
- thyroid disease
- diabetes
- heart disease
- high blood pressure
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

**Ask a doctor or pharmacist before use if your child is taking**

- the blood thinning drug warfarin
- sedatives or tranquilizers

**When using this product**

- do not exceed recommended dose (see Overdose warning)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

## READYinCASE Children's Pain Relief Multi-Symptom Cold

## PAIN RELIEF PLUS MULTI SYMPTOM COLD CHILDRENS

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, phenylephrine hcl liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49580-0304
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	1 mg in 5 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH)	DEXTROMETHORPHAN	5 mg

(DEXTROMETHORPHAN - UNII:7355X3ROTS)		HYDROBROMIDE	in 5 mL	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)		PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL	
Inactive Ingredients				
Ingredient Name			Strength	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
D&C RED NO. 33 (UNII: 9DBA0SBB0L)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
GLYCERIN (UNII: PDC6A3C0OX)				
WATER (UNII: 059QF0KO0R)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
SORBITOL (UNII: 506T60A25R)				
SUCROSE (UNII: C151H8M554)				
XANTHAN GUM (UNII: TTV12P4NEE)				
Product Characteristics				
Color		Score		
Shape		Size		
Flavor	GRAPE	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49580-0304-4	1 in 1 BOX	03/26/2021	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M012	03/26/2021	

**Labeler** - P & L Development, LLC (101896231)