

ANTI DIARRHEAL- loperamide hcl tablet
L.N.K. International, Inc.

Sound Body 44-375-Delisted

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

Heart alert: Taking more than directed can cause serious heart problems or death.

Do not use

if you have bloody or black stool.

Ask a doctor before use if you have

- a fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product,

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store between 20°-25°C (68°-77°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

Questions or comments?

Call **1-800-426-9391** 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

SOUNDBODY™

***Compare to the active ingredient in Imodium® A-D**

NDC 50844-753-08

Anti-Diarrheal

Loperamide HCl, 2 mg

Anti-Diarrheal

Controls the Symptoms of Diarrhea

24 CAPLETS

Actual Size

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Imodium® A-D.

50844 REV0619F37508

Manufactured for Big Lots Stores, Inc.

by **LNK INTERNATIONAL, INC.**

60 Arkay Drive, Hauppauge, NY 11788 USA

V#733000 ITEM#022737508BLBX

Drug Facts
Active ingredient (in each caplet)
Loperamide HCl 2 mg
Anti-diarrheal

Use
controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

SOUNDBODY™

*Compare to the active ingredient in Imodium® A-D

NDC 50844-753-08

Anti-Diarrheal
Loperamide HCl, 2 mg
Anti-Diarrheal

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

No print/No varnish
Lot & Exp date

3
508




Controls the symptoms of diarrhea

24 CAPLETS



Actual Size



B-0227-375-08-R
REV0619F37508



2. Use \gg to cut through plastic for caplet.



1. To open, tear at perforations and remove a section.

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark, Imodium[®] A-D.
50844 REV0619F37508
Manufactured for Big Lots Stores, Inc.
by LNK INTERNATIONAL, INC.
60 Arkay Drive, Hauppauge, NY 11788 USA
V#733000 ITEM#0227375088LBX

Questions or comments? Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Inactive ingredients FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide, corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate.

Other information
 ■ store between 20°-25°C (68°-77°F) ■ see end flap for expiration date and lot number
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Directions ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea. ■ find right dose on chart. If possible, use weight to dose; otherwise, use age.

Drug Facts (continued)
 Heart alert: Taking more than directed can cause serious heart problems or death.
 Do not use if you have bloody or black stool.
 Ask a doctor before use if you have ■ a fever ■ mucus in the stool ■ a history of liver disease ■ a history of abnormal heart rhythm
 Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.
 When using this product, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.
 Stop use and ask a doctor if ■ symptoms get worse ■ diarrhea lasts for more than 2 days ■ you get abdominal swelling or bloating. These may be signs of a serious condition.
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Sound Body 44-375

ANTI DIARRHEAL

loperamide hcl tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50844-753
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	green	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	44;375
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-753-08	4 in 1 CARTON	05/03/2005	11/21/2025
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:50844-753-02	2 in 1 CARTON	05/03/2005	08/31/2019
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076497	05/03/2005	11/21/2025

Labeler - L.N.K. International, Inc. (038154464)**Establishment**

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

LNK International, Inc.		832867837	manufacture(50844-753) , pack(50844-753)
-------------------------	--	-----------	--

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(50844-753)

Revised: 6/2024

L.N.K. International, Inc.