

MILK OF MAGNESIA- magnesium hydroxide suspension

H E B

heb mom CHRY

Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are taking a prescription drug.

This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a

Poison Control Center right away.

Directions

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose

age	dose
adults and children 12 years and over	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

Other information

- each tablespoonful contains: **magnesium 500 mg, sodium 6 mg**
- avoid freezing
- store at room temperature tightly closed

Inactive ingredients

carboxymethylcellulose sodium, citric acid, D & C red no. 28, flavor, glycerin, microcrystalline cellulose, purified water, sodium citrate, sodium hypochlorite, sucrose, xanthan gum

Questions or comments?

1-800-540-3765

package Label

Compare to **Phillips'**® Milk of Magnesia
active ingredient*

NDC 37808-064-12

H-E-B®

Cherry

Milk of Magnesia

Magnesium Hydroxide

Saline Laxative

- Cramp Free
- Stimulant Free



12 FL OZ (355 mL)

282-19312-0 REV GC0820

Drug Facts

TAMPER EVIDENT: DO NOT USE IF IMPRINTED NECKBAND IS MISSING OR BROKEN.

Active ingredient (in each 15 mL)	Purpose
Magnesium hydroxide 1200 mg	Saline laxative

Use

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- a magnesium-restricted diet
- a sudden change in bowel habits that lasts over 14 days
- kidney disease
- stomach pain, nausea or vomiting

Ask a doctor or pharmacist before use if you are taking a prescription drug.
This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breastfeeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose

age	dose
adults and children 12 years and over	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

Other information

- each 15 mL contains: magnesium 500 mg, sodium 6 mg
- avoid freezing
- store at room temperature tightly closed

Inactive ingredients carboxymethylcellulose sodium, citric acid, D&C red no. 28, flavor, glycerin, microcrystalline cellulose, purified water, sodium citrate, sodium hypochlorite, sucrose, xanthan gum

Questions or comments? 1-800-540-3765



MADE WITH PRIDE & CARE FOR H-E-B®
SAN ANTONIO, TX 78204



*This product is not manufactured or distributed by Bayer Healthcare LLC, the owner of the registered trademark Phillips'® Milk of Magnesia.

16269-2010

982-19312-0 REV GC0820

MILK OF MAGNESIA

magnesium hydroxide suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-064
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
GLYCERIN (UNII: PDC6A3C0OX)	
XANTHAN GUM (UNII: TTV12P4NEE)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
WATER (UNII: 059QF0KO0R)	
SODIUM HYPOCHLORITE (UNII: DY38VHM5OD)	
SUCROSE (UNII: C151H8M554)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-064-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2020	
2	NDC:37808-064-26	769 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	08/01/2020	

Labeler - H E B (007924756)

Registrant - GCP Laboratories (965480861)

Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(37808-064)