

LEVETIRACETAM- levetiracetam tablet, film coated, extended release

Apotex Corp.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use LEVETIRACETAM EXTENDED-RELEASE TABLETS safely and effectively. See full prescribing information for LEVETIRACETAM EXTENDED-RELEASE TABLETS.

LEVETIRACETAM extended-release tablets, for oral use
Initial U.S. Approval: 1999

RECENT MAJOR CHANGES -----

Warnings and Precautions (5.6) 3/2024

INDICATIONS AND USAGE -----

Levetiracetam extended-release tablets are indicated for the treatment of partial-onset seizures in patients 12 years of age and older (1)

DOSAGE AND ADMINISTRATION -----

Initiate treatment with a dose of 1,000 mg once daily; increase by 1,000 mg every 2 weeks to a maximum recommended dose of 3,000 mg once daily (2)

See full prescribing information for use in patients with impaired renal function (2.1)

DOSAGE FORMS AND STRENGTHS -----

- 500 mg white, film-coated extended-release tablet (3)
- 750 mg white, film-coated extended-release tablet (3)
- 1,000 mg white, film-coated extended-release tablet (3)

CONTRAINDICATIONS -----

Known hypersensitivity to levetiracetam; angioedema and anaphylaxis have occurred (4, 5.4)

WARNINGS AND PRECAUTIONS -----

- Behavioral abnormalities including psychotic symptoms, suicidal ideation, irritability, and aggressive behavior have been observed; monitor patients for psychiatric signs and symptoms (5.1)
- Suicidal Behavior and Ideation: Monitor patients for new or worsening depression, suicidal thoughts/behavior, and/or unusual changes in mood or behavior (5.2)
- Monitor for somnolence and fatigue and advise patients not to drive or operate machinery until they have gained sufficient experience on levetiracetam extended-release tablets (5.3)
- Serious Dermatological Reactions: Discontinue levetiracetam at the first sign of rash unless clearly not drug related (5.5)
- Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multiorgan Hypersensitivity: Discontinue if no alternative etiology (5.6)
- Coordination Difficulties: Monitor for ataxia, abnormal gait, and incoordination. Advise patients to not drive or operate machinery until they have gained experience on levetiracetam extended-release tablets (5.7)
- Withdrawal Seizures: Levetiracetam extended-release tablets must be gradually withdrawn (5.8)

ADVERSE REACTIONS -----

Most common adverse reactions (incidence \geq 5% more than placebo) include: somnolence and irritability (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Apotex Corp. at 1-800-706-5575 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

USE IN SPECIFIC POPULATIONS -----

Pregnancy: Plasma levels of levetiracetam may be decreased and therefore need to be monitored closely during pregnancy. Based on animal data, may cause fetal harm (5.10, 8.1)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 10/2024

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosing

2.2 Dosage Adjustments in Adult Patients with Renal Impairment

2.3 Discontinuation of Levetiracetam Extended-Release Tablets

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Behavioral Abnormalities and Psychotic Symptoms

5.2 Suicidal Behavior and Ideation

5.3 Somnolence and Fatigue

5.4 Anaphylaxis and Angioedema

5.5 Serious Dermatological Reactions

5.6 Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multiorgan Hypersensitivity

5.7 Coordination Difficulties

5.8 Withdrawal Seizures

5.9 Hematologic Abnormalities

5.10 Seizure Control During Pregnancy

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Renal Impairment

10 OVERDOSAGE

10.1 Signs, Symptoms and Laboratory Findings of Acute Overdosage in Humans

10.2 Management of Overdose

10.3 Hemodialysis

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

14.1 Levetiracetam Extended-Release Tablets in Adults

14.2 Immediate-Release Levetiracetam Tablets in Adults

14.3 Immediate-Release Levetiracetam Tablets in Pediatric Patients 4 Years to 16 Years

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

16.2 Storage

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Levetiracetam extended-release tablets are indicated for the treatment of partial-onset seizures in patients 12 years of age and older.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosing

For adults and adolescent patients, the recommended dosing for monotherapy and adjunctive therapy is the same; as outlined below.

Adults and Adolescents 12 Years of Age and Older Weighing 50 kg or More

Initiate treatment with a dose of 1,000 mg once daily. The once daily dosage may be adjusted in increments of 1,000 mg every 2 weeks to a maximum recommended daily dose of 3,000 mg/day once daily.

Administration

Levetiracetam extended-release tablets are administered once daily. Levetiracetam extended-release tablets should be swallowed whole. The tablets should not be chewed, broken, or crushed.

2.2 Dosage Adjustments in Adult Patients with Renal Impairment

Levetiracetam extended-release tablets dosing must be individualized according to the patient's renal function status. Recommended dosage adjustments for adults are shown in Table 1. In order to calculate the dose recommended for patients with renal impairment, creatinine clearance adjusted for body surface area must be calculated. To do this, an estimate of the patient's creatinine clearance (CLcr) in mL/min must first be calculated using the following formula:

$$\text{CLcr} = \frac{[140 - \text{age (years)}] \times \text{weight (kg)}}{72 \times \text{serum creatinine (mg/dL)}} \quad (\times 0.85 \text{ for female patients})$$

Then CLcr is adjusted for body surface area (BSA) as follows:

$$\text{CLcr (mL/min/1.73m}^2\text{)} = \frac{\text{CLcr (mL/min)}}{\text{BSA subject (m}^2\text{)}} \times 1.73$$

Table 1: Dosage Adjustment Regimen for Adult Patients with Renal Impairment

Group	Creatinine Clearance(mL/min/1.73m²)	Dosage (mg)	Frequency
Normal	> 80	1,000 to 3,000	Every 24 hours
Mild	50 - 80	1,000 to 2,000	Every 24 hours
Moderate	30 - 50	500 to 1,500	Every 24 hours
Severe	< 30	500 to 1,000	Every 24 hours

2.3 Discontinuation of Levetiracetam Extended-Release Tablets

Avoid abrupt withdrawal from levetiracetam extended-release tablets in order to reduce the risk of increased seizure frequency and status epilepticus [see *Warnings and Precautions (5.8)*].

3 DOSAGE FORMS AND STRENGTHS

Levetiracetam extended-release, USP 500 mg tablets are white, oval, biconvex film-coated tablets, engraved with "APO" on one side, "LXR 500" on the other side.

Levetiracetam extended-release, USP 750 mg tablets are white, capsule shaped, biconvex film-coated tablets, engraved with "APO" on one side, "LXR 750" on the other side.

Levetiracetam extended-release, USP 1,000 mg tablets are white, oval shaped, biconvex film-coated tablets, engraved with "APO" on one side, "LXR 1000" on the other side.

4 CONTRAINDICATIONS

Levetiracetam extended-release tablets are contraindicated in patients with a hypersensitivity to levetiracetam. Reactions have included anaphylaxis and angioedema [see *Warnings and Precautions (5.4)*].

5 WARNINGS AND PRECAUTIONS

5.1 Behavioral Abnormalities and Psychotic Symptoms

Levetiracetam extended-release tablets may cause behavioral abnormalities and psychotic symptoms. Patients treated with levetiracetam extended-release tablets should be monitored for psychiatric signs and symptoms.

Behavioral abnormalities

Levetiracetam Extended-Release Tablets

A total of 7% of levetiracetam extended-release-treated patients experienced non-psychotic behavioral disorders (reported as irritability and aggression) compared to 0% of placebo-treated patients. Irritability was reported in 7% of levetiracetam extended-release-treated patients. Aggression was reported in 1% of levetiracetam extended-release-treated patients.

No patient discontinued treatment or had a dose reduction as a result of these adverse reactions.

The number of patients exposed to levetiracetam extended-release tablets was

considerably smaller than the number of patients exposed to immediate-release levetiracetam tablets in controlled trials. Therefore, certain adverse reactions observed in the immediate-release levetiracetam controlled trials will likely occur in patients receiving levetiracetam extended-release tablets.

Immediate-Release Levetiracetam Tablets

A total of 13% of adult patients and 38% of pediatric patients (4 to 16 years of age) treated with immediate-release levetiracetam tablets experienced non-psychotic behavioral symptoms (reported as aggression, agitation, anger, anxiety, apathy, depersonalization, depression, emotional lability, hostility, hyperkinesias, irritability, nervousness, neurosis, and personality disorder), compared to 6% and 19% of adult and pediatric patients on placebo. A randomized, double-blind, placebo-controlled study was performed to assess the neurocognitive and behavioral effects of immediate-release levetiracetam tablets as adjunctive therapy in pediatric patients (4 to 16 years of age). An exploratory analysis suggested a worsening in aggressive behavior in patients treated with immediate-release levetiracetam tablets in that study [see *Use in Specific Populations (8.4)*].

A total of 1.7% of adult patients treated with immediate-release levetiracetam tablets discontinued treatment due to behavioral adverse reactions, compared to 0.2% of placebo-treated patients. The treatment dose was reduced in 0.8% of adult patients treated with immediate-release levetiracetam, compared to 0.5% of placebo-treated patients. Overall, 11% of pediatric patients treated with immediate-release levetiracetam experienced behavioral symptoms associated with discontinuation or dose reduction, compared to 6.2% of placebo-treated pediatric patients.

One percent of adult patients and 2% of pediatric patients (4 to 16 years of age) treated with immediate-release levetiracetam tablets experienced psychotic symptoms, compared to 0.2% and 2%, respectively, in adult and placebo-treated pediatric patients. In the controlled study that assessed the neurocognitive and behavioral effects of immediate-release levetiracetam tablets in pediatric patients 4 to 16 years of age, 1.6% levetiracetam-treated patients experienced paranoia, compared to no placebo-treated patients. There were 3.1% patients treated with immediate-release levetiracetam who experienced confusional state, compared to no placebo-treated patients [see *Use in Specific Populations (8.4)*].

Psychotic symptoms

Immediate-Release Levetiracetam Tablets

One percent of levetiracetam-treated adult patients experienced psychotic symptoms compared to 0.2% of placebo-treated patients.

Two (0.3%) levetiracetam-treated adult patients were hospitalized and their treatment was discontinued due to psychosis. Both events, reported as psychosis, developed within the first week of treatment and resolved within 1 to 2 weeks following treatment discontinuation. There was no difference between drug and placebo-treated patients in the incidence of pediatric patients who discontinued treatment due to psychotic and non-psychotic adverse reactions.

5.2 Suicidal Behavior and Ideation

Antiepileptic drugs (AEDs), including levetiracetam extended-release tablets, increase the risk of suicidal thoughts or behavior in patients taking these drugs for any indication.

Patients treated with any AED for any indication should be monitored for the emergence or worsening of depression, suicidal thoughts or behavior, and/or any unusual changes in mood or behavior.

Pooled analyses of 199 placebo-controlled clinical trials (mono- and adjunctive therapy) of 11 different AEDs showed that patients randomized to one of the AEDs had approximately twice the risk (adjusted Relative Risk 1.8, 95% CI:1.2, 2.7) of suicidal thinking or behavior compared to patients randomized to placebo. In these trials, which had a median treatment duration of 12 weeks, the estimated incidence rate of suicidal behavior or ideation among 27,863 AED-treated patients was 0.43%, compared to 0.24% among 16,029 placebo-treated patients, representing an increase of approximately one case of suicidal thinking or behavior for every 530 patients treated. There were four suicides in drug-treated patients in the trials and none in placebo-treated patients, but the number is too small to allow any conclusion about drug effect on suicide.

The increased risk of suicidal thoughts or behavior with AEDs was observed as early as one week after starting drug treatment with AEDs and persisted for the duration of treatment assessed. Because most trials included in the analysis did not extend beyond 24 weeks, the risk of suicidal thoughts or behavior beyond 24 weeks could not be assessed.

The risk of suicidal thoughts or behavior was generally consistent among drugs in the data analyzed. The finding of increased risk with AEDs of varying mechanisms of action and across a range of indications suggests that the risk applies to all AEDs used for any indication. The risk did not vary substantially by age (5 to 100 years) in the clinical trials analyzed. Table 2 shows absolute and relative risk by indication for all evaluated AEDs.

Table 2: Risk by Indication for Antiepileptic Drugs in the Pooled Analysis

Indication	Placebo Patients with Events Per 1,000 Patients	Drug Patients with Events Per 1,000 Patients	Relative Risk: Incidence of Events in Drug Patients/Incidence in Placebo Patients	Risk Difference: Additional Drug Patients with Events Per 1,000 Patients
Epilepsy	1.0	3.4	3.5	2.4
Psychiatric	5.7	8.5	1.5	2.9
Other	1.0	1.8	1.9	0.9
Total	2.4	4.3	1.8	1.9

The relative risk for suicidal thoughts or behavior was higher in clinical trials for epilepsy than in clinical trials for psychiatric or other conditions, but the absolute risk differences were similar for the epilepsy and psychiatric indications.

Anyone considering prescribing levetiracetam extended-release tablets or any other AED must balance the risk of suicidal thoughts or behavior with the risk of untreated illness. Epilepsy and many other illnesses for which AEDs are prescribed are themselves associated with morbidity and mortality and an increased risk of suicidal thoughts and

behavior. Should suicidal thoughts and behavior emerge during treatment, the prescriber needs to consider whether the emergence of these symptoms in any given patient may be related to the illness being treated.

5.3 Somnolence and Fatigue

Levetiracetam extended-release tablets may cause somnolence and fatigue. Patients should be monitored for these signs and symptoms and advised not to drive or operate machinery until they have gained sufficient experience on levetiracetam extended-release tablets to gauge whether it adversely affects their ability to drive or operate machinery.

Somnolence

Levetiracetam Extended-Release Tablets

In the levetiracetam extended-release tablets double-blind, controlled trial in patients experiencing partial-onset seizures, 8% of levetiracetam extended-release-treated patients experienced somnolence compared to 3% of placebo-treated patients.

No patient discontinued treatment or had a dose reduction as a result of these adverse reactions.

The number of patients exposed to levetiracetam extended-release tablets was considerably smaller than the number of patients exposed to immediate-release levetiracetam tablets in controlled trials. Therefore, certain adverse reactions observed in the immediate-release levetiracetam controlled trials will likely occur in patients receiving levetiracetam extended-release tablets.

Immediate-Release Levetiracetam Tablets

In controlled trials of adult patients with epilepsy experiencing partial-onset seizures, 15% of levetiracetam-treated patients reported somnolence, compared to 8% of placebo-treated patients. There was no clear dose response up to 3,000 mg/day. In a study where there was no titration, about 45% of patients receiving 4,000 mg/day reported somnolence. The somnolence was considered serious in 0.3% of the levetiracetam-treated patients, compared to 0% in the placebo group. About 3% of levetiracetam-treated patients discontinued treatment due to somnolence, compared to 0.7% of placebo-treated patients. In 1.4% of levetiracetam-treated patients and in 0.9% of placebo-treated patients the dose was reduced, while 0.3% of the treated patients were hospitalized due to somnolence.

Asthenia

Immediate-Release Levetiracetam Tablets

In controlled trials of adult patients with epilepsy experiencing partial-onset seizures, 15% of levetiracetam-treated patients reported asthenia, compared to 9% of placebo-treated patients. Treatment was discontinued due to asthenia in 0.8% of levetiracetam-treated patients as compared to 0.5% of placebo-treated patients. In 0.5% of levetiracetam-treated patients and in 0.2% of placebo-treated patients, the dose was reduced due to asthenia.

Somnolence and asthenia occurred most frequently within the first 4 weeks of treatment.

5.4 Anaphylaxis and Angioedema

Levetiracetam extended-release tablets can cause anaphylaxis or angioedema after the first dose or at any time during treatment. Signs and symptoms in cases reported in the postmarketing setting in patients treated with levetiracetam have included hypotension, hives, rash, respiratory distress, and swelling of the face, lip, mouth, eye, tongue, throat, and feet. In some reported cases, reactions were life-threatening and required emergency treatment. If a patient develops signs or symptoms of anaphylaxis or angioedema, levetiracetam extended-release tablets should be discontinued and the patient should seek immediate medical attention. Levetiracetam extended-release tablets should be discontinued permanently if a clear alternative etiology for the reaction cannot be established [see *Contraindications (4)*].

5.5 Serious Dermatological Reactions

Serious dermatological reactions, including Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN), have been reported in patients treated with levetiracetam. The median time of onset is reported to be 14 to 17 days, but cases have been reported at least four months after initiation of treatment. Recurrence of the serious skin reactions following rechallenge with levetiracetam has also been reported. Levetiracetam extended-release tablets should be discontinued at the first sign of a rash, unless the rash is clearly not drug-related. If signs or symptoms suggest SJS/TEN, use of this drug should not be resumed and alternative therapy should be considered.

5.6 Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multiorgan Hypersensitivity

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS), also known as multiorgan hypersensitivity, has been reported in patients taking antiepileptic drugs, including levetiracetam. These events can be fatal or life-threatening, particularly if diagnosis and treatment do not occur as early as possible. DRESS typically, although not exclusively, presents with fever, rash, lymphadenopathy, and/or facial swelling, in association with other organ system involvement, such as hepatitis, nephritis, hematological abnormalities, myocarditis, or myositis, sometimes resembling an acute viral infection. Eosinophilia is often present. Because this disorder is variable in its expression, other organ systems not noted here may be involved. It is important to note that early manifestations of hypersensitivity, such as fever or lymphadenopathy, may be present even though rash is not evident. If such signs or symptoms are present, the patient should be evaluated immediately. Levetiracetam extended-release tablets should be discontinued if an alternative etiology for the signs or symptoms cannot be established [see *Contraindications (4)*].

5.7 Coordination Difficulties

Coordination difficulties were not observed in the levetiracetam extended-release tablets controlled trial, however, the number of patients exposed to levetiracetam extended-release tablets was considerably smaller than the number of patients exposed to immediate-release levetiracetam tablets in controlled trials. However, adverse reactions observed in the immediate-release levetiracetam tablets controlled trials may also occur in patients receiving levetiracetam extended-release tablets.

Immediate-Release Levetiracetam Tablets

A total of 3.4% of adult levetiracetam-treated patients experienced coordination difficulties, (reported as either ataxia, abnormal gait, or incoordination) compared to

1.6% of placebo-treated patients. A total of 0.4% of patients in controlled trials discontinued levetiracetam treatment due to ataxia, compared to 0% of placebo-treated patients. In 0.7% of levetiracetam-treated patients and in 0.2% of placebo-treated patients, the dose was reduced due to coordination difficulties, while one of the levetiracetam-treated patients was hospitalized due to worsening of pre-existing ataxia. These events occurred most frequently within the first 4 weeks of treatment.

Patients should be monitored for these signs and symptoms and advised not to drive or operate machinery until they have gained sufficient experience on levetiracetam to gauge whether it could adversely affect their ability to drive or operate machinery.

5.8 Withdrawal Seizures

As with most antiepileptic drugs, levetiracetam extended-release tablets should generally be withdrawn gradually because of the risk of increased seizure frequency and status epilepticus. If withdrawal is needed because of a serious adverse reaction, rapid discontinuation can be considered.

5.9 Hematologic Abnormalities

Levetiracetam extended-release tablets can cause hematologic abnormalities. Hematologic abnormalities occurred in clinical trials and included decreases in white blood cell (WBC), neutrophil, and red blood cell (RBC) counts; decreases in hemoglobin and hematocrit; and increases in eosinophil counts. Cases of agranulocytosis, pancytopenia, and thrombocytopenia have also been reported in the postmarketing setting. A complete blood count is recommended in patients experiencing significant weakness, pyrexia, recurrent infections, or coagulation disorders.

In controlled trials of immediate-release levetiracetam tablets in patients experiencing partial-onset seizures, minor, but statistically significant, decreases compared to placebo in total mean RBC count ($0.03 \times 10^6/\text{mm}^3$), mean hemoglobin (0.09 g/dL), and mean hematocrit (0.38%), were seen in immediate-release levetiracetam-treated patients.

A total of 3.2% of levetiracetam-treated and 1.8% of placebo-treated patients had at least one possibly significant ($\leq 2.8 \times 10^9/\text{L}$) decreased WBC, and 2.4% of levetiracetam-treated and 1.4% of placebo-treated patients had at least one possibly significant ($\leq 1 \times 10^9/\text{L}$) decreased neutrophil count. Of the levetiracetam-treated patients with a low neutrophil count, all but one rose towards or to baseline with continued treatment. No patient was discontinued secondary to low neutrophil counts.

In pediatric patients (4 to <16 years of age), statistically significant decreases in WBC and neutrophil counts were seen in patients treated with immediate-release levetiracetam tablets, as compared to placebo. The mean decreases from baseline in the immediate-release levetiracetam tablets group were $-0.4 \times 10^9/\text{L}$ and $-0.3 \times 10^9/\text{L}$, respectively, whereas there were small increases in the placebo group. A significant increase in mean relative lymphocyte counts was observed in 1.7% of patients treated with immediate-release levetiracetam tablets compared to a decrease of 4% in patients on placebo.

In the controlled pediatric trial, a possibly clinically significant abnormal low WBC value was observed in 3% of patients treated with immediate-release levetiracetam tablets, compared to no patients on placebo. However, there was no apparent difference between treatment groups with respect to neutrophil count. No patient was

discontinued secondary to low WBC or neutrophil counts.

In the controlled pediatric cognitive and neuropsychological safety study, two subjects (6.1%) in the placebo group and 5 subjects (8.6%) in the immediate-release levetiracetam-treated group had high eosinophil count values that were possibly clinically significant ($\geq 10\%$ or $\geq 0.7 \times 10^9/L$).

5.10 Seizure Control During Pregnancy

Physiological changes may gradually decrease plasma levels of levetiracetam throughout pregnancy. This decrease is more pronounced during the third trimester. It is recommended that patients be monitored carefully during pregnancy. Close monitoring should continue through the postpartum period especially if the dose was changed during pregnancy.

6 ADVERSE REACTIONS

The following adverse reactions are discussed in more details in other sections of labeling:

- Behavioral abnormalities and Psychotic Symptoms [*see Warnings and Precautions (5.1)*]
- Suicidal Behavior and Ideation [*see Warnings and Precautions (5.2)*]
- Somnolence and Fatigue [*see Warnings and Precautions (5.3)*]
- Anaphylaxis and Angioedema [*see Warnings and Precautions (5.4)*]
- Serious Dermatological Reactions [*see Warnings and Precautions (5.5)*]
- Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multiorgan Hypersensitivity [*see Warnings and Precautions (5.6)*]
- Coordination Difficulties [*see Warnings and Precautions (5.7)*]
- Hematologic Abnormalities [*see Warnings and Precautions (5.9)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Levetiracetam Extended-Release Tablets

In the controlled clinical study in patients with partial-onset seizures [*see Clinical Studies (14.1)*], the most common adverse reactions in patients receiving levetiracetam extended-release tablets in combination with other AEDs, for events with rates greater than placebo, were irritability and somnolence.

Table 3 lists adverse reactions that occurred in at least 5% of epilepsy patients receiving levetiracetam extended-release tablets in the placebo-controlled study and were numerically more common than in patients treated with placebo. In this study, either levetiracetam extended-release tablets or placebo was added to concurrent AED therapy.

Table 3: Adverse Reactions in the Placebo-Controlled, Adjunctive Study in Patients Experiencing Partial-Onset Seizures

	Levetiracetam Extended-Release Tablets (N=77) %	Placebo (N=79) %
Influenza	8	4
Somnolence	8	3
Irritability	7	0
Nasopharyngitis	7	5
Dizziness	5	3
Nausea	5	3

Discontinuation or Dose Reduction in the Levetiracetam Extended-Release Tablets Controlled Clinical Study

In the controlled clinical study, 5% of patients receiving levetiracetam extended-release tablets and 3% receiving placebo discontinued as a result of an adverse reaction. The adverse reactions that resulted in discontinuation and that occurred more frequently in levetiracetam extended-release-treated patients than in placebo-treated patients were asthenia, epilepsy, mouth ulceration, rash, and respiratory failure. Each of these adverse reactions led to discontinuation in a levetiracetam extended-release-treated patient and no placebo-treated patients.

Immediate-Release Levetiracetam Tablets

Table 4 lists the adverse reactions in the controlled studies of immediate-release levetiracetam tablets in adult patients experiencing partial-onset seizures [see *Clinical Studies (14.2)*]. Although the pattern of adverse reactions in the levetiracetam extended-release tablets study seems somewhat different from that seen in partial-onset seizure controlled studies for immediate-release levetiracetam tablets, this is possibly due to the much smaller number of patients in this study compared to the immediate-release tablet studies. The adverse reactions for levetiracetam extended-release tablets are expected to be similar to those seen with immediate-release levetiracetam tablets.

Adults

In controlled clinical studies of immediate-release levetiracetam tablets as adjunctive therapy to other AEDs in adults with partial-onset seizures, the most common adverse reactions, for events with rates greater than placebo, were somnolence, asthenia, infection, and dizziness.

Table 4 lists adverse reactions that occurred in at least 1% of adult epilepsy patients receiving immediate-release levetiracetam tablets in placebo-controlled studies and were numerically more common than in patients treated with placebo. In these studies, either immediate-release levetiracetam tablets or placebo was added to concurrent AED therapy.

Table 4: Adverse Reactions in Pooled Placebo-Controlled, Adjunctive Studies in Adults Experiencing Partial-Onset Seizures

	Levetiracetam (N=769) %	Placebo (N=439) %

Asthenia	15	9
Somnolence	15	8
Headache	14	13
Infection	13	8
Dizziness	9	4
Pain	7	6
Pharyngitis	6	4
Depression	4	2
Nervousness	4	2
Rhinitis	4	3
Anorexia	3	2
Ataxia	3	1
Vertigo	3	1
Amnesia	2	1
Anxiety	2	1
Cough Increased	2	1
Diplopia	2	1
Emotional Lability	2	0
Hostility	2	1
Paresthesia	2	1
Sinusitis	2	1

Pediatric Patients 4 Years to <16 Years

In a pooled analysis of two controlled pediatric clinical studies in children 4 to 16 years of age with partial-onset seizures [see *Clinical Studies (14.3)*], the adverse reactions most frequently reported with the use of immediate-release levetiracetam tablets in combination with other AEDs, and with greater frequency than in patients on placebo, were fatigue, aggression, nasal congestion, decreased appetite, and irritability.

Table 5 lists adverse reactions that occurred in at least 2% of pediatric patients treated with immediate-release levetiracetam tablets and were more common than in pediatric patients on placebo. In these studies, either immediate-release levetiracetam tablets or placebo was added to concurrent AED therapy. Adverse reactions were usually mild to moderate in intensity.

Table 5: Adverse Reactions in Pooled Placebo-Controlled, Adjunctive Studies in Pediatric Patients Ages 4 to 16 Years Experiencing Partial-Onset Seizures

	Levetiracetam (N=165) %	Placebo (N=131) %
Headache	19	15
Nasopharyngitis	15	12
Vomiting	15	12
Somnolence	13	9
Fatigue	11	5

Aggression	10	5
Upper Abdominal Pain	9	8
Cough	9	5
Nasal Congestion	9	2
Decreased Appetite	8	2
Abnormal Behavior	7	4
Dizziness	7	5
Irritability	7	1
Pharyngolaryngeal Pain	7	4
Diarrhea	6	2
Lethargy	6	5
Insomnia	5	3
Agitation	4	1
Anorexia	4	3
Head Injury	4	0
Constipation	3	1
Contusion	3	1
Depression	3	1
Fall	3	2
Influenza	3	1
Mood Altered	3	1
Affect Lability	2	1
Anxiety	2	1
Arthralgia	2	0
Confusional State	2	0
Conjunctivitis	2	0
Ear Pain	2	1
Gastroenteritis	2	0
Joint Sprain	2	1
Mood Swings	2	1
Neck Pain	2	1
Rhinitis	2	0
Sedation	2	1

In controlled pediatric clinical studies in patients 4 to 16 years of age, 7% of patients treated with immediate-release levetiracetam tablets and 9% of patients on placebo discontinued as a result of an adverse event.

In addition, the following adverse reactions were seen in other controlled studies of immediate-release levetiracetam tablets: balance disorder, disturbance in attention, eczema, hyperkinesia, memory impairment, myalgia, personality disorders, pruritus, and blurred vision.

Comparison of Gender, Age and Race

There are insufficient data for levetiracetam extended-release tablets to support a

statement regarding the distribution of adverse reactions by gender, age, and race.

6.2 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of immediate-release levetiracetam tablets. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

The listing is alphabetized: abnormal liver function test, acute kidney injury, anaphylaxis, angioedema, agranulocytosis, choreoathetosis, drug reaction with eosinophilia and systemic symptoms (DRESS), dyskinesia, erythema multiforme, hepatic failure, hepatitis, hyponatremia, muscular weakness, obsessive-compulsive disorders (OCD), pancreatitis, pancytopenia (with bone marrow suppression identified in some of these cases), panic attack, thrombocytopenia, weight loss, and worsening of seizures including in patients with SCN8A mutations. Alopecia has been reported with immediate-release levetiracetam use; recovery was observed in majority of cases where immediate-release levetiracetam was discontinued.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to antiepileptic drugs (AEDs), including levetiracetam extended-release tablets, during pregnancy. Encourage women who are taking levetiracetam extended-release tablets during pregnancy to enroll in the North American Antiepileptic Drug (NAAED) pregnancy registry by calling 1-888-233-2334 or visiting <http://www.aedpregnancyregistry.org/>.

Risk Summary

Prolonged experience with levetiracetam tablets in pregnant women has not identified a drug-associated risk of major birth defects or miscarriage, based on published literature, which includes data from pregnancy registries and reflects experience over two decades [see *Human Data*]. In animal studies, levetiracetam produced developmental toxicity (increased embryofetal and offspring mortality, increased incidences of fetal structural abnormalities, decreased embryofetal and offspring growth, neurobehavioral alterations in offspring) at doses similar to human therapeutic doses [see *Animal Data*].

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively. The background risk of major birth defects and miscarriage for the indicated population is unknown.

Clinical Considerations

Levetiracetam extended-release tablets levels may decrease during pregnancy [see *Warnings and Precautions (5.10)*].

Physiological changes during pregnancy may affect levetiracetam concentration. Decrease in levetiracetam plasma concentrations has been observed during pregnancy.

This decrease is more pronounced during the third trimester. Dose adjustments may be necessary to maintain clinical response.

Data

Human Data

While available studies cannot definitively establish the absence of risk, data from the published literature and pregnancy registries have not established an association with levetiracetam use during pregnancy and major birth defects or miscarriage.

Animal Data

When levetiracetam (0, 400, 1,200, or 3,600 mg/kg/day) was administered orally to pregnant rats during the period of organogenesis, reduced fetal weights and increased incidence of fetal skeletal variations were observed at the highest dose tested. There was no evidence of maternal toxicity. The no-effect dose for adverse effects on embryofetal developmental in rats (1,200 mg/kg/day) is approximately 4 times the maximum recommended human dose (MRHD) of 3,000 mg on a body surface area (mg/m²) basis.

Oral administration of levetiracetam (0, 200, 600, or 1,800 mg/kg/day) to pregnant rabbits during the period of organogenesis resulted in increased embryofetal mortality and incidence of fetal skeletal variations at the mid and high dose and decreased fetal weights and increased incidence of fetal malformations at the high dose, which was associated with maternal toxicity. The no-effect dose for adverse effects on embryofetal development in rabbits (200 mg/kg/day) is approximately equivalent to the MRHD on a mg/m² basis.

Oral administration of levetiracetam (0, 70, 350, or 1,800 mg/kg/day) to female rats throughout pregnancy and lactation led to an increased incidence of fetal skeletal variations, reduced fetal body weight, and decreased growth in offspring at the mid and high doses and increased pup mortality and neurobehavioral alterations in offspring at the highest dose tested. There was no evidence of maternal toxicity. The no-effect dose for adverse effects on pre- and postnatal development in rats (70 mg/kg/day) is less than the MRHD on a mg/m² basis.

Oral administration of levetiracetam to rats during the latter part of gestation and throughout lactation produced no adverse developmental or maternal effects at doses of up to 1,800 mg/kg/day (6 times the MRHD on a mg/m² basis).

8.2 Lactation

Risk Summary

Levetiracetam is excreted in human milk. There are no data on the effects of levetiracetam extended-release tablets on the breastfed infant, or the effects on milk production.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Levetiracetam extended-release tablets and any potential adverse effects on the breastfed infant from Levetiracetam extended-release tablets or from the underlying maternal condition.

8.4 Pediatric Use

Safety and effectiveness in patients 12 years of age and older have been established based on pharmacokinetic data in adults and adolescents using levetiracetam extended-release tablets and efficacy and safety data in controlled pediatric studies using immediate-release levetiracetam tablets [see *Adverse Reactions (6.1)*, *Clinical Pharmacology (12.3)*, and *Clinical Studies (14.1)*].

Safety and effectiveness in pediatric patients below the age of 12 have not been established.

A 3-month, randomized, double-blind, placebo-controlled study was performed to assess the neurocognitive and behavioral effects of immediate-release levetiracetam tablets as adjunctive therapy in 98 pediatric patients with inadequately controlled partial seizures, ages 4 to 16 years (levetiracetam tablets N=64; placebo N=34). The target dose of immediate-release levetiracetam tablets was 60 mg/kg/day. Neurocognitive effects were measured by the Leiter-R Attention and Memory (AM) Battery, which assesses various aspects of a child's memory and attention. Although no substantive differences were observed between the placebo- and levetiracetam-treated groups in the median change from baseline in this battery, the study was not adequate to assess formal statistical non-inferiority between the drug and placebo. The Achenbach Child Behavior Checklist (CBCL/6-18), a standardized validated tool used to assess a child's competencies and behavioral/emotional problems, was also assessed in this study. An analysis of the CBCL/6-18 indicated a worsening in aggressive behavior, one of the eight syndrome scores, in patients treated with levetiracetam tablets [see *Warnings and Precautions (5.1)*].

Juvenile Animal Toxicity Data

Studies of levetiracetam in juvenile rats (dosed on postnatal days 4 through 52) and dogs (dosed from postnatal weeks 3 through 7) at doses of up to 1,800 mg/kg/day (approximately 7 and 24 times, respectively, the maximum recommended pediatric dose of 60 mg/kg/day on a mg/m² basis) did not demonstrate adverse effects on postnatal development.

8.5 Geriatric Use

There were insufficient numbers of elderly subjects in controlled trials of epilepsy to adequately assess the effectiveness of levetiracetam extended-release in these patients. It is expected that the safety of levetiracetam extended-release in elderly patients 65 and over would be comparable to the safety observed in clinical studies of immediate-release levetiracetam tablets.

There were 347 subjects in clinical studies of immediate-release levetiracetam that were 65 and over. No overall differences in safety were observed between these subjects and younger subjects. There were insufficient numbers of elderly subjects in controlled trials of epilepsy to adequately assess the effectiveness of immediate-release levetiracetam in these patients.

Levetiracetam is known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function [see *Clinical Pharmacology (12.3)*].

8.6 Renal Impairment

The effect of levetiracetam extended-release on renally impaired patients was not assessed in the controlled study. However, it is expected that the effect on levetiracetam extended-release-treated patients would be similar to the effect seen in controlled studies of immediate-release levetiracetam tablets. Clearance of levetiracetam is decreased in patients with renal impairment and is correlated with creatinine clearance [see *Clinical Pharmacology (12.3)*]. Dose adjustment is recommended for patients with impaired renal function [see *Dosage and Administration (2.2)*].

10 OVERDOSAGE

10.1 Signs, Symptoms and Laboratory Findings of Acute Overdosage in Humans

The signs and symptoms for levetiracetam extended-release tablets overdose are expected to be similar to those seen with immediate-release levetiracetam tablets.

The highest known dose of oral immediate-release levetiracetam received in the clinical development program was 6,000 mg/day. Other than drowsiness, there were no adverse reactions in the few known cases of overdose in clinical trials. Cases of somnolence, agitation, aggression, depressed level of consciousness, respiratory depression and coma were observed with immediate-release levetiracetam overdoses in postmarketing use.

10.2 Management of Overdose

There is no specific antidote for overdose with levetiracetam extended-release tablets. If indicated, elimination of unabsorbed drug should be attempted by emesis or gastric lavage; usual precautions should be observed to maintain airway. General supportive care of the patient is indicated including monitoring of vital signs and observation of the patient's clinical status. A Certified Poison Control Center should be contacted for up to date information on the management of overdose with levetiracetam extended-release tablets.


10.3 Hemodialysis

Standard hemodialysis procedures result in significant clearance of levetiracetam (approximately 50% in 4 hours) and should be considered in cases of overdose. Although hemodialysis has not been performed in the few known cases of overdose, it may be indicated by the patient's clinical state or in patients with significant renal impairment.

11 DESCRIPTION

Levetiracetam extended-release tablets, USP are an antiepileptic drug available as 500 mg, 750 mg and 1,000 mg (white) extended-release tablets, USP for oral administration.

The chemical name of levetiracetam, a single enantiomer, is (α S)- α -ethyl-2-oxo-1-pyrrolidineacetamide, its molecular formula is $C_8H_{14}N_2O_2$ and its molecular weight is 170.21. Levetiracetam is chemically unrelated to existing antiepileptic drugs (AEDs). It has the following structural formula:



Levetiracetam USP is a white to off-white powder. It is very soluble in water (104.0 g/100 mL). It is freely soluble in chloroform (65.3 g/100 mL) and in methanol (53.6 g/100 mL), soluble in ethanol (16.5 g/100 mL), sparingly soluble in acetonitrile (5.7 g/100 mL) and practically insoluble in n-hexane. (Solubility limits are expressed as g/100 mL solvent.)

Levetiracetam extended-release tablets, USP contain the labeled amount of levetiracetam. Inactive ingredients: colloidal silicon dioxide, hydroxypropyl cellulose, hydroxypropyl methylcellulose, polyethylene glycol, and titanium dioxide.

The medication is combined with a drug release controlling polymer that provides a drug release at a controlled rate. The biologically inert components of the tablet may occasionally remain intact during GI transit and will be eliminated in the feces as a soft, hydrated mass.

Meets USP Dissolution Test 5.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The precise mechanism(s) by which levetiracetam exerts its antiepileptic effect is unknown.

A saturable and stereoselective neuronal binding site in rat brain tissue has been described for levetiracetam. Experimental data indicate that this binding site is the synaptic vesicle protein SV2A, thought to be involved in the regulation of vesicle exocytosis. Although the molecular significance of levetiracetam binding to synaptic vesicle protein SV2A is not understood, levetiracetam and related analogs showed a rank order of affinity for SV2A which correlated with the potency of their antiseizure activity in audiogenic seizure-prone mice. These findings suggest that the interaction of levetiracetam with the SV2A protein may contribute to the antiepileptic mechanism of action of the drug.

12.2 Pharmacodynamics

Effects on QTc Interval

The effects of levetiracetam extended-release tablets on QTc prolongation is expected to be the same as that of immediate-release levetiracetam tablets. The effect of immediate-release levetiracetam tablets on QTc prolongation was evaluated in a randomized, double-blind, positive-controlled (moxifloxacin 400 mg) and placebo-controlled crossover study of levetiracetam tablets (1,000 mg or 5,000 mg) in 52 healthy subjects. The upper bound of the 90% confidence interval for the largest placebo-adjusted, baseline-corrected QTc was below 10 milliseconds. Therefore, there was no evidence of significant QTc prolongation in this study.

12.3 Pharmacokinetics

Overview

Bioavailability of levetiracetam extended-release tablets are similar to that of the immediate-release levetiracetam tablets. The pharmacokinetics (AUC and C_{max}) were shown to be dose proportional after single dose administration of 1,000 mg, 2,000 mg, and 3,000 mg extended-release levetiracetam. Plasma half-life of extended-release levetiracetam is approximately 7 hours.

Levetiracetam is almost completely absorbed after oral administration. The pharmacokinetics of levetiracetam are linear and time-invariant, with low intra- and inter-subject variability. Levetiracetam is not significantly protein-bound (<10% bound) and its volume of distribution is close to the volume of intracellular and extracellular water. Sixty-six percent (66%) of the dose is renally excreted unchanged. The major metabolic pathway of levetiracetam (24% of dose) is an enzymatic hydrolysis of the acetamide group. It is not liver cytochrome P450 dependent. The metabolites have no known pharmacological activity and are renally excreted. Plasma half-life of levetiracetam across studies is approximately 6 to 8 hours. The half-life is increased in the elderly (primarily due to impaired renal clearance) and in subjects with renal impairment.

The pharmacokinetics of levetiracetam are similar when used as monotherapy or as adjunctive therapy for the treatment of partial-onset seizures.

Absorption and Distribution

Extended-release levetiracetam peak plasma concentrations occur in about 4 hours. The time to peak plasma concentrations is about 3 hours longer with extended-release levetiracetam than with immediate-release tablets.

Single administration of two 500 mg extended-release levetiracetam tablets once daily produced comparable maximal plasma concentrations and area under the plasma concentration versus time as did the administration of one 500 mg immediate-release tablet twice daily in fasting conditions. After multiple dose extended-release levetiracetam tablets intake, extent of exposure (AUC_{0-24}) was similar to extent of exposure after multiple dose immediate-release tablets intake. C_{max} and C_{min} were lower by 17% and 26% after multiple dose extended-release levetiracetam tablets intake in comparison to multiple dose immediate-release tablets intake. Intake of a high fat, high calorie breakfast before the administration of extended-release levetiracetam tablets resulted in a higher peak concentration, and longer median time to peak. The median time to peak (T_{max}) was 2 hours longer in the fed state.

Two 750 mg extended-release levetiracetam tablets were bioequivalent to a single administration of three 500 mg extended-release levetiracetam tablets.

Metabolism

Levetiracetam is not extensively metabolized in humans. The major metabolic pathway is the enzymatic hydrolysis of the acetamide group, which produces the carboxylic acid metabolite, ucb L057 (24% of dose) and is not dependent on any liver cytochrome P450 isoenzymes. The major metabolite is inactive in animal seizure models. Two minor metabolites were identified as the product of hydroxylation of the 2-oxo-pyrrolidine ring (2% of dose) and opening of the 2-oxo-pyrrolidine ring in position 5 (1% of dose). There is no enantiomeric interconversion of levetiracetam or its major metabolite.

Elimination

Levetiracetam plasma half-life in adults is 7 ± 1 hour and is unaffected by either dose or

repeated administration. Levetiracetam is eliminated from the systemic circulation by renal excretion as unchanged drug which represents 66% of administered dose. The total body clearance is 0.96 mL/min/kg and the renal clearance is 0.6 mL/min/kg. The mechanism of excretion is glomerular filtration with subsequent partial tubular reabsorption. The metabolite ucb L057 is excreted by glomerular filtration and active tubular secretion with a renal clearance of 4 mL/min/kg. Levetiracetam elimination is correlated to creatinine clearance. Levetiracetam clearance is reduced in patients with impaired renal function [see *Dosage and Administration (2.2) and Use in Specific Populations (8.6)*].

Specific Populations

Elderly

There are insufficient pharmacokinetic data to specifically address the use of extended-release levetiracetam in the elderly population.

Pharmacokinetics of immediate-release levetiracetam were evaluated in 16 elderly subjects (age 61 to 88 years) with creatinine clearance ranging from 30 to 74 mL/min. Following oral administration of twice-daily dosing for 10 days, total body clearance decreased by 38% and the half-life was 2.5 hours longer in the elderly compared to healthy adults. This is most likely due to the decrease in renal function in these subjects.

Pediatric Patients

An open label, multicenter, parallel-group, two-arm study was conducted to evaluate the pharmacokinetics of levetiracetam extended-release tablets in pediatric patients (13 to 16 years old) and in adults (18 to 55 years old) with epilepsy. Levetiracetam extended-release oral tablets (1,000 mg to 3,000 mg) were administered once daily with a minimum of 4 days and a maximum of 7 days of treatment to 12 pediatric patients and 13 adults in the study. Dose-normalized steady-state exposure parameters, C_{max} and AUC, were comparable between pediatric and adult patients.

Pregnancy

Levetiracetam extended-release tablets levels may decrease during pregnancy [see *Warnings and Precautions (5.10) and Use in Specific Populations (8.1)*].

Gender

Extended-release levetiracetam C_{max} was 21 to 30% higher and AUC was 8 to 18% higher in women (N=12) compared to men (N=12). However, clearances adjusted for body weight were comparable.

Race

Formal pharmacokinetic studies of the effects of race have not been conducted with extended-release or immediate-release levetiracetam. Cross study comparisons involving Caucasians (N=12) and Asians (N=12), however, show that pharmacokinetics of immediate-release levetiracetam were comparable between the two races. Because levetiracetam is primarily renally excreted and there are no important racial differences in creatinine clearance, pharmacokinetic differences due to race are not expected.

Renal Impairment

The effect of levetiracetam extended-release tablets on renally impaired patients was not assessed in the controlled study. However, it is expected that the effect on levetiracetam extended-release-treated patients would be similar to that seen in controlled studies of immediate-release levetiracetam tablets. In patients with end stage renal disease on dialysis, it is recommended that immediate-release levetiracetam be

used instead of levetiracetam extended-release tablets.

The disposition of immediate-release levetiracetam was studied in adult subjects with varying degrees of renal function. Total body clearance of levetiracetam is reduced in patients with impaired renal function by 40% in the mild group (CLcr = 50 to 80 mL/min), 50% in the moderate group (CLcr = 30 to 50 mL/min) and 60% in the severe renal impairment group (CLcr <30 mL/min). Clearance of levetiracetam is correlated with creatinine clearance.

In anuric (end stage renal disease) patients, the total body clearance decreased 70% compared to normal subjects (CLcr >80 mL/min). Approximately 50% of the pool of levetiracetam in the body is removed during a standard 4-hour hemodialysis procedure [see *Dosage and Administration (2.2)*].

Hepatic Impairment

In subjects with mild (Child-Pugh A) to moderate (Child-Pugh B) hepatic impairment, the pharmacokinetics of levetiracetam were unchanged. In patients with severe hepatic impairment (Child-Pugh C), total body clearance was 50% that of normal subjects, but decreased renal clearance accounted for most of the decrease. No dose adjustment is needed for patients with hepatic impairment.

Drug Interactions

In vitro data on metabolic interactions indicate that levetiracetam is unlikely to produce, or be subject to, pharmacokinetic interactions. Levetiracetam and its major metabolite, at concentrations well above C_{max} levels achieved within the therapeutic dose range, are neither inhibitors of, nor high affinity substrates for, human liver cytochrome P450 isoforms, epoxide hydrolase or UDP-glucuronidation enzymes. In addition, levetiracetam does not affect the *in vitro* glucuronidation of valproic acid.

Potential pharmacokinetic interactions of or with levetiracetam were assessed in clinical pharmacokinetic studies (phenytoin, valproate, warfarin, digoxin, oral contraceptive, probenecid) and through pharmacokinetic screening with immediate-release levetiracetam tablets in the placebo-controlled clinical studies in epilepsy patients. The potential for drug interactions for levetiracetam extended-release tablets are expected to be essentially the same as that with immediate-release levetiracetam tablets.

Phenytoin

Immediate-release levetiracetam tablets (3,000 mg daily) had no effect on the pharmacokinetic disposition of phenytoin in patients with refractory epilepsy. Pharmacokinetics of levetiracetam were also not affected by phenytoin.

Valproate

Immediate-release levetiracetam tablets (1,500 mg twice daily) did not alter the pharmacokinetics of valproate in healthy volunteers. Valproate 500 mg twice daily did not modify the rate or extent of levetiracetam absorption or its plasma clearance or urinary excretion. There also was no effect on exposure to and the excretion of the primary metabolite, ucb L057.

Other Antiepileptic Drugs

Potential drug interactions between immediate-release levetiracetam tablets and other AEDs (carbamazepine, gabapentin, lamotrigine, phenobarbital, phenytoin, primidone and valproate) were also assessed by evaluating the serum concentrations of levetiracetam and these AEDs during placebo-controlled clinical studies. These data indicate that

levetiracetam does not influence the plasma concentration of other AEDs and that these AEDs do not influence the pharmacokinetics of levetiracetam.

Oral Contraceptives

Immediate-release levetiracetam tablets (500 mg twice daily) did not influence the pharmacokinetics of an oral contraceptive containing 0.03 mg ethinyl estradiol and 0.15 mg levonorgestrel, or of the luteinizing hormone and progesterone levels, indicating that impairment of contraceptive efficacy is unlikely. Coadministration of this oral contraceptive did not influence the pharmacokinetics of levetiracetam.

Digoxin

Immediate-release levetiracetam tablets (1,000 mg twice daily) did not influence the pharmacokinetics and pharmacodynamics (ECG) of digoxin given as a 0.25 mg dose every day. Coadministration of digoxin did not influence the pharmacokinetics of levetiracetam.

Warfarin

Immediate-release levetiracetam tablets (1,000 mg twice daily) did not influence the pharmacokinetics of R and S warfarin. Prothrombin time was not affected by levetiracetam. Coadministration of warfarin did not affect the pharmacokinetics of levetiracetam.

Probenecid

Probenecid, a renal tubular secretion blocking agent, administered at a dose of 500 mg four times a day, did not change the pharmacokinetics of levetiracetam 1,000 mg twice daily. $C_{ss_{max}}$ of the metabolite, ucb L057, was approximately doubled in the presence of probenecid while the fraction of drug excreted unchanged in the urine remained the same. Renal clearance of ucb L057 in the presence of probenecid decreased 60%, probably related to competitive inhibition of tubular secretion of ucb L057. The effect of immediate-release levetiracetam tablets on probenecid was not studied.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Rats were dosed with levetiracetam in the diet for 104 weeks at doses of 50, 300, and 1,800 mg/kg/day. Plasma exposure (AUC) at the highest dose was approximately 6 times that in humans at the maximum recommended daily human dose (MRHD) of 3,000 mg. There was no evidence of carcinogenicity. In mice, oral administration of levetiracetam for 80 weeks (doses up to 960 mg/kg/day) or 2 years (doses up to 4,000 mg/kg/day, lowered to 3,000 mg/kg/day after 45 weeks due to intolerability) was not associated with an increase in tumors. The highest dose tested in mice for 2 years (3,000 mg/kg/day) is approximately 5 times the MRHD on a body surface area (mg/m²) basis.

Mutagenesis

Levetiracetam was negative in *in vitro* (Ames, chromosomal aberration in mammalian cells) and *in vivo* (mouse micronucleus) assays. The major human metabolite of levetiracetam (ucb L057) was negative in *in vitro* (Ames, mouse lymphoma) assays.

Impairment of Fertility

No adverse effects on male or female fertility or reproductive performance were observed in rats at oral doses up to 1,800 mg/kg/day, which were associated with plasma exposures (AUC) up to approximately 6 times that in humans at the MRHD.

14 CLINICAL STUDIES

The effectiveness of levetiracetam extended-release for the treatment of partial-onset seizures in adults was established in one multicenter, randomized, double-blind, placebo-controlled clinical study in patients who had refractory partial-onset seizures with or without secondary generalization. This was supported by the demonstration of efficacy of immediate-release levetiracetam tablets (see below) in partial seizures in three multicenter, randomized, double-blind, placebo-controlled clinical studies in adults, as well as a demonstration of comparable bioavailability between the extended-release and immediate-release formulations [see *Clinical Pharmacology (12.3)*] in adults. The effectiveness for levetiracetam extended-release tablets for the treatment of partial-onset seizures in pediatric patients, 12 years of age and older, was based upon a single pharmacokinetic study showing comparable pharmacokinetics of extended-release levetiracetam tablets in adults and adolescents [see *Clinical Pharmacology (12.3)*]. All studies are described below.

14.1 Levetiracetam Extended-Release Tablets in Adults

The effectiveness of levetiracetam extended-release tablets for the treatment of partial-onset seizures in adults was established in one multicenter, randomized, double-blind, placebo-controlled clinical study across 7 countries in patients who had refractory partial-onset seizures with or without secondary generalization (Study 1).

Study 1

Patients enrolled in Study 1 had at least eight partial seizures with or without secondary generalization during the 8-week baseline period and at least two partial seizures in each 4-week interval of the baseline period. Patients were taking a stable dose regimen of at least one AED and could take a maximum of three AEDs. After a prospective baseline period of 8 weeks, 158 patients were randomized to placebo (N=79) or 1,000 mg (two 500 mg tablets) of levetiracetam extended-release tablets (N=79), given once daily over a 12-week treatment period.

The primary efficacy endpoint in Study 1 was the percent reduction over placebo in mean weekly frequency of partial-onset seizures. The median percent reduction in weekly partial-onset seizure frequency from baseline over the treatment period was 46.1% in the levetiracetam extended-release tablets 1,000 mg treatment group (N=74) and 33.4% in the placebo group (N=78). The estimated percent reduction over placebo in weekly partial-onset seizure frequency over the treatment period was 14.4% (statistically significant).

The relationship between the effectiveness of the same daily dose of levetiracetam extended-release tablets and immediate-release levetiracetam has not been studied and is unknown.

14.2 Immediate-Release Levetiracetam Tablets in Adults

The effectiveness of immediate-release levetiracetam tablets for the treatment of partial-onset seizures in adults was established in three multicenter, randomized, double-blind,

placebo-controlled clinical studies in patients who had refractory partial-onset seizures with or without secondary generalization (Studies 2, 3, and 4). The tablet formulation was used in all three studies. In these studies, 904 patients were randomized to placebo, levetiracetam tablets 1,000 mg, levetiracetam tablets 2,000 mg, or levetiracetam tablets 3,000 mg/day. Patients enrolled in Study 2 or Study 3 had refractory partial-onset seizures for at least two years and had taken two or more AEDs. Patients enrolled in Study 4 had refractory partial-onset seizures for at least 1 year and had taken one AED. At the time of the study, patients were taking a stable dose regimen of at least one AED and could take a maximum of two AEDs. During the baseline period, patients had to have experienced at least two partial-onset seizures during each 4-week period.

Study 2

Study 2 was a double-blind, placebo-controlled, parallel-group study conducted at 41 sites in the United States, comparing immediate-release levetiracetam tablets 1,000 mg/day (N=97), immediate-release levetiracetam tablets 3,000 mg/day (N=101), and placebo (N=95), given in equally divided doses twice daily. After a prospective baseline period of 12 weeks, patients in Study 2 were randomized to one of the three treatment groups described above. The 18-week treatment period consisted of a 6-week titration period, followed by a 12-week fixed dose evaluation period, during which concomitant AED regimens were held constant. The primary measure of effectiveness in Study 2 was a between-group comparison of the percent reduction in weekly partial seizure frequency relative to placebo over the entire randomized treatment period (titration + evaluation period). Secondary outcome variables included the responder rate (incidence of patients with $\geq 50\%$ reduction from baseline in partial-onset seizure frequency). The results of Study 2 are displayed in Table 6.


Table 6: Reduction In Mean Over Placebo In Weekly Frequency Of Partial-Onset Seizures In Study 2

	Placebo (N=95)	Immediate-release Levetiracetam Tablets 1,000 mg/day (N=97)	Immediate-release Levetiracetam Tablets 3,000 mg/day (N=101)
Percent reduction in partial seizure frequency over placebo	-	26.1%*	30.1%*

* statistically significant versus placebo

The percentage of patients (y-axis) who achieved $\geq 50\%$ reduction from baseline in weekly partial-onset seizure frequency over the entire randomized treatment period (titration + evaluation period) within the three treatment groups (x-axis) in Study 2 is presented in Figure 1.

Figure 1: Responder Rate ($\geq 50\%$ Reduction From Baseline) In Study 2



* statistically significant versus placebo

Study 3

Study 3 was a double-blind, placebo-controlled, crossover study conducted at 62 centers in Europe, comparing immediate-release levetiracetam tablets 1,000 mg/day (N=106), immediate-release levetiracetam tablets 2,000 mg/day (N=105), and placebo (N=111), given in equally divided doses twice daily.

The first period of the study (Period A) was designed to be analyzed as a parallel-group study. After a prospective baseline period of up to 12 weeks, patients in Study 3 were randomized to one of the three treatment groups described above. The 16-week treatment period consisted of the 4-week titration period followed by a 12-week fixed dose evaluation period, during which concomitant AED regimens were held constant. The primary measure of effectiveness in Study 3 was a between group comparison of the percent reduction in weekly partial seizure frequency relative to placebo over the entire randomized treatment period (titration + evaluation period). Secondary outcome variables included the responder rate (incidence of patients with $\geq 50\%$ reduction from baseline in partial-onset seizure frequency). The results of the analysis of Period A are displayed in Table 7.


Table 7: Reduction In Mean Over Placebo In Weekly Frequency Of Partial-Onset Seizures In Study 3: Period A

	Placebo (N=111)	Immediate-release Levetiracetam Tablets 1,000 mg/day (N=106)	Immediate-release Levetiracetam Tablets 2,000 mg/day (N=105)
Percent reduction in partial seizure frequency over placebo	-	17.1%*	21.4%*

* statistically significant versus placebo

The percentage of patients (y-axis) who achieved $\geq 50\%$ reduction from baseline in weekly partial-onset seizure frequency over the entire randomized treatment period (titration + evaluation period) within the three treatment groups (x-axis) in Study 3 is presented in Figure 2.

Figure 2: Responder Rate ($\geq 50\%$ Reduction From Baseline) In Study 3: Period A



* statistically significant versus placebo

The comparison of immediate-release levetiracetam tablets 2,000 mg/day to immediate-release levetiracetam tablets 1,000 mg/day for responder rate in Study 3 was statistically significant ($P=0.02$). Analysis of the trial as a cross-over study yielded similar results.

Study 4

Study 4 was a double-blind, placebo-controlled, parallel-group study conducted at 47 centers in Europe comparing immediate-release levetiracetam tablets 3,000 mg/day (N=180) and placebo (N=104) in patients with refractory partial-onset seizures, with or without secondary generalization, receiving only one concomitant AED. Study drug was given in two divided doses. After a prospective baseline period of 12 weeks, patients in Study 4 were randomized to one of two treatment groups described above. The 16-week treatment period consisted of a 4-week titration period, followed by a 12-week fixed dose evaluation period, during which concomitant AED doses were held constant. The primary measure of effectiveness in Study 4 was a between group comparison of the percent reduction in weekly seizure frequency relative to placebo over the entire randomized treatment period (titration + evaluation period). Secondary outcome variables included the responder rate (incidence of patients with $\geq 50\%$ reduction from baseline in partial-onset seizure frequency). Table 8 displays the results of Study 4.


Table 8: Reduction In Mean Over Placebo In Weekly Frequency Of Partial-Onset Seizures In Study 4

	Placebo (N=104)	Immediate-release Levetiracetam Tablets 3,000 mg/day (N=180)
Percent reduction in partial seizure frequency over placebo	-	23.0%*

* statistically significant versus placebo

The percentage of patients (y-axis) who achieved $\geq 50\%$ reduction from baseline in weekly partial-onset seizure frequency over the entire randomized treatment period (titration + evaluation period) within the two treatment groups (x-axis) in Study 4 is presented in Figure 3.

Figure 3: Responder Rate ($\geq 50\%$ Reduction From Baseline) In Study 4



* statistically significant versus placebo

14.3 Immediate-Release Levetiracetam Tablets in Pediatric Patients 4 Years to 16 Years

The use of levetiracetam extended-release tablets in pediatric patients 12 years of age and older is supported by Study 5, which was conducted using immediate-release levetiracetam tablets. Levetiracetam extended-release tablets are not indicated in children below 12 years of age.

Study 5

The effectiveness of immediate-release levetiracetam tablets for the treatment of partial-onset seizures in pediatric patients was established in a multicenter, randomized double-blind, placebo-controlled study, conducted at 60 sites in North America, in children 4 to 16 years of age with partial seizures uncontrolled by standard antiepileptic drugs (Study 5). Eligible patients on a stable dose of 1 to 2 AEDs, who still experienced at least 4 partial-onset seizures during the 4 weeks prior to screening, as well as at least 4 partial-onset seizures in each of the two 4-week baseline periods, were randomized to receive either immediate-release levetiracetam tablets or placebo. The enrolled population included 198 patients (levetiracetam tablets N=101; placebo N=97) with refractory partial-onset seizures, with or without secondarily generalization. Study 5 consisted of an 8-week baseline period and 4-week titration period followed by a 10-week evaluation period. Dosing was initiated at a dose of 20 mg/kg/day in two divided doses. During the treatment period, the immediate-release levetiracetam tablets doses were adjusted in 20 mg/kg/day increments, at 2-week intervals to the target dose of 60 mg/kg/day. The primary measure of effectiveness in Study 5 was a between group comparison of the percent reduction in weekly partial seizure frequency relative to placebo over the entire 14-week randomized treatment period (titration + evaluation period). Secondary outcome variables included the responder rate (incidence of patients with $\geq 50\%$ reduction from baseline in partial-onset seizure frequency per week). Table 9 displays the results of this study.


Table 9: Reduction In Mean Over Placebo In Weekly Frequency Of Partial-Onset Seizures in Study 5

	Placebo (N=97)	Immediate-release Levetiracetam Tablets (N=101)
Percent reduction in partial seizure frequency over placebo	-	26.8%*

*statistically significant versus placebo

The percentage of patients (y-axis) who achieved $\geq 50\%$ reduction in weekly partial-onset seizure frequency over the entire randomized treatment period (titration + evaluation period) within the two treatment groups (x-axis) in Study 5 is presented in Figure 4.

Figure 4: Responder Rate ($\geq 50\%$ Reduction From Baseline) in Study 5



*statistically significant versus placebo

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Levetiracetam extended-release tablets, USP 500 mg are white, oval, biconvex film-coated tablets, engraved with “APO” on one side, “LXR 500” on the other side. They are supplied as follows:

Bottles of 60s (NDC 60505-3280-6)
 Bottles of 500s (NDC 60505-3280-5)
 Bottles of 1,000s (NDC 60505-3280-8)

Levetiracetam extended-release tablets, USP 750 mg are white, capsule shaped, biconvex film-coated tablets, engraved with “APO” on one side, “LXR 750” on the other side. They are supplied as follows:

Bottles of 60s (NDC 60505-3517-6)
 Bottles of 500s (NDC 60505-3517-5)

Levetiracetam extended-release tablets, USP 1,000 mg are white, oval shaped, biconvex film-coated tablets, engraved with “APO” on one side, “LXR 1000” on the other side. They are supplied as follows:

Bottles of 60s (NDC 60505-3518-6)

16.2 Storage

Store at 20°C to 25°C (68°F to 77°F) excursions permitted from 15°C to 30°C (59°F to

86°F) [see USP Controlled Room Temperature]. Dispense in a tight, light-resistant container [see USP]. Protect from moisture.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide). The Medication Guide accompanies the product and can also be accessed on www.apotex.com or by calling Apotex Corp at 1-800-706-5575.

Psychiatric Reactions and Changes in Behavior

Advise patients that levetiracetam extended-release tablets may cause changes in behavior (e.g. irritability and aggression). In addition, patients should be advised that they may experience changes in behavior that have been seen with other formulations of levetiracetam, which include agitation, anger, anxiety, apathy, depression, hostility, and psychotic symptoms [see *Warnings and Precautions (5.1)*].

Suicidal Behavior and Ideation

Counsel patients, their caregivers, and/or families that antiepileptic drugs (AEDs), including levetiracetam extended-release tablets, may increase the risk of suicidal thoughts and behavior and advise patients to be alert for the emergence or worsening of symptoms of depression; unusual changes in mood or behavior; or suicidal thoughts, behavior, or thoughts about self-harm. Advise patients, their caregivers, and/or families to immediately report behaviors of concern to a healthcare provider [see *Warnings and Precautions (5.2)*].

Effects on Driving or Operating Machinery

Inform patients that levetiracetam extended-release tablets may cause dizziness and somnolence. Inform patients not to drive or operate machinery until they have gained sufficient experience on levetiracetam extended-release tablets to gauge whether it adversely affects their ability to drive or operate machinery [see *Warnings and Precautions (5.3)*].

Anaphylaxis and Angioedema

Advise patients to discontinue levetiracetam extended-release tablets and seek medical care if they develop signs and symptoms of anaphylaxis or angioedema [see *Warnings and Precautions (5.4)*].

Dermatological Adverse Reactions

Advise patients that serious dermatological adverse reactions have occurred in patients treated with levetiracetam and instruct them to call their physician immediately if a rash develops [see *Warnings and Precautions (5.5)*].

DRESS/Multiorgan Hypersensitivity

Instruct patients and caregivers that a fever or rash associated with signs of other organ system involvement (e.g., lymphadenopathy, hepatic dysfunction) may be drug-related and should be reported to their healthcare provider immediately. Levetiracetam extended-release tablets should be discontinued immediately if a serious hypersensitivity reaction is suspected [see *Warnings and Precautions (5.6)*].

Dosing and Administration

Patients should be instructed to only take levetiracetam extended-release tablets once daily and to swallow the tablets whole. They should not be chewed, broken, or crushed.

Inform patients that they should not be concerned if they occasionally notice something that looks like swollen pieces of the original tablet in their stool.

Withdrawal of Levetiracetam

Advise patients and caregivers not to discontinue use of levetiracetam extended-release tablets without consulting with their healthcare provider. Levetiracetam extended-release tablets should normally be gradually withdrawn to reduce the potential of increased seizure frequency and status epilepticus [see *Warnings and Precautions (5.8)*].

Pregnancy

Advise patients to notify their healthcare provider if they become pregnant or intend to become pregnant during levetiracetam tablets therapy. Encourage patients to enroll in the North American Antiepileptic Drug (NAAED) pregnancy registry if they become pregnant [see *Use in Specific Populations (8.1)*].

APOTEX INC.

LEVETIRACETAM EXTENDED-RELEASE TABLETS, USP 500 mg, 750 mg and 1,000 mg

Manufactured by	Manufactured for
Apotex Inc.	Apotex Corp.
Toronto, Ontario	Weston, Florida
Canada M9L 1T9	USA 33326

Revision: 14

MEDICATION GUIDE

Levetiracetam (lee" ve tye ra' se tam) Extended-Release Tablets, USP, for oral use

Read this Medication Guide before you start taking levetiracetam extended-release tablets and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or treatment.

What is the most important information I should know about levetiracetam extended-release tablets?

Like other antiepileptic drugs, levetiracetam extended-release tablets may cause suicidal thoughts or actions in a very small number of people, about 1 in 500 people taking it.

Call a healthcare provider right away if you have any of these symptoms, especially if they are new, worse, or worry you:

- thoughts about suicide or dying
- attempts to commit suicide
- new or worse depression

- new or worse anxiety
- feeling agitated or restless
- panic attacks
- trouble sleeping (insomnia)
- new or worse irritability
- acting aggressive, being angry, or violent
- acting on dangerous impulses
- an extreme increase in activity and talking (mania)
- other unusual changes in behavior or mood

Do not stop levetiracetam extended-release tablets without first talking to a healthcare provider.

- Stopping levetiracetam extended-release tablets suddenly can cause serious problems. Stopping a seizure medicine suddenly can cause seizures that will not stop (status epilepticus).
- Suicidal thoughts or actions can be caused by things other than medicines. If you have suicidal thoughts or actions, your healthcare provider may check for other causes.

How can I watch for early symptoms of suicidal thoughts and actions?

- Pay attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings.
- Keep all follow-up visits with your healthcare provider as scheduled.

Call your healthcare provider between visits as needed, especially if you are worried about symptoms.

What are levetiracetam extended-release tablets?

Levetiracetam extended-release tablets are a prescription medicine taken by mouth that is used to treat partial-onset seizures in people 12 years of age and older.

It is not known if levetiracetam extended-release tablets are safe or effective in people under 12 years of age.

Before taking your medicine, make sure you have received the correct medicine. Compare the name above with the name on your bottle and the appearance of your medicine with the description of levetiracetam extended-release tablets provided below. Tell your pharmacist immediately if you think you have been given the wrong medicine.

Who should not take levetiracetam extended-release tablets?

Do not take levetiracetam extended-release tablets if you are allergic to levetiracetam.

What should I tell my healthcare provider before starting levetiracetam extended-release tablets?

Before taking levetiracetam extended-release tablets, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had depression, mood problems or suicidal thoughts or behavior.
- have kidney problems.
- are pregnant or planning to become pregnant. It is not known if levetiracetam extended-release tablets will harm your unborn baby. You and your healthcare

provider will have to decide if you should take levetiracetam extended-release tablets while you are pregnant. If you become pregnant while taking levetiracetam extended-release tablets, talk to your healthcare provider about registering with the North American Antiepileptic Drug Pregnancy Registry. You can enroll in this registry by calling 1-888-233-2334 or go to <http://www.aedpregnancyregistry.org>. The purpose of this registry is to collect information about the safety of levetiracetam extended-release and other antiepileptic medicine during pregnancy.

- are breastfeeding or plan to breastfeed. Levetiracetam can pass into your breast milk. It is not known if the levetiracetam that passes into your breast milk can harm your baby. Talk to your doctor about the best way to feed your baby while you receive levetiracetam extended-release tablets.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Do not start a new medicine without first talking with your healthcare provider.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist each time you get a new medicine.

How should I take levetiracetam extended-release tablets?

- Take levetiracetam extended-release tablets exactly as your healthcare provider tells you to take it.
- Your healthcare provider will tell you how much levetiracetam extended-release tablets to take and when to take it. Levetiracetam extended-release tablets are usually taken 1 time each day.
- Your healthcare provider may change your dose. **Do not** change your dose without talking to your healthcare provider.
- Take levetiracetam extended-release tablets with or without food.
- Swallow the tablets whole. **Do not** chew, break, or crush tablets.
- The inactive part of levetiracetam extended-release tablets may not dissolve after all the medicine has been released in your body. You may sometimes notice something in your bowel movement that looks like swollen pieces of the original tablet. This is normal.
- If you take too much levetiracetam extended-release tablets, call your local Poison Control Center or go to the nearest emergency room right away.

What should I avoid while taking levetiracetam extended-release tablets?

Do not drive, operate machinery or do other dangerous activities until you know how levetiracetam extended-release tablets affect you. Levetiracetam extended-release tablets may make you dizzy or sleepy.

What are the possible side effects of levetiracetam extended-release tablets?

Levetiracetam extended-release tablets can cause serious side effects including:

- See **“What is the most important information I should know about levetiracetam extended-release tablets?”**

Call your healthcare provider right away if you have any of these symptoms:

- mood and behavior changes such as aggression, agitation, anger, anxiety, apathy,

mood swings, depression, hostility, and irritability. A few people may get psychotic symptoms such as hallucinations (seeing or hearing things that are really not there), delusions (false or strange thoughts or beliefs) and unusual behavior.

- extreme sleepiness, tiredness, and weakness.
- allergic reactions such as swelling of the face, lips, eyes, tongue, and throat, trouble swallowing or breathing, and hives.
- a skin rash. Serious skin rashes can happen after you start taking levetiracetam extended-release tablets. There is no way to tell if a mild rash will become a serious reaction.
- a serious allergic reaction that may affect your skin or other parts of your body such as your liver, kidneys, heart, or blood cells. This allergic reaction can be life-threatening and can cause death, particularly if it is not treated as early as possible. Call your healthcare provider right away if you have:
 - a skin rash
 - swelling of your face
 - dark urine
 - fever or swollen glands that do not go away
 - shortness of breath
 - yellowing of the skin or whites of the eyes
- problems with muscle coordination (problems walking and moving).

The most common side effects seen in people who take levetiracetam extended-release tablets include:

- sleepiness
- irritability

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of levetiracetam extended-release tablets. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to Apotex Corp. at 1-800-706-5575 or FDA at 1-800-FDA-1088.

How should I store levetiracetam extended-release tablets?

- Store levetiracetam extended-release tablets at room temperature, 59°F to 86°F (15°C to 30°C) away from heat and light. Protect from moisture.
- **Keep levetiracetam extended-release tablets and all medicines out of the reach of children.**

General information about the safe and effective use of levetiracetam extended-release tablets.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use levetiracetam extended-release tablets for a condition for which it was not prescribed. Do not give levetiracetam extended-release tablets to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about levetiracetam extended-release tablets that is written for health professionals.

What are the ingredients in levetiracetam extended-release tablets?

Levetiracetam extended-release tablet active ingredient: levetiracetam

Inactive ingredients: colloidal silicon dioxide, hydroxypropyl cellulose, hydroxypropyl methylcellulose, polyethylene glycol, and titanium dioxide.

Levetiracetam extended-release tablets do not contain lactose or gluten.

Rx only

This Medication Guide has been approved by the US Food and Drug Administration.

APOTEX INC.

**LEVETIRACETAM EXTENDED-RELEASE TABLETS, USP
500 mg, 750 mg and 1,000 mg**

Manufactured by **Manufactured for**

Apotex Inc. Toronto, Ontario Canada M9L 1T9	Apotex Corp. Weston, Florida USA 33326
---	--

Revised: October 2024

Revision: 14

PRINCIPAL DISPLAY PANEL 500 mg

Representative sample of labeling (see **HOW SUPPLIED** section for complete listing):

PRINCIPAL DISPLAY PANEL - 500 mg BOTTLE LABEL

APOTEX CORP. NDC 60505-3280-6

Levetiracetam Extended-Release Tablets

500 mg

Rx

60 count bottle

Each tablet contains 500 mg of levetiracetam.

Store at 20°C to 25°C (68°F to 77°F); excursions permitted from 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

Protect from moisture.

Dispense in a tight, light-resistant container [see USP].

Usual Dosage:
See package insert.

404161

60 Tablets NDC 60505-3280-6

**Levetiracetam
Extended-Release
Tablets, USP**

500 mg

ONCE DAILY DOSING

**PHARMACIST: Dispense the enclosed
Medication Guide to the patient.**

Rx Only APOTEX CORP.

Manufactured by:
Apotex Inc.
Toronto, Ontario
Canada M9L 1T9
Manufactured for:
Apotex Corp.
Weston, Florida 33326

3 60505 32806 2

Open Here

PRINCIPAL DISPLAY PANEL 750 mg

Representative sample of labeling (see **HOW SUPPLIED** section for complete listing):

PRINCIPAL DISPLAY PANEL - 750 mg BOTTLE LABEL


APOTEX CORP. NDC 60505-3517-6

Levetiracetam Extended-Release Tablets

750 mg

Rx

60 count bottle



PRINCIPAL DISPLAY PANEL 1000 mg

Representative sample of labeling (see **HOW SUPPLIED** section for complete listing):

PRINCIPAL DISPLAY PANEL - 1000 mg BOTTLE LABEL

APOTEX CORP. NDC 60505-3518-6

Levetiracetam Extended-Release Tablets

1000 mg

Rx

60 count bottle

Each tablet contains 1,000 mg of levetiracetam.

Store at 20° to 25°C (68° to 77°F); excursions permitted from 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

Protect from moisture.

Dispense in a tight, light-resistant container [see USP].

Usual Dosage:
See package insert.

000000

60 Tablets

NDC 60505-3518-6

**Levetiracetam
Extended-Release
Tablets, USP**

1,000 mg


ONCE DAILY DOSING

PHARMACIST: Dispense the enclosed Medication Guide to the patient.

R_x Only

APOTEX CORP.

Manufactured by:
Apotex Inc.
Toronto, Ontario
Canada M9L 1T9
Manufactured for:
Apotex Corp.
Weston, Florida 33326



Open Here

LEVETIRACETAM

levetiracetam tablet, film coated, extended release

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:60505-3280
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEVETIRACETAM (UNII: 44YRR34555) (LEVETIRACETAM - UNII:44YRR34555)	LEVETIRACETAM	500 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE 2208 (15000 MPA.S) (UNII: Z78RG6M2N2)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSE 2910 (5 MPA.S) (UNII: R75537T0T4)	
HYDROXYPROPYL CELLULOSE (1600000 WAMW) (UNII: RFW2ET671P)	
POLYETHYLENE GLYCOL 8000 (UNII: Q662QK8M3B)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	APO;LXR;500
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:60505-3280-6	60 in 1 BOTTLE; Type 0: Not a Combination Product	09/12/2011	
2	NDC:60505-3280-5	500 in 1 BOTTLE; Type 0: Not a Combination Product	09/12/2011	
3	NDC:60505-3280-8	1000 in 1 BOTTLE; Type 0: Not a Combination Product	09/12/2011	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091261	09/12/2011	

LEVETIRACETAM

levetiracetam tablet, film coated, extended release

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:60505-3517
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEVETIRACETAM (UNII: 44YRR34555) (LEVETIRACETAM - UNII:44YRR34555)	LEVETIRACETAM	750 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE 2208 (15000 MPA.S) (UNII: Z78RG6M2N2)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSE 2910 (5 MPA.S) (UNII: R75537T0T4)	
HYDROXYPROPYL CELLULOSE (1600000 WAMW) (UNII: RFW2ET671P)	
POLYETHYLENE GLYCOL 8000 (UNII: Q662QK8M3B)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	APO;LXR;750
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60505-	60 in 1 BOTTLE; Type 0: Not a Combination	09/12/2011	

1	3517-6	Product	09/12/2011	
2	NDC:60505-3517-5	500 in 1 BOTTLE; Type 0: Not a Combination Product	09/12/2011	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091261	09/12/2011	

LEVETIRACETAM

levetiracetam tablet, film coated, extended release

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:60505-3518
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
LEVETIRACETAM (UNII: 44YRR34555) (LEVETIRACETAM - UNII:44YRR34555)	LEVETIRACETAM	1000 mg

Inactive Ingredients	
Ingredient Name	Strength
HYPROMELLOSE 2208 (15000 MPA.S) (UNII: Z78RG6M2N2)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSE 2910 (5 MPA.S) (UNII: R75537T0T4)	
HYDROXYPROPYL CELLULOSE (1600000 WAMW) (UNII: RFW2ET671P)	
POLYETHYLENE GLYCOL 8000 (UNII: Q662QK8M3B)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	WHITE	Score	no score
Shape	OVAL	Size	22mm
Flavor		Imprint Code	APO;LXR;1000
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60505-3518-6	60 in 1 BOTTLE; Type 0: Not a Combination Product	02/25/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202958	02/25/2015	

Labeler - Apotex Corp. (845263701)

Registrant - Apotex Inc. (209429182)

Establishment

Name	Address	ID/FEI	Business Operations
Apotex Inc.		209429182	analysis(60505-3280, 60505-3517, 60505-3518) , manufacture(60505-3280, 60505-3517, 60505-3518)

Establishment

Name	Address	ID/FEI	Business Operations
Apotex Inc.		243805095	pack(60505-3280, 60505-3517, 60505-3518)

Revised: 10/2024

Apotex Corp.