MILK OF MAGNESIA- magnesium hydroxide suspension H E B

heb mom org

Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over than 14 days

Ask a doctor or pharmacist before use if you are taking a prescription drug.

This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- follow each dosage with a full glass (8 oz) of liquid

age	dose
adults and children 12 years and over	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

Other information

- each 15 mL contains: magnesium 500 mg
- avoid freezing
- store at room temperature tightly closed

Inactive ingredients

purified water, sodium hypochlorite

Questions or comments?

1-800-540-3765

package Label

Compare to Phillips' @ Milk of Magnesia

NDC 37808-901-12 Gluten Free

Original

Milk of Magnesia

Magnesium Hydroxide

Saline Laxative

- Sugar Free
- Cramp Free
- Stimulant Free

12 FL OZ (355 mL)

Drug Facts

TAMPER EVIDENT: DO NOT USE IF IMPRINTED NECKBAND IS MISSING OR BROKEN.

Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg

Purpose Saline laxative

Use • relieves occasional constipation (irregularity)

generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- · kidney disease
- a magnesium-restricted diet
 - · stomach pain, nausea or vomiting

a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are taking a prescription drug, This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • shake well before use

- do not exceed the maximum recommended daily dose in a 24 hour period dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- follow each dosage with a full glass (8 oz) of liquid

age	dose	
adults and children 12 years and over	30 mL to 60 mL	
children 6 to 11 years	15 mL to 30 mL	
children under 6 years	ask a doctor	

- Other information each 15 mL contains: magnesium 500 mg
- avoid freezing
- · store at room temperature tightly closed

REV GC0820

Inactive ingredients

purified water, sodium hypochlorite.

16269 - 2010

Questions or comments? 1-800-540-3765

*This product is not manufactured or distributed by Bayer Healthcare LLC, the owner of the registered trademark Phillips'® Milk of Magnesia.

MADEWITH PRIDE & CARE FOR H-E-B® SAN ANTONIO, TX 78204





MILK OF MAGNESIA

magnesium hydroxide suspension

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:37808-901

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

Inactive Ingredients			
Ingredient Name	Strength		
WATER (UNII: 059QF0KO0R)			
SODIUM HYPOCHLORITE (UNII: DY38VHM5OD)			

ı	Packaging				
	# Item Code Package Description		Marketing Start Date	Marketing End Date	
	1	NDC:37808-901- 12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2020	

Marketing Information			
Marketing Application Number or Monogra Category Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	08/01/2020	

Labeler - H E B (007924756)

Registrant - GCP Laboratories (965480861)

Establishment					
Name	Address	ID/FEI	Business Operations		
GCP Laboratories		965480861	manufacture(37808-901)		

Revised: 11/2023 H E B