

ACETAMINOPHEN- acetaminophen tablet
BETTER LIVING BRANDS LLC.

Extra Strength

Pain Relief
Acetaminophen USP, 500mg

Pain reliever/ Fever Reducer
FOR ADULTS

RAPID RELEASE

Active ingredient (in each gelcap)

Acetaminophen USP, 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - backache
 - minor pain of arthritis
 - the common cold
 - toothache
 - premenstrual and menstrual cramps
- temporarily reduces fever

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease.

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of the reach of children.

Keep out of the reach of children.

Overdose warning

In case of accidental overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over

- take 2 gelcaps every 6 hours while symptoms last

- do not take more than 6 gelcaps in 24 hours, unless directed by a doctor

- do not take more than 10 days unless directed by a doctor

children under 12 years

- ask a doctor

Other information

- store at 20°-25°C (68°-77°F). See USP Controlled Room Temperature
- avoid high humidity
- see end panel for expiration date and lot number

Inactive ingredients

ammonium hydroxide, black iron oxide, colloidal silicon dioxide, croscarmellose sodium, D&C red #33, FD&C blue #1, FD&C red #40, FD&C yellow #6, gelatin, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, hypromellose, iron oxide red, isopropyl alcohol, n-butyl alcohol, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glaze, stearic acid, titanium dioxide, yellow iron oxide

Questions or comments?

call **1-877-770-3183** Mon-Fri 8:00 AM EST to 5:00 PM PST.

PDP

700000002364
200000003794

IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USE, KEEP THE CARTON FOR IMPORTANT INFORMATION.

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Drug Facts

Active ingredient (in each gelcap) Purpose
Acetaminophen USP, 500 mg Pain Reliever/Fever Reducer

Uses
Temporarily relieves minor aches and pains due to:
headache muscle aches
backache minor pain of arthritis
the common cold toothache
premenstrual and menstrual cramps
temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
more than 4,000 mg of acetaminophen in 24 hours
with other drugs containing acetaminophen
3 or more alcoholic drinks every day while using this product
Heavy alcohol consumption may cause severe liver reactions. Symptoms may include:
skin reddening blisters rash
if a skin reaction occurs, stop use and seek medical help right away.
Do not use
with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
If you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have liver disease.
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
Stop use and ask a doctor if
pain gets worse or lasts more than 10 days
fever gets worse or lasts more than 3 days
new symptoms occur

Drug Facts (continued)

Redness or swelling is present
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
Do not take more than directed (see overdose warning).
Adults and children 12 years and over
Take 2 gelcaps every 6 hours while symptoms last. Do not take more than 6 gelcaps in 24 hours, unless directed by a doctor.
Do not use for more than 10 days unless directed by a doctor.
Children under 12 years
Ask a doctor

Other information
Store at 20°-25°C (68°-77°F). See USP Controlled Room Temperature.
Avoid high humidity.
See and read panel for expiration date and lot number.

Inactive ingredients ammonium hydroxide, black iron oxide, croscarmellose sodium, D&G red #33, FD&C blue #1, FD&C red #40, FD&C yellow #6, gelatin, hydroxypropyl cellulose, hydroxypropyl methylcellulose, hypromellose, iron oxide red, polypropylene glycol, polyethylene glycol, polyvinyl alcohol, polyvinylidene alcohol, polyvinylpyrrolidone, titanium dioxide, yellow iron oxide.

Questions or comments?
Call 1-877-770-3388 Mon-Fri 8:00 AM EST to 5:00 PM PST

*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Extra Strength Tylenol® Rapid Release Gels.



Extra Strength
Pain Relief
ACTAMINOPHEN
USP, 500 mg
Pain Reliever/Fever Reducer
FOR ADULTS
RAPID RELEASE
100 GELCAPS



Compare to
Extra Strength
Tylenol® Rapid
Release Gels
Active Ingredient*
NDC 21130-356-10

PMS 286
CMYK

PMS 424
CMYK

200000003794
700000002364

ASPIRIN FREE
MADE IN INDIA
Scan here for more information
1-877-770-3388
www.signaturecare.com

Lot: _____
Exp. _____

Extra Strength
Pain Relief
ACTAMINOPHEN USP, 500 mg
Pain Reliever/Fever Reducer
FOR ADULTS
RAPID RELEASE
100 GELCAPS

Extra Strength
Pain Relief
ACTAMINOPHEN
USP, 500 mg
Pain Reliever/Fever Reducer
FOR ADULTS
RAPID RELEASE
Actual Size
100 GELCAPS

POVIDONE K30 (UNII: U725QWY32X)
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)
D&C RED NO. 33 (UNII: 9DBA0SBB0L)
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
ISOPROPYL ALCOHOL (UNII: ND2M416302)
BUTYL ALCOHOL (UNII: 8PJ61P6TS3)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
HYPROMELLOSE 2910 (3 MPA.S) (UNII: 0VUT3PMY82)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
SHELLAC (UNII: 46N107B71O)
STEARIC ACID (UNII: 4ELV7Z65AP)
HYDROXYPROPYL CELLULOSE (1600000 WAMW) (UNII: RFW2ET671P)
STARCH, CORN (UNII: O8232NY3SJ)
AMMONIA (UNII: 5138Q19F1X)
FD&C RED NO. 40 (UNII: WZB9127XOA)

Product Characteristics

Color	gray (gray (Encapsulated with red opaque and blue gray opaque hard gelatin shells))	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	G1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-356-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	12/08/2022	
2	NDC:21130-356-24	24 in 1 BOTTLE; Type 0: Not a Combination Product	09/05/2023	
3	NDC:21130-356-21	225 in 1 BOTTLE; Type 0: Not a Combination Product	09/05/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/08/2022	

Labeler - BETTER LIVING BRANDS LLC. (009137209)

