


NYSTATIN- nystatin suspension
Par Pharmaceutical

NYSTATIN ORAL SUSPENSION, USP
100,000 units per mL

Rx only

DESCRIPTION

Nystatin is an antimycotic polyene antibiotic obtained from *Streptomyces noursei*. Structural formula:



$C_{47}H_{75}NO_{17}$

MW = 926.13

Nystatin Oral Suspension is a cherry-mint flavored suspension for oral administration. It contains 100,000 USP Nystatin Units per mL. Inactive ingredients: alcohol ($\leq 1\%$ v/v), carboxymethylcellulose sodium, flavor, glycerin, methylparaben, propylparaben, purified water, saccharin sodium, sodium citrate, and sucrose (50% w/v). May also contain citric acid for pH adjustment.

CLINICAL PHARMACOLOGY

Pharmacokinetics

Gastrointestinal absorption of nystatin is insignificant. Most orally administered nystatin is passed unchanged in the stool. In patients with renal insufficiency receiving oral therapy with conventional dosage forms, significant plasma concentrations of nystatin may occasionally occur.

Microbiology

Nystatin is both fungistatic and fungicidal *in vitro* against a wide variety of yeasts and yeast-like fungi. *Candida albicans* demonstrates no significant resistance to nystatin *in vitro* on repeated subculture in increasing levels of nystatin; other *Candida* species become quite resistant. Generally, resistance does not develop *in vivo*. Nystatin acts by binding to sterols in the cell membrane of susceptible *Candida* species with a resultant change in membrane permeability allowing leakage of intracellular components. Nystatin exhibits no appreciable activity against bacteria, protozoa, or viruses.

INDICATIONS AND USAGE

Nystatin Oral Suspension is indicated for the treatment of candidiasis in the oral cavity.

CONTRAINDICATIONS

The preparation is contraindicated in patients with a history of hypersensitivity to any of its components.

PRECAUTIONS

General

This medication is not to be used for the treatment of systemic mycoses.

Discontinue treatment if sensitization or irritation is reported during use.

Carcinogenesis, Mutagenesis, Impairment of Fertility

No long-term animal studies have been performed to evaluate carcinogenic potential. There also have been no studies to determine mutagenicity or whether this medication affects fertility in males or females.

Pregnancy:

Teratogenic Effects Category C.

Animal reproduction studies have not been conducted with nystatin oral suspension. It is also not known whether nystatin oral suspension can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Nystatin oral suspension should be given to a pregnant woman only if clearly needed.

Nursing Mothers

It is not known whether nystatin is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when nystatin is administered to a nursing woman.

Pediatric Use

See **DOSAGE AND ADMINISTRATION**.

ADVERSE REACTIONS

Nystatin is well tolerated even with prolonged therapy. Oral irritation and sensitization have been reported (See **PRECAUTIONS: General**).

Gastrointestinal: Diarrhea (including one case of bloody diarrhea), nausea, vomiting, gastrointestinal upset/disturbances.

Dermatologic: Rash, including urticaria has been reported rarely. Stevens-Johnson syndrome has been reported very rarely.

Other: Tachycardia, bronchospasm, facial swelling, and nonspecific myalgia have also been reported.

OVERDOSAGE

Oral doses of nystatin in excess of five million units daily have caused nausea and gastrointestinal upset. There have been no reports of serious toxic effects or superinfections (See **CLINICAL PHARMACOLOGY: Pharmacokinetics**).

DOSAGE AND ADMINISTRATION

INFANTS: 2 mL (200,000 units) four times daily (in infants and young children, use dropper to place one-half of dose in each side of mouth and avoid feeding for 5 to 10 minutes). **NOTE:** Limited clinical

studies in premature and low birth weight infants indicate that 1 mL four times daily is effective.

CHILDREN AND ADULTS: 4–6 mL (400,000 to 600,000 units) four times daily (one-half of dose in each side of mouth). The preparation should be retained in the mouth as long as possible before swallowing.

Continue treatment for at least 48 hours after perioral symptoms have disappeared and cultures demonstrate eradication of *Candida albicans*.

HOW SUPPLIED

Nystatin Oral Suspension, USP, 100,000 USP Nystatin Units per mL, is available as a cherry-mint flavored, light creamy yellow, ready-to-use suspension in 60 mL bottles with calibrated dropper and 473 mL bottles.

Storage

Store at 20°-25°C (68°-77°F) with excursions permitted between 15°-30°C (59°-86°F) [See USP Controlled Room Temperature]; avoid freezing.

Manufactured for:

QUALITEST PHARMACEUTICALS

Huntsville, AL 35811

8083097

R2/15-R1

PRINCIPAL DISPLAY PANEL



PRINCIPAL DISPLAY PANEL

NDC 0603-1481-58
NYSTATIN ORAL
SUSPENSION, USP
100,000 units per mL

SHAKE WELL BEFORE USING

Manufactured for:
QUALITEST PHARMACEUTICALS
HUNTSVILLE, AL 35811
Rev. 2/15 R1
8083097 1480

GTIN 00306031481581



N
3 0603-1481-58 1

Lift Here

NDC 0603-1481-58
NYSTATIN ORAL
SUSPENSION, USP
100,000 units per mL

SHAKE WELL BEFORE USING

EACH mL CONTAINS: 100,000
USP Nystatin Units in a vehicle
containing 50% sucrose. Not more
than 1% alcohol by volume.

USUAL DOSAGE FOR INFANTS:
2 mL (200,000 units) four times
daily (1 mL in each side of mouth).
USUAL DOSAGE FOR CHILDREN
AND ADULTS: See package insert.

DISPENSE in tight, light-resistant
container as defined in the USP.


STORE at 20°- 25°C (68°- 77°F)
with excursions permitted between
15°- 30°C (59°- 86°F) [See USP
Controlled Room Temperature];
avoid freezing.

Rx only

NET: 1 PINT (473 mL)

Qualitest®

PRINCIPAL DISPLAY PANEL



NYSTATIN

nystatin suspension

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0603-1481
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NYSTATIN (UNII: BDF101C72E) (NYSTATIN - UNII: BDF101C72E)	NYSTATIN	100000 [USPU] in 1 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
GLYCERIN (UNII: PDC6A3C0OX)	
METHYLPARABEN (UNII: A2I8C7H9T)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0K00R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SUCROSE (UNII: C151H8M554)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY (cherry-mint flavored)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0603-1481-49	1 in 1 CARTON	06/28/2005	01/31/2019
1		60 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		
2	NDC:0603-1481-58	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/28/2005	04/30/2019

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA065148	06/28/2005	04/30/2019

Labeler - Par Pharmaceutical (011103059)**Establishment**

Name	Address	ID/FEI	Business Operations
Vintage Pharmaceuticals, LLC		825839835	MANUFACTURE(0603-1481)