

**BRILLIA HEALTH COUGH CONTROL- lapine bradykinin immune globulin (12c, 30c, 50c), lapine morphine immune globulin (12c, 30c, 50c), lapine histamine immune globulin (12c, 30c, 50c) tablet  
Hadaf LLC**

*Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.*

-----

**72123-007 Cough Control**

**Active Ingredients**

Lapine bradykinin immune globulin 12C, 30C, 50C Purpose Reduces frequency and intensity of cough

Lapine morphine immune globulin 12C, 30C, 50C Purpose Reduces pain response

Lapine histamine immune globulin 12C, 30C, 50C Purpose.Reduces histamine and inflammation response

**Purpose**

Homeopathic,

Reduces frequency and intensity of cough

Reduces pain responseLapine histamine immune

Reduces histamine and inflammation response

**Uses**

- Suppresses wet and dry cough
- Reduces severity and duration of symptoms, including nasal congestion, cough and sore/itchy throat
- Antihistamine and decongestant/anti-inflammatory
- Reduces sore throat pain

**Warnings**

Stop use and ask a doctor if symptoms persist or worsen.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Allergic (hypersensitivity) reactions may occur in very rare cases.

Do not use if known sensitivity to Brillia Health Cough Control or any of its ingredients exists.

**Stop use**

Stop use and ask a doctor if symptoms persist or worsen.

## **Pregnancy**

If pregnant or breast-feeding, ask a health professional before use.

## **Keep out of reach of children**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

## **Allergies**

Allergic (hypersensitivity) reactions may occur in very rare cases.

## **Do Not Use**

Do not use if known sensitivity to Brillia Health Cough Control or any of its ingredients exists.

## **Directions**

- For adults and children 3 years and older.
- Take orally by holding tablet in the mouth until completely dissolved, without food.
- 1 to 2 tablets 3 times daily.

For severe symptoms, take 4 to 6 times daily for the first 3 days.

## **Other Information**

- Store in original package below 25°C or 77°F.
- Do not use if blister pack is torn or broken.

## **Inactive ingredients**

Isomalt, Microcrystalline cellulose, Magnesium stearate, Citric acid (anhydrous), Aspartame, Sodium saccharin.

20 Tablets NDC: 72123-007-01



<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>ISOMALT</b> (UNII: S870P55O2W)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	score with uneven pieces
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72123-007-01	2 in 1 BOX	07/31/2020	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved homeopathic		07/31/2020	

**Labeler** - Hadaf LLC (094114708)

### Establishment

Name	Address	ID/FEI	Business Operations
Santonika Uab		520214047	manufacture(72123-007)

Revised: 1/2022

Hadaf LLC