

**HYDROGEN PEROXIDE- hydrogen peroxide liquid**  
**ImKor Limited Liability Company**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

3.78L NDC: 77610-202-01

KillClean

hydrogen  
peroxide  
3%

## first aid antiseptic

Manufactured by:

ImKor LLC,

Fountainville, PA 18923

Questions or comments? Call

1-800-815-4686 Monday-Friday 9AM-5PM EST

**MADE IN USA** 

1 GALLON (128 FL OZ) 3.78 L

### Drug Facts

#### Active Ingredient(s)

Hydrogen peroxide 3%..... First aid antiseptic/oral debriding agent

#### Purpose

#### Use(s)

■ first aid to help prevent the risk of infection in minor cuts, scrapes, and burns  
■ aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

#### Warnings

For external use only.

##### Do not use

■ in the eyes or apply over large areas of the body  
■ longer than one week

##### Ask doctor before use if you have

■ deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions For use as a first aid antiseptic:

■ clean affected area  
■ apply small amount of product on affected area 1-3 times a day  
■ may be covered with sterile bandage  
■ if bandaged, let dry first

#### For use as an oral debriding agent (oral rinse):

■ **adults and children 2 years of age and over:** ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor ■ **children under 12 years** should be supervised in the use of this product. ■ **children under 2 years of age:** consult a doctor or dentist

#### Other information

■ keep tightly closed and in a dark place at room temperature.  
■ do not shake bottle. Hold away from face when opening.

#### Inactive ingredients

purified water

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND OR UNDER CAP IS BROKEN OR MISSING.



0 731199 537229

KillClean

hydrogen  
peroxide  
3%

## first aid antiseptic

Manufactured by:

ImKor LLC,

Fountainville, PA 18923

Questions or comments? Call

1-800-815-4686 Monday-Friday 9AM-5PM EST

**MADE IN USA** 

1 GALLON (128 FL OZ) 3.78 L

### Drug Facts

#### Active Ingredient(s)

Hydrogen peroxide 3%..... First aid antiseptic/oral debriding agent

#### Purpose

#### Use(s)

■ first aid to help prevent the risk of infection in minor cuts, scrapes, and burns  
■ aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

#### Warnings

For external use only.

##### Do not use

■ in the eyes or apply over large areas of the body  
■ longer than one week

##### Ask doctor before use if you have

■ deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions For use as a first aid antiseptic:

■ clean affected area  
■ apply small amount of product on affected area 1-3 times a day  
■ may be covered with sterile bandage  
■ if bandaged, let dry first

#### For use as an oral debriding agent (oral rinse):

■ **adults and children 2 years of age and over:** ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor ■ **children under 12 years** should be supervised in the use of this product. ■ **children under 2 years of age:** consult a doctor or dentist

#### Other information

■ keep tightly closed and in a dark place at room temperature.  
■ do not shake bottle. Hold away from face when opening.

#### Inactive ingredients

purified water

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND OR UNDER CAP IS BROKEN OR MISSING.



0 731199 537229

3.78L NDC: 77610-202-01

KillClean

hydrogen  
peroxide  
3%

## first aid antiseptic

Manufactured by:

ImKor LLC,

Fountainville, PA 18923

Questions or comments? Call

1-800-815-4686 Monday-Friday 9AM-5PM EST

**MADE IN USA** 

1 GALLON (128 FL OZ) 3.78 L

### Drug Facts

#### Active Ingredient(s)

Hydrogen peroxide 3%..... First aid antiseptic/oral debriding agent

#### Purpose

#### Use(s)

■ first aid to help prevent the risk of infection in minor cuts, scrapes, and burns  
■ aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

#### Warnings

For external use only.

#### Do not use

■ in the eyes or apply over large areas of the body  
■ longer than one week

#### Ask doctor before use if you have

■ deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions For use as a first aid antiseptic:

■ clean affected area  
■ apply small amount of product on affected area 1-3 times a day  
■ may be covered with sterile bandage  
■ if bandaged, let dry first

#### For use as an oral debriding agent (oral rinse):

■ **adults and children 2 years of age and over:** ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor ■ **children under 12 years** should be supervised in the use of this product. ■ **children under 2 years of age:** consult a doctor or dentist

#### Other information

■ keep tightly closed and in a dark place at room temperature.  
■ do not shake bottle. Hold away from face when opening.

#### Inactive ingredients

purified water

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND OR UNDER CAP IS BROKEN OR MISSING.**



0 731199 537229

KillClean

hydrogen  
peroxide  
3%

## first aid antiseptic

Manufactured by:

ImKor LLC,

Fountainville, PA 18923

Questions or comments? Call

1-800-815-4686 Monday-Friday 9AM-5PM EST

**MADE IN USA** 

1 GALLON (128 FL OZ) 3.78 L

### Drug Facts

#### Active Ingredient(s)

Hydrogen peroxide 3%..... First aid antiseptic/oral debriding agent

#### Purpose

#### Use(s)

■ first aid to help prevent the risk of infection in minor cuts, scrapes, and burns  
■ aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

#### Warnings

For external use only.

#### Do not use

■ in the eyes or apply over large areas of the body  
■ longer than one week

#### Ask doctor before use if you have

■ deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions For use as a first aid antiseptic:

■ clean affected area  
■ apply small amount of product on affected area 1-3 times a day  
■ may be covered with sterile bandage  
■ if bandaged, let dry first

#### For use as an oral debriding agent (oral rinse):

■ **adults and children 2 years of age and over:** ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor ■ **children under 12 years** should be supervised in the use of this product. ■ **children under 2 years of age:** consult a doctor or dentist

#### Other information

■ keep tightly closed and in a dark place at room temperature.  
■ do not shake bottle. Hold away from face when opening.

#### Inactive ingredients

purified water

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND OR UNDER CAP IS BROKEN OR MISSING.**



0 731199 537229

3.78L NDC: 77610-202-01

KillClean

hydrogen  
peroxide  
3%

## first aid antiseptic

Manufactured by:

**ImKor LLC,**

Fountainville, PA 18923

Questions or comments? Call

**1-800-815-4686 Monday-Friday 9AM-5PM EST**

**MADE IN USA** 

1 GALLON (128 FL OZ) 3.78 L

### Drug Facts

#### Active Ingredient(s)

Hydrogen peroxide 3%..... First aid antiseptic/oral debriding agent

#### Purpose

#### Use(s)

■ first aid to help prevent the risk of infection in minor cuts, scrapes, and burns  
■ aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

#### Warnings

**For external use only.**

#### Do not use

■ in the eyes or apply over large areas of the body  
■ longer than one week

#### Ask doctor before use if you have

■ deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions For use as a first aid antiseptic:

■ clean affected area  
■ apply small amount of product on affected area 1-3 times a day  
■ may be covered with sterile bandage  
■ if bandaged, let dry first

#### For use as an oral debriding agent (oral rinse):

■ **adults and children 2 years of age and over:** ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor ■ **children under 12 years** should be supervised in the use of this product. ■ **children under 2 years of age:** consult a doctor or dentist

#### Other information

■ keep tightly closed and in a dark place at room temperature.  
■ do not shake bottle. Hold away from face when opening.

#### Inactive ingredients

purified water

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND OR UNDER CAP IS BROKEN OR MISSING.**



0 731199 537229

KillClean

hydrogen  
peroxide  
3%

## first aid antiseptic

Manufactured by:

**ImKor LLC,**

Fountainville, PA 18923

Questions or comments? Call

**1-800-815-4686 Monday-Friday 9AM-5PM EST**

**MADE IN USA** 

1 GALLON (128 FL OZ) 3.78 L

### Drug Facts

#### Active Ingredient(s)

Hydrogen peroxide 3%..... First aid antiseptic/oral debriding agent

#### Purpose

#### Use(s)

■ first aid to help prevent the risk of infection in minor cuts, scrapes, and burns  
■ aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

#### Warnings

**For external use only.**

#### Do not use

■ in the eyes or apply over large areas of the body  
■ longer than one week

#### Ask doctor before use if you have

■ deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions For use as a first aid antiseptic:

■ clean affected area  
■ apply small amount of product on affected area 1-3 times a day  
■ may be covered with sterile bandage  
■ if bandaged, let dry first

#### For use as an oral debriding agent (oral rinse):

■ **adults and children 2 years of age and over:** ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor ■ **children under 12 years** should be supervised in the use of this product. ■ **children under 2 years of age:** consult a doctor or dentist

#### Other information

■ keep tightly closed and in a dark place at room temperature.  
■ do not shake bottle. Hold away from face when opening.

#### Inactive ingredients

purified water

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND OR UNDER CAP IS BROKEN OR MISSING.**



0 731199 537229

3.78L NDC: 77610-202-01



KillClean

hydrogen  
peroxide  
3%

## first aid antiseptic

Manufactured by:

**ImKor LLC,**

Fountainville, PA 18923

Questions or comments? Call

**1-800-815-4686 Monday-Friday 9AM-5PM EST**

**MADE IN USA** 

1 GALLON (128 FL OZ) 3.78 L

### Drug Facts

#### Active Ingredient(s)

Hydrogen peroxide 3%..... First aid antiseptic/oral debriding agent

#### Purpose

#### Use(s)

- first aid to help prevent the risk of infection in minor cuts, scrapes, and burns
- aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

#### Warnings

For external use only.

#### Do not use

- in the eyes or apply over large areas of the body
- longer than one week

#### Ask doctor before use if you have

- deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions For use as a first aid antiseptic:

- clean affected area
- apply small amount of product on affected area 1-3 times a day
- may be covered with sterile bandage
- if bandaged, let dry first

#### For use as an oral debriding agent (oral rinse):

■ **adults and children 2 years of age and over:** ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor ■ **children under 12 years** should be supervised in the use of this product. ■ **children under 2 years of age:** consult a doctor or dentist

#### Other information

- keep tightly closed and in a dark place at room temperature.
- do not shake bottle. Hold away from face when opening.

#### Inactive ingredients

purified water

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND OR UNDER CAP IS BROKEN OR MISSING.**



0 731199 537229

KillClean

hydrogen  
peroxide  
3%

## first aid antiseptic

Manufactured by:

**ImKor LLC,**

Fountainville, PA 18923

Questions or comments? Call

**1-800-815-4686 Monday-Friday 9AM-5PM EST**

**MADE IN USA** 

1 GALLON (128 FL OZ) 3.78 L

### Drug Facts

#### Active Ingredient(s)

Hydrogen peroxide 3%..... First aid antiseptic/oral debriding agent

#### Purpose

#### Use(s)

- first aid to help prevent the risk of infection in minor cuts, scrapes, and burns
- aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

#### Warnings

For external use only.

#### Do not use

- in the eyes or apply over large areas of the body
- longer than one week

#### Ask doctor before use if you have

- deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions For use as a first aid antiseptic:

- clean affected area
- apply small amount of product on affected area 1-3 times a day
- may be covered with sterile bandage
- if bandaged, let dry first

#### For use as an oral debriding agent (oral rinse):

■ **adults and children 2 years of age and over:** ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor ■ **children under 12 years** should be supervised in the use of this product. ■ **children under 2 years of age:** consult a doctor or dentist

#### Other information

- keep tightly closed and in a dark place at room temperature.
- do not shake bottle. Hold away from face when opening.

#### Inactive ingredients

purified water

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND OR UNDER CAP IS BROKEN OR MISSING.**



0 731199 537229

3.78L NDC: 77610-202-01

KillClean

hydrogen  
peroxide  
3%

## first aid antiseptic

Manufactured by:

**ImKor LLC,**

Fountainville, PA 18923

Questions or comments? Call

**1-800-815-4686 Monday-Friday 9AM-5PM EST**

**MADE IN USA** 

1 GALLON (128 FL OZ) 3.78 L

### Drug Facts

#### Active Ingredient(s)

Hydrogen peroxide 3%..... First aid antiseptic/oral debriding agent

#### Purpose

#### Use(s)

■ first aid to help prevent the risk of infection in minor cuts, scrapes, and burns  
■ aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

#### Warnings

**For external use only.**

#### Do not use

■ in the eyes or apply over large areas of the body  
■ longer than one week

#### Ask doctor before use if you have

■ deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions For use as a first aid antiseptic:

■ clean affected area  
■ apply small amount of product on affected area 1-3 times a day  
■ may be covered with sterile bandage  
■ if bandaged, let dry first

#### For use as an oral debriding agent (oral rinse):

■ **adults and children 2 years of age and over:** ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor ■ **children under 12 years** should be supervised in the use of this product. ■ **children under 2 years of age:** consult a doctor or dentist

#### Other information

■ keep tightly closed and in a dark place at room temperature.  
■ do not shake bottle. Hold away from face when opening.

#### Inactive ingredients

purified water

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND OR UNDER CAP IS BROKEN OR MISSING.**



0 731199 537229

KillClean

hydrogen  
peroxide  
3%

## first aid antiseptic

Manufactured by:

**ImKor LLC,**

Fountainville, PA 18923

Questions or comments? Call

**1-800-815-4686 Monday-Friday 9AM-5PM EST**

**MADE IN USA** 

1 GALLON (128 FL OZ) 3.78 L

### Drug Facts

#### Active Ingredient(s)

Hydrogen peroxide 3%..... First aid antiseptic/oral debriding agent

#### Purpose

#### Use(s)

■ first aid to help prevent the risk of infection in minor cuts, scrapes, and burns  
■ aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

#### Warnings

**For external use only.**

#### Do not use

■ in the eyes or apply over large areas of the body  
■ longer than one week

#### Ask doctor before use if you have

■ deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions For use as a first aid antiseptic:

■ clean affected area  
■ apply small amount of product on affected area 1-3 times a day  
■ may be covered with sterile bandage  
■ if bandaged, let dry first

#### For use as an oral debriding agent (oral rinse):

■ **adults and children 2 years of age and over:** ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor ■ **children under 12 years** should be supervised in the use of this product. ■ **children under 2 years of age:** consult a doctor or dentist

#### Other information

■ keep tightly closed and in a dark place at room temperature.  
■ do not shake bottle. Hold away from face when opening.

#### Inactive ingredients

purified water

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND OR UNDER CAP IS BROKEN OR MISSING.**



0 731199 537229

3.78L NDC: 77610-202-01

KillClean

hydrogen  
peroxide  
3%

## first aid antiseptic

Manufactured by:

ImKor LLC,

Fountainville, PA 18923

Questions or comments? Call

1-800-815-4686 Monday-Friday 9AM-5PM EST

**MADE IN USA** 

1 GALLON (128 FL OZ) 3.78 L

### Drug Facts

#### Active Ingredient(s)

Hydrogen peroxide 3%..... First aid antiseptic/oral debriding agent

#### Purpose

#### Use(s)

■ first aid to help prevent the risk of infection in minor cuts, scrapes, and burns  
■ aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

#### Warnings

For external use only.

#### Do not use

■ in the eyes or apply over large areas of the body  
■ longer than one week

#### Ask doctor before use if you have

■ deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions For use as a first aid antiseptic:

■ clean affected area  
■ apply small amount of product on affected area 1-3 times a day  
■ may be covered with sterile bandage  
■ if bandaged, let dry first

#### For use as an oral debriding agent (oral rinse):

■ **adults and children 2 years of age and over:** ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor ■ **children under 12 years** should be supervised in the use of this product. ■ **children under 2 years of age:** consult a doctor or dentist

#### Other information

■ keep tightly closed and in a dark place at room temperature.  
■ do not shake bottle. Hold away from face when opening.

#### Inactive ingredients

purified water

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND OR UNDER CAP IS BROKEN OR MISSING.



0 731199 537229

KillClean

hydrogen  
peroxide  
3%

## first aid antiseptic

Manufactured by:

ImKor LLC,

Fountainville, PA 18923

Questions or comments? Call

1-800-815-4686 Monday-Friday 9AM-5PM EST

**MADE IN USA** 

1 GALLON (128 FL OZ) 3.78 L

### Drug Facts

#### Active Ingredient(s)

Hydrogen peroxide 3%..... First aid antiseptic/oral debriding agent

#### Purpose

#### Use(s)

■ first aid to help prevent the risk of infection in minor cuts, scrapes, and burns  
■ aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

#### Warnings

For external use only.

#### Do not use

■ in the eyes or apply over large areas of the body  
■ longer than one week

#### Ask doctor before use if you have

■ deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions For use as a first aid antiseptic:

■ clean affected area  
■ apply small amount of product on affected area 1-3 times a day  
■ may be covered with sterile bandage  
■ if bandaged, let dry first

#### For use as an oral debriding agent (oral rinse):

■ **adults and children 2 years of age and over:** ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor ■ **children under 12 years** should be supervised in the use of this product. ■ **children under 2 years of age:** consult a doctor or dentist

#### Other information

■ keep tightly closed and in a dark place at room temperature.  
■ do not shake bottle. Hold away from face when opening.

#### Inactive ingredients

purified water

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND OR UNDER CAP IS BROKEN OR MISSING.



0 731199 537229

3.78L NDC: 77610-202-01



KillClean

hydrogen  
peroxide  
3%

## first aid antiseptic

Manufactured by:

ImKor LLC,

Fountainville, PA 18923

Questions or comments? Call

1-800-815-4686 Monday-Friday 9AM-5PM EST

**MADE IN USA** 

1 GALLON (128 FL OZ) 3.78 L

### Drug Facts

#### Active Ingredient(s)

Hydrogen peroxide 3%..... First aid antiseptic/oral debriding agent

#### Purpose

#### Use(s)

- first aid to help prevent the risk of infection in minor cuts, scrapes, and burns
- aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

#### Warnings

For external use only.

##### Do not use

- in the eyes or apply over large areas of the body
- longer than one week

##### Ask doctor before use if you have

- deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions For use as a first aid antiseptic:

- clean affected area
- apply small amount of product on affected area 1-3 times a day
- may be covered with sterile bandage
- if bandaged, let dry first

#### For use as an oral debriding agent (oral rinse):

■ **adults and children 2 years of age and over:** ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor ■ **children under 12 years** should be supervised in the use of this product. ■ **children under 2 years of age:** consult a doctor or dentist

#### Other information

- keep tightly closed and in a dark place at room temperature.
- do not shake bottle. Hold away from face when opening.

**Inactive ingredients** purified water

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND OR UNDER CAP IS BROKEN OR MISSING.



0 731199 537229

KillClean

hydrogen  
peroxide  
3%

## first aid antiseptic

Manufactured by:

ImKor LLC,

Fountainville, PA 18923

Questions or comments? Call

1-800-815-4686 Monday-Friday 9AM-5PM EST

**MADE IN USA** 

1 GALLON (128 FL OZ) 3.78 L

### Drug Facts

#### Active Ingredient(s)

Hydrogen peroxide 3%..... First aid antiseptic/oral debriding agent

#### Purpose

#### Use(s)

- first aid to help prevent the risk of infection in minor cuts, scrapes, and burns
- aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

#### Warnings

For external use only.

##### Do not use

- in the eyes or apply over large areas of the body
- longer than one week

##### Ask doctor before use if you have

- deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions For use as a first aid antiseptic:

- clean affected area
- apply small amount of product on affected area 1-3 times a day
- may be covered with sterile bandage
- if bandaged, let dry first

#### For use as an oral debriding agent (oral rinse):

■ **adults and children 2 years of age and over:** ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor ■ **children under 12 years** should be supervised in the use of this product. ■ **children under 2 years of age:** consult a doctor or dentist

#### Other information

- keep tightly closed and in a dark place at room temperature.
- do not shake bottle. Hold away from face when opening.

**Inactive ingredients** purified water

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND OR UNDER CAP IS BROKEN OR MISSING.



0 731199 537229

3.78 L NDC: 77610-202-01



KillClean

hydrogen  
peroxide  
3%

## first aid antiseptic

Manufactured by:

**ImKor LLC,**

Fountainville, PA 18923

Questions or comments? Call

**1-800-815-4686 Monday-Friday 9AM-5PM EST**

**MADE IN USA** 

1 GALLON (128 FL OZ) 3.78 L

### Drug Facts

#### Active Ingredient(s)

Hydrogen peroxide 3%..... First aid antiseptic/oral debriding agent

#### Purpose

#### Use(s)

■ first aid to help prevent the risk of infection in minor cuts, scrapes, and burns  
■ aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

#### Warnings

**For external use only.**

#### Do not use

■ in the eyes or apply over large areas of the body  
■ longer than one week

#### Ask doctor before use if you have

■ deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions For use as a first aid antiseptic:

■ clean affected area  
■ apply small amount of product on affected area 1-3 times a day  
■ may be covered with sterile bandage  
■ if bandaged, let dry first

#### For use as an oral debriding agent (oral rinse):

■ **adults and children 2 years of age and over:** ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor ■ **children under 12 years** should be supervised in the use of this product. ■ **children under 2 years of age:** consult a doctor or dentist

#### Other information

■ keep tightly closed and in a dark place at room temperature.  
■ do not shake bottle. Hold away from face when opening.

#### Inactive ingredients

purified water

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND OR UNDER CAP IS BROKEN OR MISSING.**



KillClean

hydrogen  
peroxide  
3%

## first aid antiseptic

Manufactured by:

**ImKor LLC,**

Fountainville, PA 18923

Questions or comments? Call

**1-800-815-4686 Monday-Friday 9AM-5PM EST**

**MADE IN USA** 

1 GALLON (128 FL OZ) 3.78 L

### Drug Facts

#### Active Ingredient(s)

Hydrogen peroxide 3%..... First aid antiseptic/oral debriding agent

#### Purpose

#### Use(s)

■ first aid to help prevent the risk of infection in minor cuts, scrapes, and burns  
■ aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

#### Warnings

**For external use only.**

#### Do not use

■ in the eyes or apply over large areas of the body  
■ longer than one week

#### Ask doctor before use if you have

■ deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions For use as a first aid antiseptic:

■ clean affected area  
■ apply small amount of product on affected area 1-3 times a day  
■ may be covered with sterile bandage  
■ if bandaged, let dry first

#### For use as an oral debriding agent (oral rinse):

■ **adults and children 2 years of age and over:** ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor ■ **children under 12 years** should be supervised in the use of this product. ■ **children under 2 years of age:** consult a doctor or dentist

#### Other information

■ keep tightly closed and in a dark place at room temperature.  
■ do not shake bottle. Hold away from face when opening.

#### Inactive ingredients

purified water

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND OR UNDER CAP IS BROKEN OR MISSING.**



473 mL NDC: 77610-202-16

KillClean

NDC 77610-202-16

# 3% HYDROGEN PEROXIDE

## first aid antiseptic

Manufactured by:

**ImKor LLC,**

Fountainville, PA 18923

Questions or comments? Call

**1-800-815-4686 Monday-Friday 9AM-5PM EST**

**MADE IN USA** 

16 FL OZ (473 mL)

946 mL NDC: 77610-202-32

### Drug Facts

#### Active Ingredient(s)

Hydrogen peroxide 3%..... First aid antiseptic/oral debriding agent

#### Purpose

#### Use(s)

- first aid to help prevent the risk of infection in minor cuts, scrapes, and burns
- aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

#### Warnings

For external use only.

#### Do not use

- in the eyes or apply over large areas of the body
- longer than one week

#### Ask doctor before use if you have

- deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions For use as a first aid antiseptic:

- clean affected area
- apply small amount of product on affected area 1-3 times a day
- may be covered with sterile bandage
- if bandaged, let dry first

#### For use as an oral debriding agent (oral rinse):

- **adults and children 2 years of age and over:** ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist.
- **children under 12 years** should be supervised in the use of this product.
- **children under 2 years of age:** consult a doctor or dentist

#### Other information

- keep tightly closed and in a dark place at room temperature.
- do not shake bottle. Hold away from face when opening.

**Inactive ingredients** purified water

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND OR UNDER CAP IS BROKEN OR MISSING.**



KillClean

NDC 77610-202-32

# 3% HYDROGEN PEROXIDE

## first aid antiseptic

Manufactured by:  
**ImKor LLC**,  
Fountainville, PA 18923  
Questions or comments? Call  
**1-800-815-4686 Monday-Friday 9AM-5PM EST**  
**MADE IN USA** 

32 FL OZ (946 mL)

### Drug Facts

**Active Ingredient(s)** Hydrogen peroxide 3%..... **Purpose** First aid antiseptic/oral debriding agent

### Use(s)

- first aid to help prevent the risk of infection in minor cuts, scrapes, and burns
- aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

### Warnings

For external use only.

#### Do not use

- in the eyes or apply over large areas of the body
- longer than one week

#### Ask doctor before use if you have

- deep puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens ■ swelling, rash, or fever develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### Directions

For use as a first aid antiseptic:

- clean affected area
- apply small amount of product on affected area 1-3 times a day
- may be covered with sterile bandage
- if bandaged, let dry first

For use as an oral debriding agent (oral rinse):

- adults and children 2 years of age and over: ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist.
- children under 12 years should be supervised in the use of this product.
- children under 2 years of age: consult a doctor or dentist

### Other information

- keep tightly closed and in a dark place at room temperature.
- do not shake bottle. Hold away from face when opening.

**Inactive ingredients** purified water

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**



## HYDROGEN PEROXIDE

hydrogen peroxide liquid

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:77610-202
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROGEN PEROXIDE (UNII: BBX060AN9V) (HYDROGEN PEROXIDE - UNII:BBX060AN9V)	HYDROGEN PEROXIDE	0.03 L in 0.03 L

### Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start	Marketing End
---	-----------	---------------------	-----------------	---------------

#	Item Code	Package Description	Date	Date
1	NDC:77610-202-01	1 L in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/13/2020	
2	NDC:77610-202-16	0.473 L in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/13/2020	
3	NDC:77610-202-32	0.946 L in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/13/2020	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	08/13/2020	

**Labeler** - ImKor Limited Liability Company (094772617)

### Establishment

Name	Address	ID/FEI	Business Operations
ImKor Limited Liability Company		094772617	manufacture(77610-202)

Revised: 9/2020

ImKor Limited Liability Company