

NIX- permethrin shampoo
Central Texas Community Health Centers

NIX®

Drug Facts

Active ingredient (in each fluid ounce)

Permethrin 280 mg (1%)

Purpose

Lice treatment

Use

treats head lice

Warnings

For external use only

Do not use

- on children under 2 months of age
- near the eyes
- inside the nose, ear, mouth, or vagina
- on lice in eyebrows or eyelashes. See your doctor.

Ask a doctor before use if you are

allergic to ragweed. May cause breathing difficulty or an asthmatic episode.

When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets into the eyes, immediately flush with large amounts of water
- scalp itching or redness may occur

Stop use and see a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Inspect

- all household members should be checked by another person for lice and/or nits (eggs)
- use a magnifying glass in bright light to help you see the lice and nits (eggs)
- use a tool, such as a comb or two unsharpened pencils to lift and part the hair
- look for tiny nits near the scalp, beginning at the back of the neck and behind the ears
- small sections of hair (1-2 inches wide) should be examined at a time
- unlike dandruff, nits stick to the hair. Dandruff should move when lightly touched.
- if either lice or nits (eggs) are found, treat with Nix[®] Creme Rinse

Treat

- wash hair with a shampoo without conditioner. Do not use a shampoo that contains a conditioner or a conditioner alone since this may decrease the activity of Nix[®]. Rinse with water.
- towel dry hair so it is damp but not wet
- shake the bottle of Nix[®] well
- completely saturate the hair and scalp with Nix[®]. Begin to apply Nix[®] behind the ears and at the back of the neck.
- keep Nix[®] out of the eyes. Protect the eyes with a washcloth or towel.
- leave Nix[®] on the hair for 10 minutes, but no longer
- rinse with warm water
- towel dry hair and comb out tangles
- if live lice are seen seven days or more after the first treatment, a second treatment should be given

Remove Lice/Nits

- remove nits by combing the hair with the special small tooth comb provided. Remaining nits may be removed by hand (using a throw-away glove), or cutting the nits out.
- use the nit comb provided and make sure the hair remains slightly damp while removing nits
- if the hair dries during combing, dampen it slightly with water
- part the hair into 4 sections. Work on one section at a time. Longer hair may take more time (1-2 hours).
- start at the top of the head on the section you have picked
- with one hand, lift a 1-2 inch wide strand of hair. Get the teeth of the comb as close to the scalp as possible and comb with a firm, even motion away from the scalp to the end of the hair.
- use clips to pin back each strand of hair after you have combed out the nits
- clean the comb completely as you go. Wipe the nits from the comb with a tissue and throw away the tissue in a sealed plastic bag to prevent the lice from coming back.
- after combing, recheck the entire head for nits and repeat combing if necessary
- check the affected head daily to remove any nits that you might have missed

Other information

- read all the directions in the Consumer Information Insert and warnings before use. Keep the carton. It contains important information.
- store at 20° to 25°C (68° to 77°F)

Inactive ingredients

balsam canada, cetyl alcohol, citric acid, FD&C yellow no. 6, fragrance, hydrolyzed animal protein, hydroxyethylcellulose, polyoxyethylene 10 cetyl ether, propylene glycol, stearylalkonium chloride, water, isopropyl alcohol, methylparaben, and propylparaben

Questions or comments?

call 1-888-LICE LINE (1-888-542-3546), Monday to Friday, 9 AM - 5 PM EST


Distributed by:
INSIGHT Pharmaceuticals LLC.
Tarrytown, NY 10591
A Prestige Brands Company

HOW SUPPLIED

Product: 76413-358

NDC: 76413-358-01 1 mL in a CARTON

NIX (PERMETHRIN) SHAMPOO



NIX

permethrin shampoo

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:76413-358(NDC:63736-024)
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PERMETHRIN (UNII: 509F88P9SZ) (PERMETHRIN - UNII:509F88P9SZ)	PERMETHRIN	1 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
CANADA BALSAM (UNII: K304VI98B)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYDROXYETHYL CELLULOSE (2000 MPA.S AT 1%) (UNII: S38J6RZN16)	
CETETH-10 (UNII: LF9X1PN3XJ)	

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARALKONIUM CHLORIDE (UNII: 0OUO26BB88)	
WATER (UNII: 059QF0K00R)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	

Product Characteristics			
Color	ORANGE (Light Orange)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76413-358-01	1 mL in 1 CARTON; Type 0: Not a Combination Product	01/15/2019	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA019918	11/15/2013	

Labeler - Central Texas Community Health Centers (079674019)

Establishment			
Name	Address	ID/FEI	Business Operations
Central Texas Community Health Centers		079674019	RELABEL(76413-358) , REPACK(76413-358)

Revised: 1/2019

Central Texas Community Health Centers