

**BENZETHONIUM CHLORIDE AND DYCLONINE HYDROCHLORIDE- liquid  
bandage liquid  
Rite Aid Corporation**

-----  
**Liquid Bandage 1 fl oz (30 mL)**

***Active ingredient***

Benzethonium chloride 0.2%

Dyclonine hydrochloride 0.75%

***Purpose***

First aid antiseptic

Topical analgesic

***Uses***

- first aid to prevent bacterial contamination in minor cuts, scrapes and burns
- for temporary relief of pain

***Warnings***

**For external use only.**

**Flammable:**

Keep away from fire or flame.

**Do not use**

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor

**Ask a doctor before use if you have**

- deep or puncture wounds
- deep cuts
- animal bites
- serious burns

**When using this product**

- avoid contact with eyes

**Stop use and ask a doctor if**

- condition persists or gets worse
- symptoms persists for more than 7 days or clear up and occur again within a few

days

- infection occurs

### **Keep out of reach of children.**

If swallowed, get medical help or call a Poison Control Center right away.

### **Directions**

- for adults and children 2 years of age or older
- clean affected area
- apply a small amount on the area 1 to 3 times daily and let dry
- a second coating may be applied for extra protection
- children under 2 years of age: consult a doctor

### **Other information**

- cap bottle tightly
- store between 20° to 25°C (68° to 77°F)
- may temporarily sting upon application
- to remove, apply more Liquid Bandage and quickly wipe off

### **Inactive ingredients**

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

### **Questions?**

**Call 1-866-964-0939**

### **Principal Display Panel**

Rite Aid

### **LIQUID BANDAGE**

### **TOPICAL ANALGESIC & ANTISEPTIC**

For small cuts & wounds

To help prevent infection

Antiseptic formula dries to form an invisible protective cover that is flexible

NET 1.0 FL OZ (29.6 mL)

Use Liquid Bandage for blisters, chapped & cracked fingertips, hangnails, paper cuts, shaving nicks, & help in preventing the formulation of calluses

Great for: athletes, active lifestyles, outdoor sportsmen & musicians



# LIQUID BANDAGE

TOPICAL ANALGESIC & ANTISEPTIC

For small cuts & wounds

To help prevent infection

Antiseptic formula dries to form an invisible protective cover that is flexible

NET 1.0 FL OZ (29.6 mL)

# LIQUID BANDAGE

TOPICAL ANALGESIC & ANTISEPTIC

Use Liquid Bandage for: blisters, chapped & cracked fingertips, hangnails, paper cuts, shaving nicks & help in preventing the formation of calluses

Great for: athletes, active lifestyles, outdoor sportsmen & musicians

53-033RA-B01

# LIQUID BANDAGE

TOPICAL ANALGESIC & ANTISEPTIC

For small cuts & wounds  
To help prevent infection

# LIQUID BANDAGE

TOPICAL ANALGESIC & ANTISEPTIC

For small cuts & wounds  
To help prevent  
infection.



PHARMACY

NET 1.0 FL OZ (29.6 mL)

DISTRIBUTED BY: RITE AID  
30 HUNTER LANE  
CAMP HILL, PA 17011  
www.riteaid.com  
MADE IN USA WITH U.S.  
AND IMPORTED PARTS

SATISFACTION  
GUARANTEE

If you're not satisfied, we'll  
happily refund your money.



0 11822 78323 1

Care ■ Wipe top of bottle clean after each use.

Questions? Call 1-866-964-0939

Inactive ingredients: acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

more Liquid Bandage and quickly wipe off

Other information ■ cap bottle tightly ■ store between 20° to 25°C (68° to 77°F) ■ may temporarily sting upon application ■ to remove, apply

■ children under 2 years of age: consult a doctor

■ a second coating may be applied for extra protection


Directions ■ for adults and children 2 years of age and older ■ clean affected area ■ apply a small amount on the area 1 to 3 times daily and let dry

Control Center right away.

Keep out of reach of children. If swallowed, get medical help or call a Poison

few days ■ infection occurs

■ symptoms persist for more than 7 days or clear up and occur again within a



**DO NOT USE IF NECKBAND IS BROKEN**

**Active ingredients:** Benzethonium Chloride 0.2% and Dyclonine Hydrochloride 0.75%. **Purpose:** First aid antiseptic and topical analgesic. **Uses:** First aid to prevent bacterial contamination in minor cuts, scrapes, and burns. For the temporary relief of pain. **Warnings:** For external use only. **Flammable:** Keep away from fire or flame. Do not use in eyes, over large areas of the body, and longer than 1 week unless directed by a doctor.

# LIQUID BANDAGE

**TOPICAL ANALGESIC & ANTISEPTIC**

For small cuts & wounds  
To help prevent infection

NET 1.0 FL OZ (29.6 mL)

**Ask a doctor before use if you have** deep or puncture wounds, deep cuts, animal bites and serious burns.

**When using this product** avoid contact with eyes.

**Stop use and ask a doctor** if condition persists or gets worse, symptoms persist for more than 7 days or clear up and occur again within a few days, and infection occurs.

**Keep out of reach of children.** If swallowed, get medical help or call a Poison Control Center right away.

**Directions:** For adults and children 2 years of age and older. Clean affected area. Apply a small amount on the area 1 to 3 times daily and let dry. A second coating may be applied for extra protection. Children under 2 years of age: consult a doctor.

**Other information:** Cap bottle tightly. Store between 20° to 25°C (68° to 77°F). May temporarily sting upon application. To remove, apply more Liquid Bandage and quickly wipe off.

**Inactive ingredients:** acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

**Questions? Call 1-866-964-0939**

**DISTRIBUTED BY:** RITE AID, 30 HUNTER LANE  
CAMP HILL, PA 17011 [www.riteaid.com](http://www.riteaid.com)

**MADE IN USA WITH U.S. AND IMPORTED PARTS**

**SATISFACTION GUARANTEE:** if you're not satisfied, we'll happily refund your money. 50-033RA-B01

## BENZETHONIUM CHLORIDE AND DYCLONINE HYDROCHLORIDE

liquid bandage liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-0779
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BENZETHONIUM CHLORIDE</b> (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	0.2 mg in 30 mL
<b>DYCLONINE HYDROCHLORIDE</b> (UNII: ZEC193879Q) (DYCLONINE -	DYCLONINE	0.75 mg

UNII:078A24Q300)

HYDROCHLORIDE

in 30 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>PYROXYLIN</b> (UNII: KYR8BR2X6O)	
<b>ETHYL ACETATE</b> (UNII: 76845O8NMZ)	
<b>ACETONE</b> (UNII: 1364PS73AF)	
<b>AMYL ACETATE</b> (UNII: 92Q24NH7AS)	
<b>CASTOR OIL</b> (UNII: D5340Y2I9G)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0779-3	30 mL in 1 BOTTLE, WITH APPLICATOR; Type 0: Not a Combination Product	02/01/2020	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	02/01/2020	

**Labeler** - Rite Aid Corporation (014578892)

Revised: 2/2024

Rite Aid Corporation