

**NICOTINE TRANSDERMAL- nicotine**  
**Rite Aid Corporation**

-----  
**Drug Facts**

***Active ingredient Step 1 (in each patch)***

Nicotine, 21 mg delivered over 24 hours

***Active ingredient Step 2 (in each patch)***

Nicotine, 14 mg delivered over 24 hours

***Active ingredient Step 3 (in each patch)***

Nicotine, 7 mg delivered over 24 hours

***Purpose***

Stop smoking aid

***Use***

reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

***Warnings***

**If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.** Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Do not use**

- if you continue to smoke, chew tobacco, use snuff, use nicotine gum, or use another nicotine patch or other nicotine containing products

**Ask a doctor before use if you have**

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to

get rashes.

### **Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

### **When using this product**

- do not smoke even when not wearing the patch. The nicotine in your skin will still be entering your blood stream for several hours after you take off the patch.
- if you have vivid dreams or other sleep disturbances remove this patch at bedtime

### **Stop use and ask a doctor if**

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat

### **Keep out of reach of children and pets.**

Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

### **Directions**

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed self-help guide for complete directions and other information
- stop smoking completely when you begin using the patch
- **if you smoke more than 10 cigarettes per day**, use the following schedule below:

STEP 1	STEP 2	STEP 3
Use one 21mg patch/day	Use one 14mg patch/day	Use one 7mg patch/day
Weeks 1-4	Weeks 5-6	Weeks 7-8

- if you smoke **10 or less cigarettes per day**, start with **Step 2** for 6 weeks, then **Step 3** for 2 weeks and then stop
- apply one new patch every 24 hours on skin that is dry, clean and hairless
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

- the used patch should be removed and a new one applied to a different skin site at the same time each day
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- to avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures
- stop using the patch at the end of 8 weeks. If you still feel the need to use the patch talk to your doctor.

### ***Other information***

store at 20-25°C (68-77°F)

### ***Inactive ingredients***

Acrylate adhesive, aluminized polyester, cellulose paper, methacrylic acid copolymer

### ***Questions or comments?***

call **1-800-585-8682**

Weekdays (9am-8pm ET) or visit us at [www.habitrol.com](http://www.habitrol.com)

### **TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. Use one patch daily according to directions.
3. It is important to complete treatment.
4. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.
5. Use patch with a behavioural support program, such as the one described in the enclosed booklet.

For your family's protection, patches are supplied in child-resistant pouches. **Do not use if individual pouch is open or torn.**

- **Not for sale to persons under 18 years of age.**
- **Proof of age required.**
- **Not for sale in vending machines or from any source where proof of age cannot be verified.**

### **Principal Display Panel Kit**

**Nicotine Transdermal System**PATCH Kit



# NICOTINE TRANSDERMAL

nicotine kit

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-3627(NDC:43598-445)
---------------------	----------------	---------------------------	-------------------------------

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-3627-3	56 in 1 CARTON	03/31/2021	
1		1 in 1 PATCH; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

## Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1		1
Part 2		1
Part 3		1

Part 1 of 3

NICOTINE

nicotine transdermal patch, extended release

### Product Information

**Route of Administration** TRANSDERMAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Nicotine</b> (UNII: 6M3C89ZY6R) (Nicotine - UNII:6M3C89ZY6R)	Nicotine	21 mg in 24 h

### Inactive Ingredients

Ingredient Name	Strength
<b>Methacrylic Acid</b> (UNII: 1CS02G8656)	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020076	01/30/2020	

## Part 2 of 3

### NICOTINE

nicotine transdermal patch, extended release

### Product Information

**Route of Administration** TRANSDERMAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Nicotine</b> (UNII: 6M3C89ZY6R) (Nicotine - UNII:6M3C89ZY6R)	Nicotine	14 mg in 24 h

### Inactive Ingredients

Ingredient Name	Strength
<b>Methacrylic Acid</b> (UNII: 1CS02G8656)	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020076	01/30/2020	

### Part 3 of 3

### NICOTINE

nicotine transdermal patch, extended release

### Product Information

**Route of Administration** TRANSDERMAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Nicotine</b> (UNII: 6M3C89ZY6R) (Nicotine - UNII:6M3C89ZY6R)	Nicotine	7 mg in 24 h

### Inactive Ingredients

Ingredient Name	Strength
<b>Methacrylic Acid</b> (UNII: 1CS02G8656)	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020076	01/30/2020	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020076	03/31/2021	

**Labeler** - Rite Aid Corporation (014578892)