

**CORICIDIN HBP MAXIMUM STRENGTH MULTI SYMPTOM FLU- acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide tablet  
Bayer HealthCare LLC.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Coricidin HBP Max Strength Multi-symptom Flu (tablet) new product name  
Perrigo Old name Coricidin HBP Max Strength (tablet)**

**Coricidin HBP**

**Maximum Strength Multi-Symptom Flu**

***Drug Facts***

**Active Ingredient**

***Active ingredients (in each tablet) Purposes***

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Chlorpheniramine maleate 2 mg.....Antihistamine  
Dextromethorphan hydrobromide 10 mg.....Cough suppressant

**Uses**

***Uses***

- temporarily relieves these symptoms due to a cold or flu:
- minor aches and pains
- headache
- cough
- runny nose
- sneezing
- sore throat
- temporarily reduces fever

**Warnings**

**Liver warning**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert**

**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

### **Sore throat warning**

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with excessive phlegm (mucus)
- a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** Abuse of this product can lead to serious injury.

### **Overdose warning**

**Overdose warning:** Taking more than the recommended dose may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

### **Directions**

do not use more than directed (see overdose warning)

adults and children 12 years and over: take 2 tablets every 4 hours, while symptoms persist. Do not take more than 10 tablets in 24 hours or as directed by a doctor.

children under 12 years of age: ask a doctor


### **Other information**

store at room temperature. Avoid excessive heat.

**Inactive ingredients** FD&C red #40 aluminum lake, lecithin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

**Questions or comments?** 1-800-317-2165 (Mon-Fri 9AM - 5PM EST)

**Carton**



**Coricidin®**  
**HBP COLD RELIEF FOR PEOPLE WITH**  
**HIGH BLOOD PRESSURE†**  
**MAXIMUM STRENGTH**  
**MULTI-SYMPTOM**  
**FLU**

Relieves

- Body Aches & Pains
- Headache
- Fever
- Cough
- Runny Nose & Sneezing
- Sore Throat

†Decongestant Free

24 tablets

**Acetaminophen - Pain Reliever/Fever Reducer**  
**Chlorpheniramine Maleate - Antihistamine**

# Dextromethorphan HBr - Cough Suppressant

LEARN MORE

American Heart Association

Information Enclosed

## CORICIDIN HBP MAXIMUM STRENGTH MULTI SYMPTOM FLU

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11523-0038
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

### Product Characteristics

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	c;flu
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11523-0038-1	1 in 1 CARTON	09/04/2020	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination		

1		Product		
2	NDC:11523-0038-2	2 in 1 CARTON	09/04/2020	
2		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/01/2020	

**Labeler** - Bayer HealthCare LLC. (112117283)

Revised: 3/2023

Bayer HealthCare LLC.