

**PENTREXCILINA- acetaminophen, phenylephrine hydrochloride,  
dextromethorphan hydrobromide tablet  
Healthlife of USA**

-----  
**PENTREXCILINA**

**Drug Facts**

Active Ingredients (in each tablet)

Acetaminophen USP 325mg.....Pain reliever/fever  
reducer

Guaifenesin USP 200mg.....Expectorant

Phenylephrine HCL USP 5mg.....Decongestant

Dextromethorphan HBr USP 15mg.....Antitussive

**Purpose**

Pain reliever/ fever reducer

Expectorant

Decongestant

Antitussive

**Uses**

Temporarily

- Relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- Relieves sinus congestion and pressure, helps decongest sinus openings and passages
- Restores freer breathing
- Helps loosen bothersome mucus, drain bronchial tubes, and make coughs more productive
- Suppresses cough due to minor throat and bronchial irritation associated with a cold or inhaled irritants
- Temporarily relieves minor aches, pains and fever associated with: headache, common cold, toothache, backache, muscular aches, menstrual cramps

**Warnings: Liver Warning:**

This product contains acetaminophen. Severe liver damage may occur if you take:

- More than 8 tablets in 24 hours
- With other drugs containing acetaminophen (prescription or nonprescription). Ask a doctor or pharmacist before using with other drugs if you are not sure

- 3 or more alcoholic drinks every day while using this product

**Allergy alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash if a skin reaction occurs, stop use and seek medical help right away.

**Warning:** A persistent cough may be a sign of a serious condition

**Do not**

- use with any other product containing acetaminophen this will provide more than the recommended dose (overdose) of acetaminophen and could cause serious health concerns.
- use more than the recommended dose
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor
- when using this product do not exceed recommended dose
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping MAOI drug. If you do not know if your prescription drug contains an MAOI, consult a doctor or pharmacist before taking this product
- take this product for persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema or where cough is accompanied by excessive phlegm (mucus) unless directed by a doctor.

**Stop use and ask a doctor if:**

symptoms do not improve, pain or fever persists or gets worse, new symptoms occur, redness or swelling is present, symptoms do not improve within 7 days or are accompanied by fever, cough persists for more than 1 week, tends to recur, or is accompanied by a fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition

**Ask a doctor before use if you have:**

heart disease, high blood pressure, thyroid disease, diabetes, difficulty in urination due to enlargement of the prostate gland, liver disease

**Ask a doctor or pharmacist if you are**

taking the blood thinning drug warfarin.

**Do not exceed recommended dosage**

if nervousness, dizziness, or sleeplessness occur, discontinue use consult a doctor.

**If pregnant or breast-feeding,**

ask a health professional before use.

**KEEP OUT OF REACH OF CHILDREN**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms

**Directions:**

Adults and children 12 years of age and older: Take 2 tablets every 6 to 8 hours as needed, do not exceed 8 tablets in 24 hours, or as directed by a doctor. Children under 12 years, consult a doctor

**Other Information:**

- Tamper evident. Do not use if packet is torn, cut or opened
- Store at controlled room temperature 15° to 30°C (59° to 86°F)
- Avoid excessive heat and humidity

**Inactive Ingredients**

maltodextrin, microcrystalline cellulose, povidone, silicon dioxide\*, sodium starch glycolate, starch, stearic acid

(\*may contain)

**Questions or comments?**

Call: 619-600-5632 (Mon-Fri 9am - 5pm EST)

**Pentrexilina****2 Tablets in a pouch**

**2 TABLETS**



Acetaminophen / Dextromethorphan HBr  
Guaifenesin / Phenylephrine HCl

**PENTREXILINA.**

**Directions:** Adults and children 12 years and over:  
Take 2 tablets every 6 to 8 hours as needed.  
Do not take more than 8 tablets in 24 hours.  
Children under 12 years of age: ask a doctor.

## Cold, Cough & Flu Relief

Acetaminophen 325mg  
Pain reliever / Fever Reducer  
Dextromethorphan HBr 15mg  
Cough suppressant  
Guaifenesin 200mg  
Expectorant  
Phenylephrine HCl 5mg  
Nasal Decongestant

**BEFORE USE READ ALL WARNINGS  
AND DIRECTIONS IN PRODUCT CARTON  
YOU WILL FIND COMPLETE INFORMATION**

### DRUG FACTS

ACTIVE INGREDIENTS	PURPOSE
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestion

### Uses: temporarily

• relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies • relieves sinus congestion and pressure • helps decongest sinus openings and passages • restores free breathing • helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passages of bothersome mucus, drain bronchial tubes, and make coughs more productive. • suppresses cough due to minor throat and bronchial irritation associated with a cold or inhaled irritants

temporarily relieves minor aches, pains, and fever associated with headache, common cold, toothache, backache, muscular aches, menstrual cramps

### Warnings:

**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if you take • more than 8 tablets in 24 hours • with other drugs containing acetaminophen (prescription or nonprescription). Ask a doctor or pharmacist before using with other drugs if you are not sure. • 3 or more alcoholic drinks every day while using this product.

Manufactured for:  
PRX San Diego, CA  
Exclusively distributed by:  
OPMX San Diego, CA 92154

**DO NOT USE IF  
POUCH IS BROKEN  
OR TORN**

**COLD, FLU & COUGH**  
**PENTREXCILINA®**

**Drug Facts** (in each tablet)  
**ACTIVE INGREDIENTS**  
 Acetaminophen 325 mg  
 Dextromethorphan HBr 15 mg  
 Guaifenesin 200 mg  
 Phenylephrine HCl 5 mg  
 Pseudoephedrine HCl 15 mg  
**Uses:** temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies ■ relieves sinus congestion and pressure ■ helps decompress sinus openings and passages ■ restores nasal breathing ■ helps loosen phlegm (mucus) and suppresses cough due to minor throat and bronchial irritation associated with a cold or influenza ■ temporarily relieves minor aches, pains, and fever associated with headache, common cold, toothache, backache, muscular aches, menstrual cramps  
**Warnings:** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 8 tablets in 24 hours ■ with other drugs containing acetaminophen (prescription or nonprescription). Ask a doctor or pharmacist before using with other drugs if you are not sure. ■ 3 or more alcoholic drinks every day while using this product  
**Caution:** acetaminophen may cause severe skin reactions. Symptoms may include skin redness, blisters, rash if a skin reaction occurs. Stop use and seek medical help right away.  
**Warning:** A persistent cough may be a sign of a serious condition.  
**Do not:**  
 ■ use with any other product containing acetaminophen as this will provide more than the recommended dose (avoidance of acetaminophen and other products containing acetaminophen) ■ use more than the recommended dose ■ for more than 10 days for pain unless directed by a doctor ■ for more than 5 days for fever unless directed by a doctor ■ when using this product do not exceed recommended dose. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or 2 weeks after stopping MAOI drug. ■ take this product for persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema or where cough is accompanied by excessive phlegm (mucus) unless directed by a doctor.  
**Stop use and ask a doctor if:** symptoms do not improve within 7 days or are accompanied by fever, cough persists for more than 1 week, tends to recur, or is accompanied by a fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.  
**Ask a doctor before use if you have:** heart disease, high blood pressure, thyroid disease, diabetes, difficulty in urination due to enlargement of the prostate gland, liver disease.  
**Ask a doctor or pharmacist if you are taking the blood thinning drug warfarin.**  
**Do not exceed recommended dosage.** If nervousness, dizziness, or sleepiness occur, discontinue use and consult a doctor.  
**If pregnant or breast-feeding,** ask a health professional before use.  
**KEEP OUT OF REACH OF CHILDREN.** In case of overdose, get medical help or contact a Poison Control Center right away.  
 Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**PENTREXCILINA®**

Exclusively distributed by:  
  
 San Diego, CA 92154  
 Phone: 619-600-5632

Made for:  
  
 San Diego, CA



**COLD, FLU & COUGH EXTRA STRENGTH**

NDC 69517-155-06

Acetaminophen 325 mg / Dextromethorphan HBr 15 mg  
 Guaifenesin 200 mg / Phenylephrine HCl 5 mg


**PENTREXCILINA®**

- Cough Suppressant
- Expectoant
- Nasal Congestion
- Sneezing
- Itchy, Watery Eyes & Runny Nose

**12 Tablets**

**Drug Facts (Continued)**  
**Directions:**  
 Adults and children 12 years of age and older: Take 2 tablets every 6 to 8 hours as needed, do not exceed 8 tablets in 24 hours, or as directed by a doctor.  
 Children under 12 years, consult a doctor  
**Other Information:** ■ Tamper Evident. Do not use if packet is torn, cut or opened  
 ■ Store at controlled room temperature 15° to 30° C (59° to 86° F) ■ Avoid excessive heat and humidity  
**Inactive Ingredients:** mallo-dextrin, microcrystalline cellulose, povidone, silicon dioxide\*, sodium starch glycolate, starch, stearic acid (\*May contain)  
 Questions or comments? Call: 619-600-5632 (Mon-Fri 9am-5pm EST)


Code No.: GUJDRUGS/G125/2025  
 Batch #  
 Exp. date



0 39853 98970 5

75 mm  
 2.952 in

30 r  
 1.18



## PENTREXCILINA

acetaminophen, phenylephrine hydrochloride, dextromethorphan hydrobromide tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69517-155
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>SODIUM STARCH GLYCOLATE TYPE A CORN</b> (UNII: AG9B65PV6B)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	CAPSULE	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	A15
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69517-155-06	6 in 1 PACKAGE	01/05/2018	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:69517-155-72	72 in 1 PACKAGE	01/05/2018	
2		2 in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/05/2018	

**Labeler** - Healthlife of USA (079656178)

## Establishment

Name	Address	ID/FEI	Business Operations
Vovantis Laboratories Private Limited		650502151	manufacture(69517-155)

Revised: 10/2023

Healthlife of USA