

**ASPIRIN PAIN RELIEF- aspirin tablet, delayed release**  
**Rite Aid Corporation**

-----  
**Rite Aid 44-227**

***Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

***Uses***

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - headache
  - menstrual pain
  - a cold
  - toothache
  - muscle pain
- or as recommended by your doctor

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- hives
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- new symptoms occur
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

**Inactive ingredients**

black iron oxide, cellulose, colloidal anhydrous silica, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, propylene glycol, shellac wax, simethicone, sodium alginate, sodium bicarbonate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

**Questions or comments?**

**1-800-426-9391**

**Principal Display Panel**

NDC 11822-2270-4

\*\*Compare to the active ingredient  
in **Ecotrin®** Regular Strength

**Talk to your doctor or other healthcare provider  
before using this product for your heart.**

REGULAR STRENGTH  
PAIN RELIEF  
**ASPIRIN**

**ASPIRIN 325 mg  
PAIN RELIEVER (NSAID)**

aspirin regimen  
safety coated

**500** ENTERIC  
COATED TABLETS

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

\*\*This product is not manufactured or distributed by Medtech Products Inc.,  
owner of the registered trademark Ecotrin® Regular Strength. 50844  
REV0122C22714

**DISTRIBUTED BY: RITE AID  
30 HUNTER LANE  
CAMP HILL, PA 17011  
www.riteaid.com**

**SATISFACTION GUARANTEE**

**If you're not satisfied, we'll happily refund your money.**

**Drug Facts (continued)**

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

\*\*This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Ecotrin® Regular Strength. 50844 REV0122C22714

**DISTRIBUTED BY: RITE AID**  
30 HUNTER LANE,  
CAMP HILL, PA 17011  
www.riteaid.com

**SATISFACTION GUARANTEE**  
If you're not satisfied, we'll happily refund your money.

0 11822 42300 7

**PEEL HERE FOR MORE DRUG FACTS**

NDC 11822-2270-4  
Talk to your doctor or other healthcare provider before using this product for your heart.

\*\*Compare to the active ingredient in Ecotrin® Regular Strength

**REGULAR STRENGTH PAIN RELIEF**

**ASPIRIN**

**ASPIRIN 325 mg**

**PAIN RELIEVER (NSAID)**

aspirin regimen  
safety coated

**500 ENTERIC COATED TABLETS**

**No Print/No Varnish Area**  
Lot no. & Exp. Date

**Drug Facts**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

Active ingredient (in each tablet)	Purpose
Aspirin 325 mg (NSAID)	Pain reliever
*nonsteroidal anti-inflammatory drug	

**Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis    ■ headache    ■ menstrual pain    ■ a cold
- toothache    ■ muscle pain
- or as recommended by your doctor

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- facial swelling    ■ hives    ■ shock
- asthma (wheezing)

**STOP PEELING**

**Drug Facts (continued)**

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are taking a prescription drug for**

- gout    ■ diabetes    ■ arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child

**Drug Facts (continued)**

or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

**Inactive ingredients** black iron oxide, cellulose, colloidal anhydrous silica, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, propylene glycol, shellac wax, simethicone, sodium alginate, sodium bicarbonate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

**Questions or comments?**  
1-800-426-9391

**Rite Aid 44-227**

## ASPIRIN PAIN RELIEF

aspirin tablet, delayed release

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-2270
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>FERROSO FERRIC OXIDE</b> (UNII: XM0M87F357)	

<b>POWDERED CELLULOSE</b> (UNII: SMD1X3XO9M)
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)
<b>D&amp;C YELLOW NO. 10 ALUMINUM LAKE</b> (UNII: CQ3XH3DET6)
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)
<b>Polyvinyl Acetate Phthalate</b> (UNII: 58QVG85GW3)
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)
<b>SHELLAC</b> (UNII: 46N107B71O)
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)
<b>WATER</b> (UNII: 059QF0KO0R)
<b>SODIUM ALGINATE</b> (UNII: C269C4G2ZQ)
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)
<b>TALC</b> (UNII: 7SEV7J4R1U)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)
<b>TRIACETIN</b> (UNII: XHX3C3X673)
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;227
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-2270-4	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/14/1992	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/14/1992	

**Labeler** - Rite Aid Corporation (014578892)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	manufacture(11822-2270) , pack(11822-2270)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		832867837	manufacture(11822-2270)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		832867894	manufacture(11822-2270)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		868734088	manufacture(11822-2270)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		967626305	pack(11822-2270)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		117025878	manufacture(11822-2270)

Revised: 12/2023

Rite Aid Corporation