


**SORBITOL- sorbitol irrigant**  
**Baxter Healthcare Corporation**

-----  
**3% Sorbitol Urologic Irrigating Solution**  
**in UROMATIC Plastic Container**

**DESCRIPTION**

3% Sorbitol Urologic Irrigating Solution is a sterile, nonpyrogenic, nonhemolytic, electrically nonconductive solution in a single dose UROMATIC Plastic Container for use as a urologic irrigating solution. Each liter contains 30 g Sorbitol in Water for Injection. pH 5.0 (4.5 to 6.5). Osmolarity 165 mOsmol/L (calc.). No antimicrobial agent has been added.

Sorbitol is a reduced form of dextrose and is designated chemically as D-Glucitol,



The UROMATIC plastic container is fabricated from a specially formulated polyvinyl chloride (PL 146 plastic). The amount of water that can permeate from inside the container into the overwrap is insufficient to affect the solution significantly. Solutions in contact with the plastic container can leach out certain of its chemical components in very small amounts within the expiration period, e.g., di-2-ethylhexyl phthalate (DEHP), up to 5 parts per million; however, the safety of the plastic has been confirmed in tests in animals according to USP biological tests for plastic containers as well as by tissue culture toxicity studies.

**CLINICAL PHARMACOLOGY**

3% Sorbitol Urologic Irrigating Solution is useful as an irrigating fluid for the urinary bladder because the sorbitol solution is nonhemolytic, electrically nonconductive, and provides a high degree of visibility for urologic procedures requiring endoscopy. During transurethral surgical procedures, the solution acts as a lavage for removing blood and tissue fragments. It is also useful as an irrigating fluid to maintain the patency of an indwelling catheter in the immediate postoperative period. If absorbed either intravascularly or extravascularly during transurethral resections, the sorbitol will be either metabolized to carbon dioxide and water via the fructose pathway, or excreted by a normally functioning kidney.

**INDICATIONS AND USAGE**

3% Sorbitol Urologic Irrigating Solution is indicated for use as a urologic irrigating fluid with endoscopic instruments during transurethral procedures requiring distension, irrigation, and lavage of the urinary bladder. It may be used for lavage of an indwelling catheter to maintain patency.

**CONTRAINDICATIONS**

Anuria.

**WARNINGS**

Not for injection.

**Not for use in patients with documented or suspected fructose intolerance.**

Solutions for urologic irrigation must be used with caution in patients with severe cardiopulmonary or renal dysfunction.

Irrigating fluids used during transurethral prostatectomy have been demonstrated to enter the systemic circulation in relatively large volumes; thus sorbitol irrigating solution must be regarded as a systemic drug. Absorption of large amounts of fluids containing sorbitol may significantly alter cardiopulmonary and renal dynamics. **Appropriate patient monitoring should be maintained due to the possibility of fluid overload. Should fluid overload occur, intensive fluid and electrolyte management is necessary. Monitoring of fluid and electrolyte levels beyond the acute phase may be considered due to the possibility of delayed fluid absorption.** (See ADVERSE REACTIONS, Post-Marketing Experience)

Hyperglycemia from metabolism of absorbed sorbitol may occur in patients with diabetes mellitus.

The contents of an opened container should be used promptly to minimize the possibility of bacterial growth or pyrogen formation. Discard the unused portion of irrigating solution since no antimicrobial agent has been added.

## **PRECAUTIONS**

The cardiovascular status, especially of the patient with cardiac disease, should be carefully observed before and during transurethral resection of the prostate when using 3% Sorbitol Urologic Irrigating Solution, because the quantity of fluid absorbed into the systemic circulation by opened prostatic veins may produce significant expansion of the intravascular fluid and lead to fulminating congestive heart failure.

Shift of sodium free intracellular fluid into the extracellular compartment, following systemic absorption may lower serum concentration and aggravate preexisting hyponatremia.

Excessive loss of water and electrolytes may lead to serious imbalances. With continuous administration of solution, loss of water may occur in excess of electrolytes, producing hypernatremia.

Sustained diuresis that results from transurethral irrigation with sorbitol irrigating solutions may obscure and intensify inadequate hydration or hypovolemia.

### **Pregnancy**

#### Teratogenic Effects

Animal reproduction studies have not been conducted with 3% Sorbitol Urologic Irrigating Solution. It is not known whether 3% Sorbitol Urologic Irrigating Solution can cause fetal harm when administered to a pregnant woman or can affect reproductive capacity. 3% Sorbitol Urologic Irrigating Solution should be given to a pregnant woman only if clearly needed.

### **Pediatric Use**

Safety and effectiveness in pediatric patients have not been established.

### **Geriatric Use**

Clinical studies of Irrigation Solutions did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from other younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients.

Do not administer unless the solution is clear and the seal is intact.

## **ADVERSE REACTIONS**

**Life threatening** adverse reactions with intravenous sorbitol infusions have been reported in patients with fructose intolerance.

The literature reports occasional adverse reactions for intravenous sorbitol infusions. These include disturbances such as acidosis, electrolyte loss, marked diuresis, urinary retention, edema, dryness of mouth and thirst, and dehydration; cardiovascular/pulmonary disorders such as pulmonary congestion, hypotension, tachycardia, angina-like pains, and other general reactions such as blurred vision, convulsions, nausea, vomiting, diarrhea, rhinitis, chills, vertigo, and backache. Allergic reactions reported to occur from sorbitol include urticaria.

Should adverse reactions occur, discontinue the irrigant and reevaluate the clinical status of the patient.

## **Post-Marketing Experience**

Following off-label use of 3% Sorbitol Urologic Irrigating Solution for hysteroscopic procedures in women, life-threatening adverse events related to fluid overload have been reported.

## **DOSAGE AND ADMINISTRATION**

The volume of solution needed will vary with the nature and duration of the urologic procedure.

If desired, warm in overwrap to near body temperature in a water bath or oven heated to not more than 45°C.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit.

## **HOW SUPPLIED**

3% Sorbitol Urologic Irrigating Solution in UROMATIC plastic containers is available as follows:

2B7357	3000mL	NDC 0338-0295-47
--------	--------	------------------

Exposure of pharmaceutical products to heat should be minimized. Avoid excessive heat. It is recommended the product be stored at room temperature (25°C); brief exposure up to 40°C does not adversely affect the product.

## **DIRECTIONS FOR USE**

Tear overwrap down side at slit and remove solution container. Visually inspect the container. If the outlet port protector is damaged, detached, or not present, discard container as solution path sterility may be impaired. Some opacity of the plastic due to moisture absorption during the sterilization process may be observed. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually. Check for minute leaks by squeezing bag firmly. If leaks are found, discard solution as sterility may be impaired.

Use Aseptic Technique.

1. Suspend container using hanger hole.
2. Remove protector from outlet port.
3. Attach irrigation set. Refer to complete directions accompanying set.

**Baxter Healthcare Corporation**  
Deerfield, IL 60015 USA

Printed in USA

Baxter and Uromatic are trademarks of Baxter International Inc.

07-19-00-0227

Rev. January 2018

## PRINCIPAL DISPLAY PANEL - PACKAGING LABELING

LOT	EXP		
⊗	⊗	<b>3000 mL</b>	<b><u>2700</u></b>
		<b>2B7357</b>	
NOT FOR INJECTION	NDC 0338-0295-47		
<b>3%</b> <b>Sorbitol Urologic</b> <b>Irrigating Solution</b>			<b><u>2400</u></b>
EACH 100 mL CONTAINS 3 g SORBITOL NO ANTIMICROBIAL AGENT HAS BEEN ADDED pH 5.0 (4.5 TO 6.5) HYPOTONIC OSMOLARITY 165 mOsmol/L (CALC) STERILE NONPYROGENIC SINGLE DOSE CONTAINER			<b><u>1500</u></b>
DO NOT USE UNLESS SOLUTION IS CLEAR DISCARD UNUSED PORTION HYPOTONIC SOLUTION FOR BLADDER IRRIGATION IN CONNECTION WITH CYSTOSCOPIC EXAMINATION AND UROLOGICAL SURGICAL PROCEDURES SEE DIRECTION SHEET			<b><u>1200</u></b>
CAUTIONS SQUEEZE AND INSPECT INNER BAG WHICH MAINTAINS PRODUCT STERILITY DISCARD IF LEAKS ARE FOUND Rx ONLY STORE UNIT IN MOISTURE BARRIER OVERWRAP AT ROOM TEMPERATURE (25°C) UNTIL READY TO USE AVOID EXCESSIVE HEAT SEE INSERT			<b><u>900</u></b>
UROMATIC CONTAINER PL 146 PLASTIC			
<b>Baxter</b> BAXTER HEALTHCARE CORPORATION DEERFIELD IL 60015 USA MADE IN USA	BAXTER UROMATIC AND PL 146 ARE TRADEMARKS OF BAXTER INTERNATIONAL INC	FOR PRODUCT INFORMATION 1-800-933-0300	<b><u>600</u></b>
			<b><u>300</u></b>

**Container Label**

Container Label

LOT

EXP

**3000 mL**

**2B7357**

**NDC 0338-0295-47**

**NOT FOR INJECTION**

**3%**  
**Sorbitol Urologic**  
**Irrigating Solution**

EACH 100 mL CONTAINS 3 g SORBITOL  
NO ANTIMICROBIAL AGENT HAS BEEN ADDED  
pH 5.0 (4.5 TO 6.5) HYPOTONIC  
OSMOLARITY 165 mOsmol/L (CALC)  
STERILE NONPYROGENIC SINGLE DOSE  
CONTAINER

DO NOT USE UNLESS SOLUTION IS CLEAR  
DISCARD UNUSED PORTION HYPOTONIC  
SOLUTION FOR BLADDER IRRIGATION IN  
CONNECTION WITH CYSTOSCOPIC EXAMINATION  
AND UROLOGICAL SURGICAL PROCEDURES SEE  
DIRECTION SHEET

CAUTIONS SQUEEZE AND INSPECT INNER BAG  
WHICH MAINTAINS PRODUCT STERILITY  
DISCARD IF LEAKS ARE FOUND RX ONLY  
STORE UNIT IN MOISTURE BARRIER OVERWRAP  
AT ROOM TEMPERATURE (25°C) UNTIL READY TO  
USE AVOID EXCESSIVE HEAT SEE INSERT

UROMATIC CONTAINER

PL 146 PLASTIC

BAXTER UROMATIC AND PL 146 ARE  
TRADEMARKS OF BAXTER INTERNATIONAL INC

***Baxter Logo***  
**BAXTER HEALTHCARE CORPORATION**  
DEERFIELD IL 60015 USA

MADE IN USA

FOR PRODUCT INFORMATION  
1-800-933-0303

2B7357

4-3000 ML  
UROMATIC CONTAINERS

3% SORBITOL UROLOGIC IRRIGATING SOLUTION

EXP  
XXXXX

SECONDARY BAR CODE

(17) YYMM00 (10) XXXXX

LOT  
XXXXX

PRIMARY BAR CODE

(01) 50303380295475

**Carton Label**

Carton Label

2B7357

4-3000 ML  
UROMATIC CONTAINERS

3% SORBITOL UROLOGIC IRRIGATING SOLUTION

EXP  
XXXXX

SECONDARY BAR CODE  
(17) YYMM00 (10) XXXXX

LOT  
XXXXX

PRIMARY BAR CODE  
(01) 50303380295475

**SORBITOL**

sorbitol irrigant

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0338-0295
<b>Route of Administration</b>	IRRIGATION		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
SORBITOL (UNII: 506T60A25R) (SORBITOL - UNII:506T60A25R)	SORBITOL	30.00 g in 1000.00 mL

**Inactive Ingredients**

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0295-47	3000 mL in 1 BAG; Type 0: Not a Combination Product	05/30/1980	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA017863	05/30/1980	

**Labeler** - Baxter Healthcare Corporation (005083209)**Establishment**

Name	Address	ID/FEI	Business Operations
Baxter Healthcare Corporation		059140764	MANUFACTURE(0338-0295) , ANALYSIS(0338-0295) , PACK(0338-0295) , LABEL(0338-0295) , STERILIZE(0338-0295)

**Establishment**

Name	Address	ID/FEI	Business Operations
Baxter Healthcare Corporation		194684502	ANALYSIS(0338-0295)

Revised: 12/2019

Baxter Healthcare Corporation