

# **GUNA-ANTI IL 1- anti-interleukin-1.alpha. immunoglobulin g rabbit-canakinumab solution/ drops**

**Guna spa**

*Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.*

-----

## **GUNA-ANTI IL1**

### **ACTIVE INGREDIENTS/PURPOSE**

ANTI INTERLEUKIN 1 ALPHA	4C	ACUTE PAIN AND FEVER RELIEF
ANTI INTERLEUKIN 1 BETA	4C	ACUTE PAIN AND FEVER RELIEF

### **USES**

Temporary relief of:

- **Acute pain**
- **Fever**

### **WARNINGS**

- **Stop use and ask doctor if symptoms persist** more than 5 days.
- **If pregnant or breast-feeding** ask a health professional before use.
- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.
- Contains ethyl alcohol 30%

### **PREGNANCY**

**If pregnant or breast-feeding** ask a doctor before use.

### **WARNINGS**

**Keepout of reach of children.**

### **DIRECTIONS**

**Adults and children 12 years and older** 20 drops twice a day in a little water. Hold in the mouth for about 30 seconds then swallow.

**Children between 12 years and 6 years of age** 10 drops twice a day in a little water. Hold in the mouth for about 30 seconds then swallow.

**Children under 6 years** 5 drops twice a day in a glass of water.

## QUESTIONS

**Questions?:** info@gunainc.com  
Tel. (484) 223-3500

Take 15 minutes before meals.

**Inactive ingredient:** Ethyl alcohol 30%.

## PRINCIPAL DISPLAY PANEL

NDC 17089-299-18

**Uses:** Temporary relief of: acute pain, fever.

These statements have not been evaluated by the FDA. They are based upon homeopathic principles.

**Drug Facts**

Active ingredients	Purpose
30 ml contains:	
Anti interleukin 1 alpha 4C 15 ml	Acute pain
Anti interleukin 1 beta 4C 15 ml	and fever relief

**Inactive ingredient:** Ethyl alcohol 30%.

**Warnings:**

- Stop use and ask doctor if symptoms persist more than 5 days.
- If pregnant or breast-feeding ask a health professional before use.
- Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Drug Facts (continued)**

**Directions:**  
Take 15 minutes before meals.

Adults and children 12 years and older	20 drops in a little water, 2 times per day
Children between 12 years and 6 years of age	10 drops in a little water, 2 times per day
Children under 6 years	5 drops in a glass of water, 2 times per day

**Other information:**


- Store at 20°-25°C (68°-77°F).
- Excursions permitted between 15°-30°C (59°-86°F).

**Questions?:** info@gunainc.com  
tel. (484) 223-3500

**Do not use if tamper-evident seal is broken.**

Manufactured by  
GUNA S.p.a. - Italy

**GUNA**  
biotherapeutics  
GUNA Inc.



## GUNA-ANTI IL 1

anti-interleukin-1.alpha. immunoglobulin g rabbit- canakinumab solution/ drops

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:17089-398
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ANTI-INTERLEUKIN-1.ALPHA. IMMUNOGLOBULIN G RABBIT</b> (UNII: ML4QRZ1HCL) (ANTI-INTERLEUKIN-1.ALPHA. IMMUNOGLOBULIN G RABBIT - UNII:ML4QRZ1HCL)	ANTI-INTERLEUKIN-1.ALPHA. IMMUNOGLOBULIN G RABBIT	4 [hp_C] in 30 mL
<b>CANAKINUMAB</b> (UNII: 37CQ2C7X93) (CANAKINUMAB - UNII:37CQ2C7X93)	CANAKINUMAB	4 [hp_C] in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ALCOHOL</b> (UNII: 3K9958V90M)	9 mL in 30 mL

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17089-398-18	1 in 1 BOX	12/21/2018	
1		30 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved homeopathic		06/17/2008	

**Labeler** - Guna spa (430538264)

## Establishment

Name	Address	ID/FEI	Business Operations
Guna spa		338587646	manufacture(17089-398)

Revised: 11/2022

Guna spa