

**NAPROXEN SODIUM- naproxen sodium tablet, film coated**  
**Rite Aid Corporation**

-----  
**Rite Aid Corporation Naproxen Sodium Tablets, 220 mg Drug Facts**

**Active ingredient (in each caplet)**

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

**Warnings**

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - you have difficulty swallowing
  - it feels like the pill is stuck in your throat
  - redness or swelling is present in the painful area
  - any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older:

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period

Children under 12 years:

- ask a doctor

**Other information**

- each caplet contains: sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

**Inactive ingredients**

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

**Questions or comments?**

1-800-719-9260

**Principal Display Panel**

Compare to the active ingredient of Aleve® Caplets

pain reliever/fever reducer

naproxen sodium tablets, 220 mg

pain reliever/fever reducer (NSAID)

strength to last 12 hours

actual size

50 CAPLETS†

†capsule-shaped tablets

**KEEP CARTON  
FOR REFERENCE**

**DO NOT USE IF PRINTED  
FOIL UNDER CAP IS  
BROKEN OR MISSING**

ACTUAL SIZE



**\*\*Compare to the active ingredient of Aleve® Caplets**



**pain reliever/fever reducer  
naproxen sodium**

**tablets, 220 mg**



**pain reliever/fever reducer  
(NSAID)**


**50 CAPLETS†**  
†capsule-shaped tablets

**strength to  
last 12 hours**



actual size





EXP. DATE :

EXP.

LOT NO.



RA# 0398521

### Drug Facts

**Active ingredient (in each caplet)**      **Purposes**  
Naproxen sodium 220 mg      Pain reliever/  
(naproxen 200 mg) (NSAID)\* .....fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ temporarily relieves minor aches and pains due to:  
■ min or pain of arthritis    ■ muscular aches    ■ backache  
■ menstrual cramps        ■ headache        ■ toothache  
■ the common cold        ■ temporarily reduces fever

### Warnings

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

■ hives    ■ facial swelling    ■ asthma (wheezing)  
■ shock   ■ skin reddening   ■ rash   ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you: ■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if** ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have asthma

**Ask a doctor or pharmacist before use if you are**  
■ under a doctor's care for any serious condition  
■ taking any other drug

### Drug Facts (continued)

#### When using this product

■ take with food or milk if stomach upset occurs  
■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint

■ vomit blood    ■ have bloody or black stools  
■ have stomach pain that does not get better  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ you have difficulty swallowing  
■ it feels like the pill is stuck in your throat  
■ redness or swelling is present in the painful area  
■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

■ do not take more than directed  
■ the smallest effective dose should be used  
■ drink a full glass of water with each dose

#### Adults and children 12 years and older:

■ take 1 caplet every 8 to 12 hours while symptoms last  
■ for the first dose you may take 2 caplets within the first hour  
■ do not exceed 2 caplets in any 8- to 12-hour period  
■ do not exceed 3 caplets in a 24-hour period

**Children under 12 years:** ■ ask a doctor

### Other information

■ each caplet contains: sodium 21 mg  
■ store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

**Drug Facts (continued)**  
**Inactive ingredients:** FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

**Questions or comments? 1-800-719-9200**

This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Aleve® Caplets.

**DISTRIBUTED BY: RITE AID**  
**30 HUNTER LANE, CAMP HILL, PA 17011**

**IF YOU'RE NOT SATISFIED, WE'LL HAPPILY REFUND YOUR MONEY.**

**RITE AID**  
**NOIŃE SATEFACTO**  
**GUARANTEE**

## NAPROXEN SODIUM

naproxen sodium tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-0507
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

### Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6B30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POVIDONES (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	BLUE (Light Blue)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (oval)	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	L368
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0507-1	1 in 1 CARTON		
1		50 in 1 BOTTLE		
2	NDC:11822-0507-7	1 in 1 CARTON		
2		100 in 1 BOTTLE		
3	NDC:11822-0507-8	1 in 1 CARTON		
3		200 in 1 BOTTLE		
4	NDC:11822-0507-0	300 in 1 BOTTLE		
5	NDC:11822-0507-2	1 in 1 CARTON		
5		120 in 1 BOTTLE		
6	NDC:11822-0507-3	1 in 1 CARTON		
6		24 in 1 BOTTLE		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074661	05/16/2001	

**Labeler** - Rite Aid Corporation (014578892)

Revised: 6/2014

Rite Aid Corporation