

## **CHILDRENS IBUPROFEN - ibuprofen suspension 7-Eleven**

-----

### ***Drug Facts***

#### ***Active ingredient (in each 5 mL)***

Ibuprofen USP 100 mg (NSAID)\*

#### ***Purpose***

Pain reliever/fever reducer

\*nonsteroidal anti-inflammatory drug

#### ***Uses***

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

#### ***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Do not use**

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
  - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech

- leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6 to 8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

**Dosing Chart**

<b>Weight (lb)</b>	<b>Age (year)</b>	<b>Dose (mL)*</b>
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

\* or as directed by a doctor

**Other information**

- **each 5 mL contains:** sodium 2 mg
- store between 20 to 25°C (68 to 77°F)
- **do not use if carton is opened or seal under cap opening is broken or missing**
- see bottom panel for lot number and expiration date

**Inactive ingredients**

Acesulfame potassium, art raspberry flavor, citric acid anhydrous, D&C yellow No. 10,

FD&C red No. 40, glycerin, hypromellose, polysorbate 80, pregelatinized starch (potato), purified water, sodium benzoate, sucrose and xanthan gum

**Questions or comments?**

Call **1-855-274-4122**

Distributed by:  
**7-ELEVEN, INC.**  
 IRVING, TX 75063  
 www.7-ELEVEN.COM

Made in India

Code: TS/DRUGS/19/1993

**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL 4 FL OZ (120 mL)  
 Container Label**

**24/7 life BY 7-ELEVEN™  
 CHILDREN'S  
 Ibuprofen  
 Oral Suspension USP  
 100 mg per 5 mL  
 Pain Reliever/  
 Fever Reducer (NSAID)  
 Alcohol Free . Lasts up to 8 Hours  
 AGES 2 TO 11 YEARS  
 Berry Flavored 4FL OZ (120 mL)**

**Important:** Read all product information before using. Keep this box for important information. This product is intended for use in children.

**Purpose**  
 Ibuprofen USP (100 mg (NSAID)).....Pain reliever/fever reducer  
 nonsteroidal anti-inflammatory drug

**Uses temporarily:** ■ relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache ■ reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin redness ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child: ■ has had stomach ulcers or bleeding problems ■ takes a blood thinning (anticoagulant) or steroid drug ■ takes other drugs containing prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ takes more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart failure, and stroke. These warnings: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than directed or for longer than directed. **Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. **Do not use:** ■ if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer, ■ right before or after heart surgery. **Ask a doctor before use if:** ■ stomach bleeding warning applies to your child ■ child has a history of stomach problems, such as heartburn ■ child has problems or serious side effects from taking pain relievers or fever reducers ■ child has not been drinking fluids ■ child has lost a lot of fluid due to vomiting or diarrhea ■ child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke ■ child has asthma ■ child is taking a diuretic. **Ask a doctor or pharmacist before use if the child is:** ■ under a doctor's care for any serious condition ■ taking any other drug. When using this product ■ take with food or milk if stomach upset occurs. **Stop use and ask a doctor if:** ■ child experiences any of the following signs of stomach bleeding: ■ feels faint ■ vomits blood ■ has bloody or black stools ■ has stomach pain that does not get better ■ child has symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ the child does not get any relief within first day (24 hours) of treatment ■ fever or pain gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions** ■ This product does not contain directions or complete warnings for adult use ■ do not give more than directed ■ shake well before using ■ ml = milliliter ■ find right dose on chart below. If possible, use weight to dose; otherwise use age ■ use only enclosed dosing cup. Do not use any other dosing device ■ if needed, repeat dose every 6 to 8 hours ■ do not use more than 4 times a day ■ replace original bottle cap to maintain child resistance ■ wash dosing cup after each use

Dosing Chart	
Weight (lb)	Dose (mL)*
under 24	ask a doctor
under 2 years	ask a doctor
24-35 lbs	2-3 years 5 mL
36-47 lbs	4-5 years 7.5 mL
48-59 lbs	6-8 years 10 mL
60-71 lbs	9-10 years 12.5 mL
72-95 lbs	11 years 15 mL

\* or as directed by a doctor

**Other information** ■ each 5 mL contains: sodium 2 mg ■ store between 20 to 25°C (68 to 77°F) ■ do not use if seal under cap is broken or missing

**Inactive ingredients:** Acesulfame potassium, art raspberry flavor, citric acid anhydrous, D&C yellow No. 10, FD&C red No. 40, glycerin, hypromellose, polysorbate 80, pregelatinized starch (potato), purified water, sodium benzoate, sucrose and xanthan gum.

**Questions or comments?** 1-855-274-4122

DISTRIBUTED BY  
 7-ELEVEN, INC.  
 IRVING, TX 75063  
 www.7-ELEVEN.COM  
 Made in India  
 Code: TS/DRUGS/19/1993

LM-3571

LOT \_\_\_\_\_  
 EXP \_\_\_\_\_

**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL 4 FL OZ (120 mL) Carton Label**

**24/7 life BY 7-ELEVEN™**  
**CHILDREN'S**  
**Ibuprofen**  
**Oral Suspension USP**  
**100 mg per 5 mL**  
**Pain Reliever/  
Fever Reducer (NSAID)**  
**Alcohol Free . Lasts up to 8 Hours**  
**AGES 2 TO 11 YEARS**  
**Use ONLY enclosed dosage cup**      **Berry Flavor**  
**4FL OZ (120mL)**

**KEEP IN OUTER CARTON UNTIL CONTENTS USED**

**Drug Facts (continued)**

**Dosing Chart**

Weight (lb)	Age (year)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

\* or as directed by a doctor

**Other information**

- each 5 mL contains sodium 2 mg
- store between 20 to 25°C (68 to 77°F)
- do not use if carton is opened or seal under cap opening is broken or missing
- see bottom panel for lot number and expiration date

**Inactive ingredients**


Acesulfame potassium, art raspberry flavor, citric acid anhydrous, D&C yellow No. 10, FD&C red No. 40, glycerin, hypromellose, polysorbate 80, pregelatinized starch (potato), purified water, sodium benzoate, sucrose and xanthan gum

**Questions or comments?** ☎ 1-855-274-4122

DISTRIBUTED BY:  
7-ELEVEN, INC.  
IRVING, TX 75062  
www.7-ELEVEN.COM

Made in India

Code: TS/DRUGS/19/1993




**CHILDREN'S  
Ibuprofen**

Oral Suspension USP  
100 mg per 5 mL

Berry Flavor      4 FL OZ (120 mL)

LM-3572




**CHILDREN'S  
Ibuprofen**

Oral Suspension USP  
100 mg per 5 mL

Pain Reliever/  
Fever Reducer (NSAID)


Alcohol Free . Lasts up to 8 Hours

**AGES 2 TO 11 YEARS**




Use ONLY enclosed dosage cup

**Berry Flavor**



4 FL OZ (120 mL)



0 52548 69064 9

LOT:  
EXP:

**Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.**

**Drug Facts**

**Active ingredient (in each 5 mL)**  
Ibuprofen USP 100 mg (NSAID)\*.....Pain reliever/fever reducer

\*nonsteroidal anti-inflammatory drug

**Uses temporarily:**

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- stomach pain
- reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Do not use**

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids

**Drug Facts (continued)**

- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding:
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling

- child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every 6 to 8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

Unvarnished Zone  
(dotted line not for printing)

Unvarnished Zone  
(dotted line not for printing)

Unvarnished Zone  
(dotted line not for printing)

Unvarnished Zone  
(dotted line not for printing)

**CHILDRENS IBUPROFEN**  
ibuprofen suspension

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:10202-711
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg in 5 mL

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>ACESULFAME POTASSIUM</b> (UNII: 230V73Q5G9)	
<b>RASPBERRY</b> (UNII: 4N14V5R27W)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>HYPROMELLOSE 2208 (4000 MPA.S)</b> (UNII: 39J80LT57T)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>STARCH, POTATO</b> (UNII: 8I089SAH3T)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Product Characteristics

<b>Color</b>	ORANGE (Light orange to orange)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10202-711-24	1 in 1 CARTON	05/06/2019	
1		120 mL in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA209179	05/06/2019	

**Labeler** - 7-Eleven (007347602)

## **Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Aurobindo Pharma Limited		918917642	ANALYSIS(10202-711) , MANUFACTURE(10202-711)

Revised: 11/2022

7-Eleven