

APEXICON E- diflorasone diacetate cream
PharmaDerm a division of Fougera Pharmaceuticals Inc.

ApexiCon[®] E Cream
(diflorasone diacetate cream USP, 0.05% [emollient])

Rx only


NOT FOR OPHTHALMIC USE

DESCRIPTION

Each gram of ApexiCon[®] E Cream (diflorasone diacetate cream USP, 0.05% [emollient]) contains 0.5 mg diflorasone diacetate in a cream base for topical dermatological use.

Chemically, diflorasone diacetate is: 6 α ,9-difluoro-11 β ,17,21-trihydroxy-16 β -methylpregna-1,4-diene-3,20-dione 17,21-diacetate.

The structural formula is represented below:



Each gram of ApexiCon[®] E Cream (diflorasone diacetate cream USP, 0.05% [emollient]) contains: 0.5 mg diflorasone diacetate in a hydrophilic vanishing cream base of propylene glycol, stearyl alcohol, cetyl alcohol, sorbitan monostearate, polysorbate 60, mineral oil and purified water.

CLINICAL PHARMACOLOGY

Topical corticosteroids share anti-inflammatory, antipruritic and vasoconstrictive actions. The mechanism of anti-inflammatory activity of the topical corticosteroids is unclear. Various laboratory methods, including vasoconstrictor assays, are used to compare and predict potencies and/or clinical efficacies of the topical corticosteroids. There is some evidence to suggest that a recognizable correlation exists between vasoconstrictor potency and therapeutic efficacy in man.

Pharmacokinetics: The extent of percutaneous absorption of topical corticosteroids is determined by many factors including the vehicle, the integrity of the epidermal barrier, and the use of occlusive dressings. Topical corticosteroids can be absorbed from normal intact skin. Inflammation and/or other disease processes in the skin increase percutaneous absorption. Occlusive dressings substantially increase the percutaneous absorption of topical corticosteroids. Thus, occlusive dressings may be a valuable therapeutic adjunct for treatment of resistant dermatoses. (See **DOSAGE AND ADMINISTRATION.**)

Once absorbed through the skin, topical corticosteroids are handled through pharmacokinetic pathways similar to systemically administered corticosteroids. Corticosteroids are bound to plasma proteins in varying degrees. They are metabolized primarily in the liver and are then excreted by the kidneys. Some of the topical corticosteroids and their metabolites are also excreted into the bile.

INDICATIONS AND USAGE

Topical corticosteroids are indicated for relief of the inflammatory and pruritic manifestations of corticosteroid responsive dermatoses.

CONTRAINDICATIONS

Topical steroids are contraindicated in those patients with a history of hypersensitivity to any of the components of the preparation.

PRECAUTIONS

General: Systemic absorption of topical corticosteroids has produced reversible hypothalamic-pituitary-adrenal (HPA) axis suppression, manifestations of Cushing's syndrome, hyperglycemia, and glucosuria in some patients.

Conditions which augment systemic absorption include the application of the more potent steroids, use over large surface areas, prolonged use, and the addition of occlusive dressings.

Therefore, patients receiving a large dose of a potent topical steroid applied to a large surface area or under an occlusive dressing should be evaluated periodically for evidence of HPA axis suppression by using the urinary free cortisol and ACTH stimulation tests. If HPA axis suppression is noted, an attempt should be made to withdraw the drug, to reduce the frequency of application, or to substitute a less potent steroid.

Recovery of HPA axis function is generally prompt and complete upon discontinuation of the drug. Infrequently, signs and symptoms of steroid withdrawal may occur, requiring supplemental systemic corticosteroids.

Pediatric patients may absorb proportionally larger amounts of topical corticosteroids and thus be more susceptible to systemic toxicity. (See **PRECAUTIONS: Pediatric Use**).

If irritation develops, topical corticosteroids should be discontinued and appropriate therapy instituted.

In the presence of dermatological infections, the use of an appropriate antifungal or antibacterial agent should be instituted. If a favorable response does not occur promptly, the corticosteroid should be discontinued until the infection has been adequately controlled.

Information for the Patient: Patients using topical corticosteroids should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes.
2. Patients should be advised not to use this medication for any disorder other than that for which it was prescribed.
3. The treated skin area should not be bandaged or otherwise covered or wrapped as to be occlusive unless directed by the physician.
4. Patients should report any signs of local adverse reactions especially under occlusive dressing.
5. Parents of pediatric patients should be advised not to use tight-fitting diapers or plastic pants on an infant or child being treated in the diaper area, as these garments may constitute occlusive dressings.

Laboratory Tests: The following tests may be helpful in evaluating the HPA axis suppression: Urinary free-cortisol test; ACTH stimulation test.

Carcinogenesis, Mutagenesis, and Impairment of Fertility: Long-term animal studies have not been

performed to evaluate the carcinogenic potential or the effect on fertility of topical corticosteroids. Studies to determine mutagenicity with prednisolone and hydrocortisone have revealed negative results.

Pregnancy Category C: Corticosteroids are generally teratogenic in laboratory animals when administered systemically at relatively low dosage levels. The more potent corticosteroids have been shown to be teratogenic after dermal application in laboratory animals. There are no adequate and well-controlled studies in pregnant women on teratogenic effects from topically applied corticosteroids. Therefore, topical corticosteroids should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Drugs of this class should not be used extensively on pregnant patients, in large amounts, or for prolonged periods of time.

Nursing Mothers: It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in breast milk. Systemically administered corticosteroids are secreted into breast milk in quantities **not** likely to have a deleterious effect on the infant. Nevertheless, caution should be exercised when topical corticosteroids are administered to a nursing woman.

Pediatric Use: Safety and effectiveness of diflorasone diacetate emollient cream in pediatric patients have not been established. Because of a higher ratio of skin surface area to body mass, pediatric patients are at a greater risk than adults of HPA-axis suppression when they are treated with topical corticosteroids. They are, therefore, also at greater risk of glucocorticosteroid insufficiency after withdrawal of treatment and of Cushing's syndrome while on treatment. Adverse effects including striae have been reported with inappropriate use of topical corticosteroids in pediatric patients.

HPA axis suppression, Cushing's syndrome, and intracranial hypertension have been reported in pediatric patients receiving topical corticosteroids. Manifestations of adrenal suppression in pediatric patients include linear growth retardation, delayed weight gain, low plasma cortisol levels, and absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema.

Administration of topical corticosteroids to pediatric patients should be limited to the least amount compatible with an effective therapeutic regimen. Chronic corticosteroid therapy may interfere with the growth and development of pediatric patients.

ADVERSE REACTIONS

The following local adverse reactions have been reported with topical corticosteroids, but may occur more frequently with the use of occlusive dressings. These reactions are listed in an approximate decreasing order of occurrence: burning, itching, irritation, dryness, folliculitis, hypertrichosis, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, maceration of the skin, secondary infection, skin atrophy, striae, and miliaria.

OVERDOSAGE

Topically applied corticosteroids can be absorbed in sufficient amounts to produce systemic effects. (See **PRECAUTIONS**).

DOSAGE AND ADMINISTRATION

ApexiCon[®] E Cream (diflorasone diacetate cream USP, 0.05% [emollient]) should be applied to the affected areas as a thin film from one to three times daily depending on the severity or resistant nature of the condition.

Occlusive dressings may be used for the management of psoriasis or recalcitrant conditions. If an infection develops, the use of occlusive dressings should be discontinued and appropriate antimicrobial therapy initiated.

HOW SUPPLIED

ApexiCon[®] E Cream (diflorasone diacetate cream USP, 0.05% [emollient]) is available in the following size tubes:

NDC 10337-395-30 30 gram tube

NDC 10337-395-60 60 gram tube

Store at controlled room temperature 20° to 25°C (68° to 77°F) [see USP].

PharmaDerm[®]
A division of
FougeraPharmaceuticals Inc.
Melville, New York 11747 USA
www.pharmaderm.com

I8395F/IF8395F

R04/16

#35

PACKAGE LABEL – PRINCIPAL DISPLAY PANEL – 60 G LABEL

NDC 10337-395-60


PharmaDerm[®]

ApexiCon[®] E Cream

(diflorasone diacetate cream USP 0.05% [emollient])

FOR EXTERNAL USE ONLY NOT FOR OPHTHALMIC USE

Rx only 60 g



Each gram contains: 0.5 mg of diflorasone diacetate in a vehicle consisting of propylene glycol, stearyl alcohol, cetyl alcohol, sorbitan monostearate, polysorbate 60, mineral oil and purified water.

USUAL DOSAGE: See package insert for full prescribing information.

WARNING: Keep out of reach of children.

Keep tightly closed.

Store at controlled room temperature, 20° to 25°C (68° to 77°F) [see USP].

See crimp of tube for Lot No. and Exp. Date.

46131768B
R03/15

PharmaDerm[®]
A division of Fougera Pharmaceuticals Inc.
Melville, NY 11747 USA
www.pharmaderm.com



PACKAGE LABEL – PRINCIPAL DISPLAY PANEL – 60 G CARTON

NDC 10337-395-60


PharmaDerm®

ApexiCon® E Cream

(diflorasone diacetate cream USP 0.05% [emollient])

FOR EXTERNAL USE ONLY NOT FOR OPHTHALMIC USE

Rx only 60 g



APEXICON E

diflorasone diacetate cream

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:10337-395
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
diflorasone diacetate (UNII: 7W2J09SCWX) (diflorasone - UNII:T2DHJ9645W)	diflorasone diacetate	0.5 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
Stearyl Alcohol (UNII: 2KR89I4H1Y)	

Cetyl Alcohol (UNII: 936JST6JCN)	
Sorbitan Monostearate (UNII: NVZ4I0H58X)	
Mineral Oil (UNII: T5L8T28FGP)	
Water (UNII: 059QF0KO0R)	
Polysorbate 60 (UNII: CAL22UVI4M)	
Propylene Glycol (UNII: 6DC9Q167V3)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10337-395-30	1 in 1 CARTON	12/20/2002	
1		30 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:10337-395-60	1 in 1 CARTON	12/20/2002	
2		60 g in 1 TUBE; Type 0: Not a Combination Product		
3	NDC:10337-395-66	6 in 1 CARTON	12/20/2002	
3		2 g in 1 TUBE; Type 0: Not a Combination Product		
4	NDC:10337-395-08	2 g in 1 TUBE; Type 0: Not a Combination Product	12/20/2002	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076263	12/20/2002	

Labeler - PharmaDerm a division of Fougera Pharmaceuticals Inc. (043838424)

Revised: 4/2016

PharmaDerm a division of Fougera Pharmaceuticals Inc.