MILK OF MAGNESIA- magnesium hydroxide suspension CARDINAL HEALTH

ldr mom CHRY

Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are presently taking a prescription drug.

This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- follow each dosage with a full glass (8 oz) of liquid

age	dose
adults and children 12 years and over	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

Other information

- each 15 mL contains: magnesium 500 mg, sodium 6 mg
- do not freeze
- store at room temperature tightly closed

Inactive ingredients

carboxymethylcellulose sodium, citric acid, D & C red no. 28, flavor, glycerin, microcrystalline cellulose, purified water, sodium citrate, sodium hypochlorite, sucrose, xanthan gum

Questions or comments?

1-800-540-3765

package Label



TAMPER EVIDENT: Do not use if imprinted neckband is Drug Facts missing or broken.

Active ingredient (in each 15 mL) Magnesium hydroxide 1200 mg

Purpose Saline laxative

Use • relieves occasional constipation (irregularity)

generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have • a magnesium-restricted diet

kidney disease

· stomach pain, nausea or vomiting

a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. This product may interact with certain prescription drugs

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor

follow each dosage with a full glass (8 oz) of liquid

age	dose
adults and children 12 years and over	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

Other information

- each 15 mL contains: magnesium 500 mg, sodium 6 mg
- do not freeze store at room temperature tightly closed

Inactive ingredients carboxymethylcellulose sodium, citric acid, D&C red no. 28, flavor, glycerin, microcrystalline cellulose, punified water, sodium citrate, sodium hypochlorite, sucrose, xanthan gum

Questions or comments? 1-800-540-3765

"This product is not manufactured or distributed by Bayer Healthcare LLC, distributors of Phillips® brand Milk of Magnesia. 995-19312-1 REV GC-0722

©2022 CARDINAL HEALTH, LEADER, and LOGOs are trademarks of Cardinal Health. All other marks are the property of their respective owners.



100% Money Back Guarantee Return to place of purchase if not satisfied.

MILK OF MAGNESIA

magnesium hydroxide suspension

Product Information

MAGN UNII:T

Product Type HUMAN OTC DRUG Item Code (Source) NDC:70000-0064

DISTRIBUTED BY CARDI
DUBLIN, OHIO 43017
www.myleader.com

BY CARDINAL HEALTH

ORAL **Route of Administration**

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION -	MAGNESIUM	1200 mg
T6V3LHY838. HYDROXIDE ION - UNII:9159UV381P)	HYDROXIDE	in 15 mL

Inactive Ingredients				
Ingredient Name	Strength			
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
D&C RED NO. 28 (UNII: 767IP0Y5NH)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
GLYCERIN (UNII: PDC6A3C0OX)				
XANTHAN GUM (UNII: TTV12P4NEE)				
SODIUM CITRATE (UNII: 1Q73Q2JULR)				
WATER (UNII: 059QF0KO0R)				
SODIUM HYPOCHLORITE (UNII: DY38VHM5OD)				
SUCROSE (UNII: C151H8M554)				
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K6790BS311)				

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

l	Packaging				
	# Item Code Package Description		Marketing Start Date	Marketing End Date	
	1	NDC:70000- 0064-1	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/01/2020	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	09/01/2020	

Labeler - CARDINAL HEALTH (063997360)

Registrant - GCP Laboratories (965480861)

Establishment				
Name	Address	ID/FEI	Business Operations	
GCP Laboratories		965480861	manufacture(70000-0064)	

Revised: 11/2023 CARDINAL HEALTH