

FIBER CAPLETS- calcium polycarbophil tablet

Strategic Sourcing Services LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Fiber Laxative

Active ingredient (in each caplet)

Calcium polycarbophil 625 mg equivalent to 500 mg polycarbophil

Purpose

Bulk-forming laxative

uses

- relieves occasional constipation
- this product generally produces bowel movement in 12 to 72 hours

Warnings

Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Ask a doctor before use if you have

- abdominal pain, nausea or vomiting
- difficulty in swallowing
- a sudden change in bowel habits that persists over a period of 2 weeks

Ask a doctor or pharmacist before use if you are

taking any other drug. Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work.

When using this product

- do not use for more than 7 days unless directed by a doctor
- do not take more than 8 caplets in a 24 hour period unless directed by a doctor.

Stop use and ask a doctor if

- you experience chest pain, abdominal pain, nausea, vomiting, difficulty in breathing or swallowing
- you fail to have a bowel movement after use or have rectal bleeding. These could be

signs of a serious condition.

Keep out of the reach of children

In case of overdose, get medical help or contact a Poison Control Center immediately.

Directions

- take this product (child or adult dose) with a full glass of water (8 oz.) or other fluid. Taking this product without enough liquid may cause choking. See choking warning.
- dosage will vary according to diet, exercise, previous laxative use or severity of constipation.
- continued use for 1 or 3 days is normally required to provide full benefit.

| | |
|---------------------------------------|-------------------------------|
| adults and children 12 years and over | 2 caplets, 1 to 4 times a day |
| children under 12 years | ask a doctor |

Other information

- **each caplet contains:** calcium 135mg
- store at room temperature
- protect contents from moisture

Inactive ingredients

Calcium carbonate, caramel, colloidal silicon dioxide, crospovidone, hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate

PDP

sunmark

NDC 49348-190-13

*COMPARE TO FIBERCON CAPLETS

fiber caplets

Calcium Polycarbophil 625 mg

Relieves occasional constipation

Bulk forming laxative


90 Caplets

sunmark®
*COMPARE TO
 FIBERCON® CAPLETS
 NDC 49348-190-13

fiber caplets

Calcium Polycarbophil 625 mg

Relieves occasional constipation
 Bulk forming laxative



90 CAPLETS

WARNING: Consuming this product can expose you to chemicals including lead, which is known to the State of California to cause cancer and birth defects or other reproductive harm. For more information go to www.P65Warnings.ca.gov/food.

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING.

Drug Facts

Active ingredient (in each caplet) Purpose
 Calcium polycarbophil 625 mg equivalent to 500 mg polycarbophil...Bulk-forming laxative

Uses ■ relieves occasional constipation
 ■ this product generally produces bowel movement in 12 to 72 hours

Warnings

Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing, if you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Ask a doctor before use if you have

- abdominal pain, nausea or vomiting
- difficulty in swallowing
- a sudden change in bowel habits that persists over a period of 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work.

When using this product

- do not use for more than 7 days unless directed by a doctor
- do not take more than 8 caplets in a 24-hour period unless directed by a doctor

LOT:

MFR#53041

REV 0419



EXP:

▶ Peel Here for Continuing Drug Facts

Stop Peeling Here

MEKESON
 Active Quality Product Available by McKesson
 855 Star Line, Inc., Irving, TX 75039
 Many local pharmacies
 Please visit us at www.mckessonpharm.com

*This product is not manufactured or distributed by Pfizer Consumer Healthcare, the distributor of Fibercon®.

Inactive ingredients calcium carbonate, caramel, colloidal silicon dioxide, croscopolone, hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate.

Other information ■ each caplet contains: calcium 135 mg ■ store at room temperature ■ protect contents from moisture

Directions ■ take this product (child or adult dose) with a full glass of water (8oz.) or other fluid. Taking this product without enough liquid may cause choking. See choking warning. ■ dosage will vary according to diet, exercise, previous laxative use or severity of constipation ■ continued use for 1 to 3 days is normally required to provide full benefit

| | |
|-----------------------------------|-------------------------------|
| adults & children 12 years & over | 2 caplets, 1 to 4 times a day |
| children under 12 years | ask a doctor |

Stop use and ask a doctor if
 ■ you experience chest pain, abdominal pain, nausea, vomiting, difficulty in breathing or swallowing
 ■ you fail to have a bowel movement after use or have rectal bleeding. These could be signs of a serious condition.

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center immediately.

Drug Facts (continued)

| FIBER CAPLETS | | | |
|--|-----------------------|--------------------|---------------|
| calcium polycarbophil tablet | | | |
| Product Information | | | |
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:49348-190 |
| Route of Administration | ORAL | | |
| Active Ingredient/Active Moiety | | | |
| Ingredient Name | Basis of Strength | Strength | |
| CALCIUM POLYCARBOPHIL (UNII: 8F049NKY49) (POLYCARBOPHIL - UNII:W25LM17A4W) | CALCIUM POLYCARBOPHIL | 625 mg | |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| CALCIUM CARBONATE (UNII: H0G9379FGK) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| CROSPVIDONE (UNII: 2S7830E561) | |
| HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| POLYETHYLENE GLYCOL 1000 (UNII: U076Q6Q621) | |
| SODIUM LAURYL SULFATE (UNII: 368GB5141J) | |
| CARAMEL (UNII: T9D99G2B1R) | |

Product Characteristics

| | | | |
|-----------------|---------|---------------------|----------|
| Color | white | Score | no score |
| Shape | capsule | Size | 19mm |
| Flavor | | Imprint Code | G147 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:49348-190-13 | 90 in 1 BOTTLE; Type 0: Not a Combination Product | 09/18/2019 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|-------------------------|--|----------------------|--------------------|
| OTC monograph not final | part334 | 09/18/2019 | |

Labeler - Strategic Sourcing Services LLC (116956644)

Registrant - Guardian Drug Company (119210276)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-----------------------|---------|-----------|------------------------|
| Guardian Drug Company | | 119210276 | manufacture(49348-190) |