

**SENNA SYRUP- sennosides liquid
AVPAK**

Senna Syrup

Drug Facts

Active Ingredient (in each 5 mL = 1 teaspoonful)

Sennosides 8.8 mg

Purpose

Laxative

Use: • relieves occasional constipation (irregularity), generally causes bowel movement in 6-12 hours.

WARNINGS: Do not use • laxative products for a period longer than 1 week unless directed by a doctor

Ask a doctor before use if you have • stomach pain • nausea • vomiting • noticed a sudden change in bowel movement that continues over a period of 2 weeks

Stop use and ask a doctor if you have • rectal bleeding • no bowel movement after use of laxative

These could be signs of a serious condition.

If pregnant or breast feeding, ask a health care professional before use.

Keep this and all drugs out of reach of children. In case of overdose, get medical help or contact a Poison Control Center immediately. In case of eye contact, flush with water.

Directions: <ul style="list-style-type: none">• Shake well before use• Do not exceed recommended dose		
Age	Starting Dose	Maximum Dosage
Adults and children 12 years	2 to 3 teaspoonfuls (10 mL to 15 mL) once a day preferably at bedtime.	3 teaspoonfuls (15 mL) in the morning and 3

of age and over	preferably at bedtime, increase as needed or as recommended by a doctor	teaspoonfuls (15 mL) at bedtime
Children under 12 years of age	Ask a doctor	Ask a doctor

1 teaspoonful = 5 mL

Other information: Store at room temperature 15°C to 25°C (59°F to 77°F). Excursions between 26°C to 30°C (78°F to 86°F) are allowed. Protect from excessive heat. For more info call 1- 800-447-1006.

Inactive ingredients: Artificial and natural chocolate flavor, methylparaben, propylparaben, propylene glycol, sucrose and purified water.

Principal Display Panel

NDC 50268-731-24

SENNA Syrup

Natural
Vegetable Laxative

24 Units of 8.8 mg/5 mL



5026873124

Drug Facts

Active Ingredient (per teaspoonful-5mL)	Purpose
Sennosides 8.8 mg	Laxative

Uses

- relieves occasional constipation (irregularity)
- generally causes bowel movement in 6 to 12 hours

Warnings

Do not use

- laxative products for longer than 1 week unless directed by a doctor

Ask a doctor a pharmacist before use if you have

- stomach pain ■ nausea ■ vomiting
- noticed a sudden change in bowel movement that continues over a period of 2 weeks

Stop use and ask a doctor if

you have rectal bleeding or failure to have a bowel movement after use of laxative. These may indicate a serious condition.

If pregnant or breast-feeding ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center immediately.

Mfg. Rev. 03/19

AV Rev. 05/21 (P)



Peel
Here

SENNA SYRUP

sennosides liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50268-731
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SENNOSIDES (UNII: 3FYP5M0JJX) (SENNOSIDES - UNII:3FYP5M0JJX)	SENNOSIDES	8.8 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0K00R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50268-731-24	24 in 1 BOX	05/03/2021	
1	NDC:50268-731-11	5 mL in 1 CUP, UNIT-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	05/03/2021	

Labeler - AVPAK (832926666)

Revised: 3/2024

AVPAK