

**FIRST AID SHOT THERAPY ALLERGY RELIEF- diphenhydramine hydrochloride solution**  
**First Aid Beverages, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**First Aid Shot Therapy® Allergy Relief®**

***Drug Facts***

**Active ingredient**

Diphenhydramine HCl 50 mg

**Purpose (in each bottle)**

Antihistamine

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose and throat
- temporarily relieves these symptoms due to the common cold:
  - runny nose
  - sneezing

**Warnings**

**Do not use**

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are** taking sedatives or tranquilizers

**When using this product**

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**If pregnant or breast-feeding,** ask a health care professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control

Center right away.

### **Directions**

- Adults and children 12 years and over:
  - swallow the contents of one bottle
  - repeat every 4-6 hours while symptoms persist, or as directed by doctor
  - do not take more than 6 bottles in 24 hours
- Children under 12 years: Do not use

### **Other information**

- each 40 ml contains: sodium 44 mg
- store between 20-25C°(68-77°F)

### **Inactive ingredients**

citric acid, flavor, glycerol, polyoxyl 40 stearate, purified water, sodium benzoate, sodium citrate, sucrose

### **Questions or comments?**

1-877-958-8331 (Monday-Friday 9AM-5PM PST). You may also report serious side effects to this number.

Distributed by  
First Aid Shot Therapy®  
Burlingame, CA 94010

### **PRINCIPAL DISPLAY PANEL - 40mL Bottle Label**

**ALLERGY**

**FAST  
-LIQUID-  
RELIEF®**

**FIRST AID  
SHOT THERAPY®**

**TROPICAL**

**FDA  
COMPLIANT**

Diphenhydramine HCl  
(Antihistamine)

1.35 FL OZ (40mL)

# • ALLERGY • ALLERGY • ALLERGY

Do not use if seal is broken

We hope F.A.S.T.  
relieved your  
allergy symptoms.

For more info visit:  
[www.firstaidshottherapy.com](http://www.firstaidshottherapy.com)

1018-010-000

## FAST — LIQUID — RELIEF®



FIRST AID  
SHOT THERAPY®

TROPICAL

FDA  
COMPLIANT

Diphenhydramine HCl  
(Antihistamine)

1.35 FL. OZ (40mL)



Single Dose  
Relieves:

- Sneezing
- Itchy Eyes
- Runny Nose
- Itchy Throat

### Drug Facts

**Active ingredients Purpose**  
(in each bottle)

Diphenhydramine HCl 50 mg..  
.....Antihistamine

1017-010-000



LIFT PANEL FOR MORE DRUG FACTS  
INFORMATION. READ ALL WARNINGS  
UNDER LABEL BEFORE USE.

### Drug Facts (continued)

#### Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose and throat
- temporarily relieves these symptoms due to the common cold:
  - runny nose
  - sneezing

#### Warnings

##### Do not use

- to make a child sleepy

### Drug Facts (continued)

- with any other product containing diphenhydramine, even one used on skin

#### Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

### Drug Facts (continued)

- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

If pregnant or breast-feeding, ask a health care professional before use.

**Keep out of reach of children.** In case of

<ul style="list-style-type: none"> <li>• a sodium-restricted diet</li> </ul>
<b>Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers</b>
<b>When using this product</b> <ul style="list-style-type: none"> <li>• marked drowsiness may occur</li> <li>• avoid alcoholic drinks</li> </ul>

overdose, get medical help or contact a Poison Control Center right away.
<b>Directions</b>
<ul style="list-style-type: none"> <li>• Adults and children 12 years and over: <ul style="list-style-type: none"> <li>• swallow the contents of one bottle</li> </ul> </li> </ul>

<b>Drug Facts (continued)</b>
<ul style="list-style-type: none"> <li>• repeat every 4-6 hours while symptoms persist, or as directed by doctor</li> <li>• do not take more than 6 bottles in 24 hours</li> <li>• Children under 12 years: Do not use</li> </ul>
<b>Other information</b>
<ul style="list-style-type: none"> <li>•each 40 ml contains: sodium 44 mg</li> <li>•store between 20-25°C(68-77°F)</li> </ul>
<b>Inactive ingredients</b>
citric acid, flavor, glycerol, polyoxyl 40 stearate, purified water, sodium benzoate, sodium citrate, sucrose

<b>Drug Facts (continued)</b>
<b>Questions or comments?</b>
<b>1-877-958-8331</b> (Monday-Friday 9AM-5PM PST). You may also report serious side effects to this number.
Distributed by First Aid Shot Therapy® Burlingame, CA 94010 First Aid Shot Therapy® and cross symbol are registered trademarks of First Aid Shot Therapy®. <a href="http://www.firstaidshottherapy.com">www.firstaidshottherapy.com</a>

## FIRST AID SHOT THERAPY ALLERGY RELIEF

diphenhydramine hydrochloride solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:578 15-0 10
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Diphenhydramine Hydrochloride</b> (UNII: TC2D6JAD40) (Diphenhydramine - UNII:8GTS82S83M)	Diphenhydramine Hydrochloride	50 mg in 40 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>POLYOXYL 40 STEARATE</b> (UNII: 13A4J4NH9I)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>WATER</b> (UNII: 059QF0K00R)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	

<b>Flavor</b>	MANGO, PEACH		<b>Imprint Code</b>	
<b>Contains</b>				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:578 15-010-01	1 in 1 CARTON	07/01/2016	
1		40 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC MONOGRAPH FINAL	part341	07/01/2016		

**Labeler** - First Aid Beverages, Inc. (078618157)

Revised: 6/2016

First Aid Beverages, Inc.