

**ACETAMINOPHEN, ASPIRIN, CAFFEINE- acetaminophen, aspirin,
caffeine tablet, film coated
Proficient Rx LP**

5004-Rugby

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID*)

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever

Pain reliever aid

Uses

■ temporarily relieves minor aches and pains due to:

■ headache

■ muscular aches

■ a cold

■ arthritis

■ toothache

■ premenstrual and menstrual cramps

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ with other drugs containing acetaminophen ■ more than 8 caplets in 24 hours, which is the maximum daily amount ■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ take more or for a longer time than directed ■ have had stomach ulcers or bleeding problems

■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing

prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
 - an allergic reaction occurs. Seek medical help right away.
 - ringing in the ears or loss of hearing occurs
 - pain gets worse or lasts for more than 10 days
 - fever gets worse or lasts for more than 3 days
 - painful area is red or swollen
 - any new symptoms appear.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over:
 - take 2 caplets every 6 hours
 - do not take more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

hypromellose, maltodextrin, microcrystalline cellulose, polydextrose, polyvinylpyrrolidone, pregelatinized starch, silica, starch, stearic acid, talc, titanium dioxide, triglycerides

Questions or comments?

1-800-645-2158

RETAIN THIS CARTON FOR COMPLETE PRODUCT INFORMATION

Distributed by: RUGBY® LABORATORIES, Livonia, MI 48152
www.rugbylaboratories.com

Repackaged by: Proficient Rx LP Thousand Oaks, CA 91320

†This product is not manufactured or distributed by GlaxoSmithKline, owner of the registered trademark Excedrin®

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Pain Reliever Plus

NDC 82804-008-60

Compare to the active ingredients in Excedrin® Extra Strength Caplets†

Acetaminophen, Aspirin (NSAID)* and Caffeine

250mg/250mg/65mg

Pain Reliever/Pain Reliever Aid

NOT FOR HOUSEHOLDS WITH YOUNG CHILDREN

60 Caplets



Scan Here



NDC 82804-008-60

Packaged By: Proficient Rx LP
Thousand Oaks, CA 91320



5

Pain Reliever Plus 250mg /250mg /65mg #60 Caplets

Each caplet contains: Acetaminophen 250 mg Pain reliever / Aspirin 250 mg (NSAID*) Pain reliever / Caffeine 65 mg Pain reliever aid *nonsteroidal anti-inflammatory drug

White, unscored, oval shaped caplet, with imprint code 2793

Product ID: ZP000860

Dist. By: RUGBY LABORATORIES, Livonia, MI 48152

Store at 25°C (77°F)

Keep medication out of the reach of children

Pain Reliever Plus 250mg /250mg /65mg #60 Caplets
Lot #:00000 SN# MASTER
NDC 82804-008-60 Exp:00/00/00

Pain Reliever Plus 250mg /250mg /65mg #60 Caplets
Lot #:00000 SN# MASTER
NDC 82804-008-60 Exp:00/00/00

Pain Reliever Plus 250mg /250mg /65mg #60 Caplets
Lot #:00000 SN# MASTER
NDC 82804-008-60 Exp:00/00/00



GTIN: 00382804008605
SN# MASTER
Exp. 00/00/00
Lot #:00000

ACETAMINOPHEN, ASPIRIN, CAFFEINE

acetaminophen, aspirin, caffeine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82804-008(NDC:0536-1326)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg

Inactive Ingredients

Ingredient Name	Strength
POLYDEXTROSE (UNII: VH2XOU12IE)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE K30 (UNII: U725QWY32X)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
STARCH, CORN (UNII: O8232NY3SJ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	2793
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82804-008-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	10/05/2023	
2	NDC:82804-008-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	10/05/2023	
3	NDC:82804-008-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	10/05/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	06/03/2021	

Labeler - Proficient Rx LP (079196022)

Establishment

Name	Address	ID/FEI	Business Operations
Proficient Rx LP		079196022	REPACK(82804-008) , RELABEL(82804-008)

Revised: 11/2023

Proficient Rx LP