

# MYXREDLIN- insulin human injection, solution

## Baxter Healthcare Corporation

-----

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use MYXREDLIN safely and effectively. See full prescribing information for MYXREDLIN.

**MYXREDLIN (insulin human) in sodium chloride injection, for intravenous use**  
**Initial U.S. Approval: 2019**

### INDICATIONS AND USAGE

MYXREDLIN is a short-acting human insulin indicated to improve glycemic control in adults and pediatric patients with diabetes mellitus (1).

### DOSAGE AND ADMINISTRATION

- Inspect MYXREDLIN visually before use. It should appear clear and colorless. Do not use MYXREDLIN if particulate matter or coloration is seen. (2.1)
- Administer MYXREDLIN intravenously ONLY under medical supervision with close monitoring of blood glucose and potassium levels.(2.1)
- Do not add supplementary medication or additives. (2.1)
- Do not use in series connections. (2.1)
- Do not shake or freeze. Discard unused portion. (2.1)
- Individualize dose based on metabolic needs, blood glucose monitoring results, and glycemic control goal. (2.2)
- Dosage adjustments may be needed with changes in nutrition, renal, or hepatic function or during acute illness. (2.2)

### DOSAGE FORMS AND STRENGTHS

Injection: 100 units insulin human in 100 mL of 0.9% sodium chloride (1 unit/mL) in a single-dose container (3)

### CONTRAINDICATIONS

- During episodes of hypoglycemia (4)
- Hypersensitivity to insulin human or any of the excipients in MYXREDLIN (4)

### WARNINGS AND PRECAUTIONS

- *Hyper- or Hypoglycemia with Changes in Insulin Regimen:* Carry out under close medical supervision and increase frequency of blood glucose monitoring. (5.1)
- *Hypoglycemia:* May be life-threatening. Factors which may increase the risk include changes in nutrition and co-administered medication and patients with renal or hepatic impairment. Increased frequency of blood glucose monitoring is recommended in patients at increased risk. (5.2)
- *Hypersensitivity Reactions:* Severe, life-threatening, generalized allergy, including anaphylaxis, can occur. Discontinue MYXREDLIN, monitor, and treat if indicated. (5.3)
- *Hypokalemia:* May be life-threatening. Monitor potassium levels and treat if indicated. (5.4)
- *Fluid Retention and Heart Failure with Concomitant Use of Thiazolidinediones (TZDs):* Observe for signs and symptoms of heart failure; consider dosage reduction or discontinuation if heart failure occurs. (5.5)

### ADVERSE REACTIONS

Adverse reactions observed with insulin human injection include hypoglycemia, allergic reactions, weight gain and edema (6).

**To report SUSPECTED ADVERSE REACTIONS, contact Baxter Healthcare at 1-866-888-2472 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### DRUG INTERACTIONS

- *Drugs that may increase the risk of hypoglycemia:* antidiabetic agents, ACE inhibitors, angiotensin II receptor blocking agents, disopyramide, fibrates, fluoxetine, monoamine oxidase inhibitors, pentoxifylline, pramlintide, salicylates, somatostatin analog (e.g., octreotide), and sulfonamide antibiotics. (7)
- *Drugs that may decrease the blood glucose lowering effect:* atypical antipsychotics, corticosteroids,

danazol, diuretics, estrogens, glucagon, isoniazid, niacin, oral contraceptives, phenothiazines, progestogens (e.g., in oral contraceptives), protease inhibitors, somatropin, sympathomimetic agents (e.g., albuterol, epinephrine, terbutaline), and thyroid hormones. (7)

- *Drugs that may increase or decrease the blood glucose lowering effect:* Alcohol, beta-blockers, clonidine, lithium salts, and pentamidine. (7)
- *Drugs that may blunt the signs and symptoms of hypoglycemia:* beta-blockers, clonidine, guanethidine, and reserpine. (7)

**See 17 for PATIENT COUNSELING INFORMATION.**

**Revised: 6/2020**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

### **2 DOSAGE AND ADMINISTRATION**

2.1 Important Administration Instructions

2.2 Dosage Information

2.3 Dosage Adjustment due to Drug Interactions

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

5.1 Hyperglycemia or Hypoglycemia with Changes in Insulin Regimen

5.2 Hypoglycemia

5.3 Hypersensitivity and Allergic Reactions

5.4 Hypokalemia

5.5 Fluid Retention and Heart Failure with Concomitant Use of PPAR-gamma agonists

### **6 ADVERSE REACTIONS**

### **7 DRUG INTERACTIONS**

### **8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Renal Impairment

8.7 Hepatic Impairment

### **10 OVERDOSAGE**

### **11 DESCRIPTION**

### **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

### **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

### **16 HOW SUPPLIED/STORAGE AND HANDLING**

16.1 How Supplied

16.2 Storage and Handling

### **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

MYXREDLIN is indicated to improve glycemic control in adults and pediatric patients with diabetes mellitus.

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Important Administration Instructions**

- Inspect MYXREDLIN visually before use. It should appear clear and colorless. Do not use MYXREDLIN if particulate matter or coloration is seen.
- Administer MYXREDLIN intravenously ONLY under medical supervision with close monitoring of blood glucose and potassium levels [see Warnings and Precautions (5.2, 5.4)].
- Do not add supplementary medication or additives.
- Do not use in series connections.
- Do not shake. Do not freeze. Discard any unused portion.

#### **2.2 Dosage Information**

- Individualize and adjust the dosage of MYXREDLIN based on the individual's metabolic needs, blood glucose monitoring results, and glycemic control goal.
- Dosage adjustments may be needed with changes in nutrition, changes in renal or hepatic function or during acute illness [see Warnings and Precautions (5.1, 5.2) and Use in Specific Populations (8.6, 8.7)].

#### **2.3 Dosage Adjustment due to Drug Interactions**

- Dosage adjustment may be needed when MYXREDLIN is used concomitantly with certain drugs [see Drug Interactions (7)].

### **3 DOSAGE FORMS AND STRENGTHS**

Injection: 100 units insulin human in 100 mL of 0.9% sodium chloride (1 unit per mL) as a clear, colorless solution in a single-dose container.

### **4 CONTRAINDICATIONS**

MYXREDLIN is contraindicated:

- During episodes of hypoglycemia
- In patients with hypersensitivity to insulin human or any of the excipients in MYXREDLIN

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Hyperglycemia or Hypoglycemia with Changes in Insulin Regimen**

Changes in insulin strength, manufacturer, type, or method of administration may affect glycemic control and predispose to hypoglycemia [see Warnings and Precautions (5.2)] or hyperglycemia. These changes should be made cautiously and under close medical supervision and the frequency of blood glucose monitoring should be increased.

### **5.2 Hypoglycemia**

Hypoglycemia is the most common adverse reaction of all insulins, including MYXREDLIN. Severe hypoglycemia can cause seizures, may lead to unconsciousness, may be life threatening or cause death. Hypoglycemia can impair concentration ability and reaction time.

Hypoglycemia can happen suddenly and symptoms may differ in each individual and change over time in the same individual. Symptomatic awareness of hypoglycemia may be less pronounced in patients with longstanding diabetes, in patients with diabetic nerve disease, in patients using medications that block the sympathetic nervous system (e.g., beta-blockers) [see Drug Interactions (7)], or in patients who experience recurrent hypoglycemia.

#### Risk Factors for Hypoglycemia

Factors which may increase the risk of hypoglycemia include changes in nutrition and co-administered medication [see Drug Interactions (7)]. Patients with renal or hepatic impairment may be at higher risk of hypoglycemia [see Use in Specific Populations (8.6, 8.7)].

#### Risk Mitigation Strategies for Hypoglycemia

Patients and caregivers must be educated to recognize hypoglycemia. In patients at higher risk for hypoglycemia and patients who have reduced symptomatic awareness of hypoglycemia, increased frequency of blood glucose monitoring is recommended.

### **5.3 Hypersensitivity and Allergic Reactions**

Severe, life-threatening, generalized allergy, including anaphylaxis, can occur with MYXREDLIN. Generalized allergy to insulin may manifest as a whole body rash (including pruritus), dyspnea, wheezing, hypotension, tachycardia, or diaphoresis. If hypersensitivity reactions occur, discontinue MYXREDLIN; treat per standard of care and monitor until symptoms and signs resolve. MYXREDLIN is contraindicated in patients who have had hypersensitivity reactions to insulin human or any of the excipients in MYXREDLIN [see Contraindications (4)].

### **5.4 Hypokalemia**

All insulins, including MYXREDLIN, cause a shift in potassium from the extracellular to intracellular space, possibly leading to hypokalemia. Untreated hypokalemia may cause respiratory paralysis, ventricular arrhythmia, and death. Monitor potassium levels and treat if indicated.

### **5.5 Fluid Retention and Heart Failure with Concomitant Use of PPAR-gamma**

## agonists

Thiazolidinediones (TZDs), which are peroxisome proliferator-activated receptor (PPAR)-gamma agonists, can cause dose-related fluid retention, particularly when used in combination with insulin. Fluid retention may lead to or exacerbate heart failure. Patients treated with insulin, including MYXREDLIN, and a PPAR-gamma agonist should be observed for signs and symptoms of heart failure. If heart failure develops, it should be managed according to current standards of care, and discontinuation or dose reduction of the PPAR-gamma agonist must be considered.

## 6 ADVERSE REACTIONS

The following adverse reactions are also discussed elsewhere in the labeling:

- Hypoglycemia [see Warnings and Precautions (5.2)]
- Hypersensitivity Reactions [see Warnings and Precautions (5.3)]
- Hypokalemia [see Warnings and Precautions (5.4)]

### Adverse Reactions from Clinical Studies or Postmarketing Reports

The following additional adverse reactions have been identified during clinical studies or from postmarketing reports with use of insulin human injection. Because some of these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or to establish a causal relationship to drug exposure.

*Adverse reactions associated with insulin initiation and glucose control intensification*  
Intensification or rapid improvement in glucose control has been associated with a transitory, reversible ophthalmologic refraction disorder, worsening of diabetic retinopathy, and acute painful peripheral neuropathy. Over the long-term, improved glycemic control decreases the risk of diabetic retinopathy and neuropathy.

#### *Hypersensitivity reactions*

Severe, life-threatening, generalized allergy, including anaphylaxis.

#### *Hypoglycemia*

Hypoglycemia is the most commonly observed adverse reaction with MYXREDLIN.

#### *Hypokalemia*

MYXREDLIN can cause a shift in potassium from the extracellular to intracellular space, possibly leading to hypokalemia.

#### *Peripheral edema*

Insulins, including MYXREDLIN, may cause sodium retention and edema, particularly if previously poor metabolic control is improved by intensified insulin therapy.

#### *Weight gain*

Weight gain can occur with insulin therapies, including MYXREDLIN, and has been attributed to the anabolic effects of insulin and the decrease in glucosuria.

#### *Immunogenicity*

As with all therapeutic peptides, insulin administration may cause anti-insulin antibodies to form. Increases in titers of anti-insulin antibodies that react with human insulin have been observed in patients treated with subcutaneous insulin human injection.

## 7 DRUG INTERACTIONS

**Table 1: Clinically Significant Drug Interactions with MYXREDLIN**

<b>Drugs that May Increase the Risk of Hypoglycemia</b>	
<i>Drugs:</i>	Antidiabetic agents, ACE inhibitors, angiotensin II receptor blocking agents, disopyramide, fibrates, fluoxetine, monoamine oxidase inhibitors, pentoxifylline, pramlintide, salicylates, somatostatin analog (e.g., octreotide), and sulfonamide antibiotics.
<i>Intervention:</i>	Dose adjustment and increased frequency of glucose monitoring may be required when MYXREDLIN is co-administered with these drugs.
<b>Drugs that May Decrease the Blood Glucose Lowering Effect of MYXREDLIN</b>	
<i>Drugs:</i>	Atypical antipsychotics (e.g., olanzapine and clozapine), corticosteroids, danazol, diuretics, estrogens, glucagon, isoniazid, niacin, oral contraceptives, phenothiazines, progestogens (e.g., in oral contraceptives), protease inhibitors, somatropin, sympathomimetic agents (e.g., albuterol, epinephrine, terbutaline), and thyroid hormones.
<i>Intervention:</i>	Dose adjustment and increased frequency of glucose monitoring may be required when MYXREDLIN is co-administered with these drugs.
<b>Drugs that May Increase or Decrease the Blood Glucose Lowering Effect of MYXREDLIN</b>	
<i>Drugs:</i>	Alcohol, beta-blockers, clonidine, and lithium salts. Pentamidine may cause hypoglycemia, which may sometimes be followed by hyperglycemia.
<i>Intervention:</i>	Dose adjustment and increased frequency of glucose monitoring may be required when MYXREDLIN is co-administered with these drugs.
<b>Drugs that May Blunt Signs and Symptoms of Hypoglycemia</b>	
<i>Drugs:</i>	Beta-blockers, clonidine, guanethidine, and reserpine.
<i>Intervention:</i>	Increased frequency of glucose monitoring may be required when MYXREDLIN is co-administered with these drugs.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

Available data from published studies over decades have not established an association with human insulin use during pregnancy and major birth defects, miscarriage or adverse maternal or fetal outcomes (*see Data*). There are risks to the mother and fetus associated with poorly controlled diabetes in pregnancy (*see Clinical Considerations*). Animal reproduction studies were not performed.

The estimated background risk of major birth defects is 6-10% in women with pre-gestational diabetes with a HbA1c >7 and has been reported to be as high as 20-25% in women with a HbA1c >10. The estimated background risk of miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

#### Clinical Considerations

### *Disease-associated maternal and/or embryo/fetal risk*

Poorly controlled diabetes in pregnancy increases the maternal risk for diabetic ketoacidosis, preeclampsia, spontaneous abortions, preterm delivery, and delivery complications. Poorly controlled diabetes increases the fetal risk for major birth defects, stillbirth, and macrosomia-related morbidity.

### Data

#### *Human Data*

While available studies cannot definitively establish the absence of risk, published data from retrospective studies, open-label, randomized, parallel studies and meta-analyses have not established an association with human insulin use during pregnancy and major birth defects, miscarriage, or adverse maternal or fetal outcomes. All available studies have methodological limitations including lack of blinding, unclear methods of randomization, and small sample size.

## **8.2 Lactation**

### Risk Summary

Available data from published literature suggest that exogenous human insulin products, including insulin human injection, are transferred into human milk. There are no adverse reactions reported in the breastfed infants in the literature. There are no data on the effects of exogenous human insulin products, including MYXREDLIN, on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for MYXREDLIN and any potential adverse effects on the breastfed infant from MYXREDLIN or from the underlying maternal condition.

## **8.4 Pediatric Use**

MYXREDLIN is indicated to improve glycemic control in pediatric patients with diabetes mellitus.

The dosage of MYXREDLIN must be individualized in pediatric patients based on metabolic needs and frequent monitoring of blood glucose to reduce the risk of hypoglycemia [see Dosage and Administration (2.2) and Warnings and Precautions (5.2)].

## **8.5 Geriatric Use**

The effect of age on the pharmacokinetics and pharmacodynamics of insulin human injection has not been studied.

Elderly patients using MYXREDLIN, may be at increased risk of hypoglycemia due to co-morbid disease [see Warnings and Precautions (5.2)].

## **8.6 Renal Impairment**

The effect of renal impairment on the pharmacokinetics and pharmacodynamics of MYXREDLIN has not been studied. Patients with renal impairment are at increased risk of hypoglycemia and may require more frequent MYXREDLIN dose adjustment and more frequent blood glucose monitoring [see Warnings and Precautions (5.2)].

## **8.7 Hepatic Impairment**

The effect of hepatic impairment on the pharmacokinetics and pharmacodynamics of MYXREDLIN has not been studied. Patients with hepatic impairment are at increased risk of hypoglycemia and may require more frequent MYXREDLIN dose adjustment and more frequent blood glucose monitoring [see Warnings and Precautions (5.2)].


## 10 OVERDOSAGE

Excess insulin administration may cause hypoglycemia and hypokalemia. More severe episodes with coma, seizure, or neurologic impairment can be treated with intramuscular or subcutaneous glucagon or intravenous glucose. Sustained monitoring may be necessary because hypoglycemia may recur after apparent clinical recovery. Hypokalemia must be corrected appropriately. [see Warnings and Precautions (5.2, 5.4)]

## 11 DESCRIPTION

Insulin human is a short-acting human insulin. It is a polypeptide hormone and is produced by recombinant DNA technology, utilizing *Pichia pastoris* (a yeast) as the production organism. Insulin human is regular human insulin and has the empirical formula  $C_{257}H_{383}N_{65}O_{77}S_6$  and a molecular weight of 5808.

Figure 1: Structural formula of Insulin Human



MYXREDLIN (insulin human) in 0.9% sodium chloride injection for intravenous use is a sterile, preservative-free, nonpyrogenic, clear, aqueous, and colorless solution supplied



in a 100 mL GALAXY single-dose container. MYXREDLIN contains 100 units of insulin human in 100 milliliters of 0.9% sodium chloride injection. Each milliliter of solution contains 1 unit Insulin Human, USP; 0.412 mg Dibasic Sodium Phosphate Anhydrous, USP; 0.29 mg Monobasic Sodium Phosphate Monohydrate, USP; 9 mg Sodium Chloride, USP; and Water for Injection, USP. The pH range is 6.5-7.2.

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

The primary activity of insulin, including MYXREDLIN is the regulation of glucose metabolism. Insulins lower blood glucose by stimulating peripheral glucose uptake, especially by skeletal muscle and fat, and by inhibiting hepatic glucose production. Insulin inhibits lipolysis and proteolysis, and enhances protein synthesis.

### **12.2 Pharmacodynamics**

MYXREDLIN is a short-acting insulin. The time course of insulin action (i.e., glucose lowering) may vary considerably in different individuals, within the same individual, and different doses. In a double-blind, randomized, crossover, euglycemic glucose clamp study described below, the average onset of action, defined as start of intravenous glucose infusion during the clamp, was approximately 21 minutes after starting of the intravenous infusion administration. The glucose infusion rate gradually increased to a maximum response of 13.7 mg/kg/min after 5 hours of human insulin infusion.

### **12.3 Pharmacokinetics**

In a double-blind, randomized, crossover, euglycemic glucose clamp study, fifty eight healthy male subjects between 19 and 50 years of age received an intravenous infusion of either MYXREDLIN or another human insulin at 1 milliUnit/kg/min for 6 hours (0.36 units/kg total dose). On average, insulin concentrations of about 300 pM were attained approximately from 1.5 hours to 6 hours after starting intravenous infusion of MYXREDLIN, followed by a return to the baseline level 1.5 hours after stopping intravenous infusion.

#### Metabolism and Elimination

The mean terminal half-life from concentration data after stopping the intravenous infusion was estimated to be 23.4 minutes in healthy volunteers.

#### Specific Population

The effects of gender, age, obesity, renal and hepatic impairment on the pharmacodynamics and pharmacokinetics of insulin human injection have not been studied.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

Standard 2-year carcinogenicity studies in animals have not been performed to evaluate the carcinogenic potential of insulin human injection.

Human insulin is not mutagenic in the following *in vitro* tests: The chromosomal aberration assay in human lymphocytes, the micronucleus assay in mouse polychromatic erythrocytes, and the mutation frequency assay in Chinese hamster cells. Standard reproduction and teratology studies in animals, including fertility assessments have not been conducted with insulin human injection.

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

### **16.1 How Supplied**

MYXREDLIN (insulin human) in 0.9% sodium chloride injection contains 100 units/100 mL (1 unit/mL) of insulin human in 0.9% sodium chloride and is a clear, colorless solution available as:

100 mL single-dose GALAXY container, package of 12, NDC 0338-0126-12

### **16.2 Storage and Handling**

Store MYXREDLIN in the refrigerator (36°F to 46°F [2°C to 8°C]) in the original carton to protect from light. Do not use after the expiration date printed on the carton and container label.

If needed, MYXREDLIN may be removed from the original carton and stored at room temperature up to 77°F (25°C) for up to 30 days. Once stored at room temperature, do not place back in the refrigerator. Discard MYXREDLIN after 30 days if stored at room temperature.

Do not freeze and do not use MYXREDLIN if it has been frozen. Do not shake.

## **17 PATIENT COUNSELING INFORMATION**

### **Hypoglycemia**

Inform patients that hypoglycemia is the most common adverse reaction with insulin. Inform patients that their ability to concentrate and react may be impaired as a result of hypoglycemia.

### **Hypersensitivity Reactions**

Advise patients that hypersensitivity reactions have occurred with insulin human injection [see Warnings and Precautions (5.3)].

Manufactured by:

**Baxter Healthcare Corporation**  
Deerfield, IL 60015 USA

U.S. License Number 0140

Baxter, Galaxy and Myxredlin are registered trademarks of Baxter International Inc.  
07-19-01-720

## **PACKAGE/LABEL PRINCIPAL DISPLAY PANEL**

NDC 0338-0126-12

**MYXREDLIN**

**Insulin Human**  
in 0.9% Sodium Chloride Injection

**100 units per 100 mL**  
(1 unit / mL)

100 mL Single-Dose GALAXY container (REGULAR) Insulin Human  
Discard unused portion Sterile Nonpyrogenic

**For Intravenous Infusion Only**

Cautions: Do not add supplemental medication or additives.  
Must not be used in series connections. Check for minute leaks and solution clarity.

Each mL contains: 1 unit Insulin Human, USP; 0.412 mg Dibasic Sodium Phosphate Anhydrous, USP; 0.29 mg Monobasic Sodium Phosphate Monohydrate, USP; 9 mg Sodium Chloride, USP; and Water for Injection, USP.

Dosage: See prescribing information.

**Rx only**

**Store refrigerated at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. Do Not Shake. Do Not Freeze.**

If needed, may store at room temperature up to 77°F (25°C) up to 30 days. Discard after 30 days if stored at room temperature.

**Baxter**

**Code 2G3322**

PL 2501 Plastic

Baxter, Galaxy and Myxredlin are registered trademarks of  
Baxter International Inc.

**Baxter Healthcare Corporation**


Deerfield, IL 60015 USA

U.S. License no. 0140

Product of USA

07-34-00-0510

**CONTAINS INSULIN**



**NDC 0338-0126-12**

**MYXREDLIN**  
**Insulin Human**  
**in 0.9% Sodium Chloride Injection**

**100 units per 100 mL**  
**(1 unit / mL)**

**100 mL Single-Dose GALAXY container**

**Discard unused portion**  
**For Intravenous Infusion Only**

**(REGULAR) Insulin Human**  
**Sterile Nonpyrogenic**

Cautions: Do not add supplemental medication or additives.

Must not be used in series connections. Check for minute leaks and solution clarity.

Each mL contains: 1 unit Insulin Human, USP; 0.412 mg Dibasic Sodium Phosphate Anhydrous, USP; 0.29 mg Monobasic Sodium Phosphate Monohydrate, USP; 9 mg Sodium Chloride, USP; and Water for Injection, USP.

Dosage: See prescribing information.

**Rx only**

**Store refrigerated at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. Do Not Shake. Do Not Freeze.**

If needed, may store at room temperature up to 77°F (25°C) up to 30 days. Discard after 30 days if stored at room temperature.

**Baxter logo**

Baxter, Galaxy and Myxredlin are trademarks of Baxter International Inc.

**Baxter Healthcare Corporation**

Deerfield, IL 60015 USA  
U.S. License no. 0140

**Code 2G3322**

PL 2501 Plastic

Product of USA  
07-34-00-0510



**FROM LIGHT WHILE REFRIGERATED**

Rx Only

1 Single-Dose Galaxy container  
Discard unused portion

***Baxter Logo***

**For Intravenous Infusion Only**

**MYXREDLIN**

**Insulin Human**

**in 0.9% Sodium Chloride Injection**

**CONTAINS INSULIN HUMAN  
100 units per 100 mL**

(1 unit/mL)

**NDC 0338-0126-12**

**MYXREDLIN**

**Insulin Human**

**in 0.9% Sodium Chloride Injection**

(1 unit/mL)

UNVARNISHED AREA FOR ON-LINE  
PRINTING OF LOT AND EXP

UPC-A  
Bar Code  
Placement

Sterile, Nonpyrogenic

Each mL contains: 1 unit Insulin Human, USP; 0.412 mg Dibasic Sodium Phosphate Anhydrous,

USP; 0.29 mg Monobasic Sodium Phosphate Monohydrate, USP; 9 mg Sodium Chloride, USP;

and Water for Injection, USP. No preservative.

**Dosage:** For Intravenous Infusion Only. See prescribing information.

**Caution:** Do not add supplemental medication or additives.

**Rx only**

**Store refrigerated at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light.**

**Do Not Shake. Do Not Freeze.**

If needed, may store MYXREDLIN at room temperature up to 77°F (25°C) up to 30 days. Once stored at room temperature, do not place back in the refrigerator.

Discard after 30 days if stored at room temperature.

**Discard 30 days after storing at room temperature. Discard After:  
(Month) (Day) (Year)**

**(to be completed by dispensing pharmacist)**

Baxter, Galaxy and Myxredlin are registered trademarks of Baxter International Inc.

**Baxter Healthcare Corporation**

Deerfield, IL 60015 USA

U.S. License no. 0140

Product of USA

**07-01-00-0251**

**MYXREDLIN**  
Insulin Human  
in 0.9% Sodium Chloride Injection  
100 units per 100 mL  
(1 unit/mL)

EXP: JUN 21      LOT #: NC125840



(01)50303380126120(17)210630(21)100000002582(10)NC125840

Contains: 12 x 100 mL Single-Dose GALAXY Containers  
NDC 0338-0126-12  
Code 2G3322

Store refrigerated at 36°F to 46°F (2°C to 8°C). Do Not Shake. Do Not Freeze.  
If needed, may store at room temperature up to 77°F (25°C) up to 30 days.  
Once stored at room temperature, do not place back in the refrigerator.  
Discard after 30 days if stored at room temperature.



GTIN 50303380126120  
SERIAL 100000002582  
EXP 210630  
LOT NC125840

Rx Only  
**Baxter**  
Baxter Healthcare Corporation  
Deerfield, IL 60015 USA  
07-06-00-0789

**MYXREDLIN**  
**Insulin Human**  
**in 0.9% Sodium Chloride Injection**  
**100 units per 100 mL**  
(1 unit/mL)

EXP: JUN 21

LOT #: NC125840

**Bar Code**

(01)50803380126120(17)210630(21)100000002582(10)NC125840

**Contains: 12 x 100 mL Single-Dose GALAXY Containers**  
**NDC 0338-0126-12**

Code 2G3322

Store refrigerated at 36°F to 46°F (2°C to 8°C) Do Not Shake. Do Not Freeze.

If needed, may store at room temperature up to 77°F (25°C) up to 30 days.

Once stored at room temperature, do not place back in the refrigerator.

Discard after 30 days if stored at room temperature.

**Bar Code**

GTIN 50303380126120

SERIAL 100000002582

EXP 210630



LOT NC125840

**Rx Only**

**Baxter Logo**

**Baxter Healthcare Corporation**

Deerfield, IL 60015 USA

07-06-00-0789

G

## MYXREDLIN

insulin human injection, solution

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0338-0126
<b>Route of Administration</b>	INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>INSULIN HUMAN</b> (UNII: 1Y17CTI5SR) (INSULIN HUMAN - UNII:1Y17CTI5SR)	INSULIN HUMAN	1.00 [iU] in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM PHOSPHATE, DIBASIC, ANHYDROUS</b> (UNII: 22ADO53M6F)	0.412 mg in 1 mL
<b>SODIUM PHOSPHATE, MONOBASIC, MONOHYDRATE</b> (UNII: 593YOG76RN)	0.29 mg in 1 mL
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	9 mg in 1 mL
<b>WATER</b> (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0126-12	12 in 1 CARTON	07/12/2019	
1		100 mL in 1 CONTAINER; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA208157	07/12/2019	

**Labeler** - Baxter Healthcare Corporation (005083209)

## Establishment

Name	Address	ID/FEI	Business Operations
Baxter Healthcare Corporation		194684502	ANALYSIS(0338-0126) , MANUFACTURE(0338-0126) , LABEL(0338-0126) , PACK(0338-0126) , STERILIZE(0338-0126)

Revised: 6/2020

Baxter Healthcare Corporation