

DULOXETINE HCL DR- duloxetine hcl capsule, delayed release
Advanced Rx Pharmacy of Tennessee, LLC

Duloxetine HCL DR 30mg Capsules

Boxed Warning

WARNING: SUICIDAL THOUGHTS AND BEHAVIORS

Antidepressants increased the risk of suicidal thoughts and behavior in children, adolescents, and young adults in short-term studies. These studies did not show an increase in the risk of suicidal thoughts and behavior with antidepressant use in patients over age 24; there was a reduction in risk with antidepressant use in patients aged 65 and older [see WARNINGS AND PRECAUTIONS (5.1)].

In patients of all ages who are started on antidepressant therapy, monitor closely for worsening, and for emergence of suicidal thoughts and behaviors. Advise families and caregivers of the need for close observation and communication with the prescriber [see WARNINGS AND PRECAUTIONS (5.1)].

1. Indications and Usage Section

1 INDICATIONS AND USAGE

Duloxetine delayed-release capsules are indicated for the treatment of:

Major Depressive Disorder [SEE CLINICAL STUDIES (14.1)]
Generalized Anxiety Disorder [SEE CLINICAL STUDIES (14.2)]
Diabetic Peripheral Neuropathy [SEE CLINICAL STUDIES (14.3)]
Fibromyalgia [see CLINICAL STUDIES (14.4)]
Chronic Musculoskeletal Pain [SEE CLINICAL STUDIES (14.5)]

2. Dosage and Administration Section

2 DOSAGE AND ADMINISTRATION

Swallow duloxetine delayed-release capsules whole. Do not chew or crush. Do not open the capsule and sprinkle its contents on food or mix with liquids. All of these might affect the enteric coating. Duloxetine delayed-release capsules can be given without regard to meals. If a dose of duloxetine delayed-release capsules is missed, take the missed dose as soon as it is remembered. If it is almost time for the next dose, skip the missed dose and take the next dose at the regular time. Do not take two doses of duloxetine delayed-release capsules at the same time.

2.1 Dosage for Treatment of Major Depressive Disorder

Administer duloxetine delayed-release capsules at a total dose of 40 mg/day (given as 20 mg twice daily) to 60 mg/day (given either once daily or as 30 mg twice daily). For some patients, it may be desirable to start at 30 mg once daily for 1 week, to allow patients to adjust to the medication before increasing to 60 mg once daily. While a 120 mg/day

dose was shown to be effective, there is no evidence that doses greater than 60 mg/day confer any additional benefits. The safety of doses above 120 mg/day has not been adequately evaluated. Periodically reassess to determine the need for maintenance treatment and the appropriate dose for such treatment [see CLINICAL STUDIES (14.1)].

2.2 Dosage for Treatment of Generalized Anxiety Disorder

Adults — For most patients, initiate duloxetine delayed-release capsules 60 mg once daily. For some patients, it may be desirable to start at 30 mg once daily for 1 week, to allow patients to adjust to the medication before increasing to 60 mg once daily. While a 120 mg once daily dose was shown to be effective, there is no evidence that doses greater than 60 mg/day confer additional benefit. Nevertheless, if a decision is made to increase the dose beyond 60 mg once daily, increase dose in increments of 30 mg once daily. The safety of doses above 120 mg once daily has not been adequately evaluated. Periodically reassess to determine the continued need for maintenance treatment and the appropriate dose for such treatment [see CLINICAL STUDIES (14.2)].

Elderly — Initiate duloxetine delayed-release capsules at a dose of 30 mg once daily for 2 weeks before considering an increase to the target dose of 60 mg. Thereafter, patients may benefit from doses above 60 mg once daily. If a decision is made to increase the dose beyond 60 mg once daily, increase dose in increments of 30 mg once daily. The maximum dose studied was 120 mg per day. Safety of doses above 120 mg once daily has not been adequately evaluated [see CLINICAL STUDIES (14.2)].

Children and Adolescents (7 to 17 years of age) — Initiate duloxetine delayed-release capsules at a dose of 30 mg once daily for 2 weeks before considering an increase to 60 mg. The recommended dose range is 30 to 60 mg once daily. Some patients may benefit from doses above 60 mg once daily. If a decision is made to increase the dose beyond 60 mg once daily, increase dose in increments of 30 mg once daily. The maximum dose studied was 120 mg per day. The safety of doses above 120 mg once daily has not been evaluated [see CLINICAL STUDIES (14.2)].

2.3 Dosage for Treatment of Diabetic Peripheral Neuropathic Pain

Administer duloxetine delayed-release capsules 60 mg once daily. There is no evidence that doses higher than 60 mg confer additional significant benefit and the higher dose is clearly less well tolerated [see CLINICAL STUDIES (14.3)]. For patients for whom tolerability is a concern, a lower starting dose may be considered.

Since diabetes is frequently complicated by renal disease, consider a lower starting dose and gradual increase in dose for patients with renal impairment [see DOSAGE AND ADMINISTRATION (2.6), USE IN SPECIFIC POPULATIONS (8.10), and CLINICAL PHARMACOLOGY (12.3)].

2.4 Dosage for Treatment of Fibromyalgia

Administer duloxetine delayed-release capsules 60 mg once daily. Begin treatment at 30 mg once daily for 1 week, to allow patients to adjust to the medication before increasing to 60 mg once daily. Some patients may respond to the starting dose. There is no evidence that doses greater than 60 mg/day confer additional benefit, even in patients who do not respond to a 60 mg dose, and higher doses are associated with a higher rate of adverse reactions [see CLINICAL STUDIES (14.4)].

2.5 Dosage for Treatment of Chronic Musculoskeletal Pain

Administer duloxetine delayed-release capsules 60 mg once daily. Begin treatment at 30 mg for one week, to allow patients to adjust to the medication before increasing to 60 mg once daily. There is no evidence that higher doses confer additional benefit, even in patients who do not respond to a 60 mg dose, and higher doses are associated with a higher rate of adverse reactions [see CLINICAL STUDIES (14.5)].

2.6 Dosing in Special Populations

Hepatic Impairment — Avoid use in patients with chronic liver disease or cirrhosis [see WARNINGS AND PRECAUTIONS (5.14) and USE IN SPECIFIC POPULATIONS (8.9)].

Severe Renal Impairment — Avoid use in patients with severe renal impairment, GFR <30 mL/min [see WARNINGS AND PRECAUTIONS (5.14) and USE IN SPECIFIC POPULATIONS (8.10)].

2.7 Discontinuing Duloxetine Delayed-Release Capsules

Adverse reactions after discontinuation of duloxetine delayed-release capsules, after abrupt or tapered discontinuation, include: dizziness, headache, nausea, diarrhea, paresthesia, irritability, vomiting, insomnia, anxiety, hyperhidrosis, and fatigue. A gradual reduction in dosage rather than abrupt cessation is recommended whenever possible [see WARNINGS AND PRECAUTIONS (5.7)].

2.8 Switching a Patient to or from a Monoamine Oxidase Inhibitor (MAOI) Intended to Treat Psychiatric Disorders

At least 14 days should elapse between discontinuation of an MAOI intended to treat psychiatric disorders and initiation of therapy with duloxetine delayed-release capsules. Conversely, at least 5 days should be allowed after stopping duloxetine delayed-release capsules before starting an MAOI intended to treat psychiatric disorders [see CONTRAINDICATIONS (4)].

2.9 Use of Duloxetine Delayed-Release Capsules with Other MAOIs such as Linezolid or Methylene Blue

Do not start duloxetine delayed-release capsules in a patient who is being treated with linezolid or intravenous methylene blue because there is an increased risk of serotonin syndrome. In a patient who requires more urgent treatment of a psychiatric condition, other interventions, including hospitalization, should be considered [see CONTRAINDICATIONS (4)].

In some cases, a patient already receiving duloxetine delayed-release capsules therapy may require urgent treatment with linezolid or intravenous methylene blue. If acceptable alternatives to linezolid or intravenous methylene blue treatment are not available and the potential benefits of linezolid or intravenous methylene blue treatment are judged to outweigh the risks of serotonin syndrome in a particular patient, duloxetine delayed-release capsules should be stopped promptly, and linezolid or intravenous methylene blue can be administered. The patient should be monitored for symptoms of serotonin syndrome for 5 days or until 24 hours after the last dose of linezolid or intravenous methylene blue, whichever comes first. Therapy with duloxetine delayed-release capsules may be resumed 24 hours after the last dose of linezolid or intravenous methylene blue [see WARNINGS AND PRECAUTIONS (5.4)].

The risk of administering methylene blue by non-intravenous routes (such as oral tablets or by local injection) or in intravenous doses much lower than 1 mg/kg with duloxetine

delayed-release capsules is unclear. The clinician should, nevertheless, be aware of the possibility of emergent symptoms of serotonin syndrome with such use [see WARNINGS AND PRECAUTIONS (5.4)].

3. Dosage Forms and Strengths

Duloxetine hydrochloride is available as delayed-release capsules:

20 mg is green opaque/green opaque, size '4' hard gelatin capsule filled with white to off-white pellets and imprinted with 'X' on green opaque cap and '01' on green opaque body with black ink.

30 mg is blue opaque/white opaque, size '3' hard gelatin capsule filled with white to off-white pellets and imprinted with 'X' on blue opaque cap and '02' on white opaque body with black ink.

60 mg is blue opaque/green opaque, size '1' hard gelatin capsule filled with white to off-white pellets and imprinted with 'X' on blue opaque cap and '03' on green opaque body with black ink.

4. Contraindications

The use of MAOIs intended to treat psychiatric disorders with duloxetine delayed-release capsules or within 5 days of stopping treatment with duloxetine delayed-release capsules is contraindicated because of an increased risk of serotonin syndrome. The use of duloxetine delayed-release capsules within 14 days of stopping an MAOI intended to treat psychiatric disorders is contraindicated [see DOSAGE AND ADMINISTRATION (2.8) and WARNINGS AND PRECAUTIONS (5.4)].

Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome [see DOSAGE AND ADMINISTRATION (2.9) and WARNINGS AND PRECAUTIONS (5.4)].

5. Warnings and Precautions

5.1 Suicidal Thoughts and Behaviors in Children, Adolescents, and Young Adults

Patients with major depressive disorder (MDD), both adult and pediatric, may experience worsening of their depression and/or the emergence of suicidal ideation and behavior (suicidality) or unusual changes in behavior, whether or not they are taking antidepressant medications, and this risk may persist until significant remission occurs. Suicide is a known risk of depression and certain other psychiatric disorders, and these disorders themselves are the strongest predictors of suicide. There has been a long-standing concern, however, that antidepressants may have a role in inducing worsening of depression and the emergence of suicidality in certain patients during the early phases of treatment.

Pooled analyses of short-term placebo-controlled trials of antidepressant drugs (SSRIs and others) showed that these drugs increase the risk of suicidal thinking and behavior

(suicidality) in children, adolescents, and young adults (ages 18 to 24) with major depressive disorder (MDD) and other psychiatric disorders. Short-term studies did not show an increase in the risk of suicidality with antidepressants compared to placebo in adults beyond age 24; there was a reduction with antidepressants compared to placebo in adults aged 65 and older.

The pooled analyses of placebo-controlled trials in children and adolescents with MDD, obsessive compulsive disorder (OCD), or other psychiatric disorders included a total of 24 short-term trials of 9 antidepressant drugs in over 4400 patients. The pooled analyses of placebo-controlled trials in adults with MDD or other psychiatric disorders included a total of 295 short-term trials (median duration of 2 months) of 11 antidepressant drugs in over 77,000 patients. There was considerable variation in risk of suicidality among drugs, but a tendency toward an increase in the younger patients for almost all drugs studied. There were differences in absolute risk of suicidality across the different indications, with the highest incidence in MDD. The risk of differences (drug vs placebo), however, were relatively stable within age strata and across indications. These risk differences (drug-placebo difference in the number of cases of suicidality per 1000 patients treated) are provided in Table 1.

Table 1

Age Range	Drug-Placebo Difference in Number of Cases of Suicidality per 1000 Patients Treated
<18	Increases Compared to Placebo 14 additional cases
18 to 24	5 additional cases
25 to 64	Decreases Compared to Placebo 1 fewer case
≥65	6 fewer cases

No suicides occurred in any of the pediatric duloxetine delayed-release capsule trials. There were suicides in the adult duloxetine delayed-release capsule trials, but the number was not sufficient to reach any conclusion about duloxetine delayed-release capsules effect on suicide.

It is unknown whether the suicidality risk extends to longer-term use, i.e., beyond several months. However, there is substantial evidence from placebo-controlled maintenance trials in adults with depression that the use of antidepressants can delay the recurrence of depression.

All patients being treated with antidepressants for any indication should be monitored appropriately and observed closely for clinical worsening, suicidality, and unusual changes in behavior, especially during the initial few months of a course of drug therapy, or at times of dose changes, either increases or decreases.

The following symptoms, anxiety, agitation, panic attacks, insomnia, irritability, hostility, aggressiveness, impulsivity, akathisia (psychomotor restlessness), hypomania, and mania, have been reported in adult and pediatric patients being treated with

antidepressants for major depressive disorder as well as for other indications, both psychiatric and nonpsychiatric. Although a causal link between the emergence of such symptoms and either the worsening of depression and/or the emergence of suicidal impulses has not been established, there is concern that such symptoms may represent precursors to emerging suicidality.

Consideration should be given to changing the therapeutic regimen, including possibly discontinuing the medication, in patients whose depression is persistently worse, or who are experiencing emergent suicidality or symptoms that might be precursors to worsening depression or suicidality, especially if these symptoms are severe, abrupt in onset, or were not part of the patient's presenting symptoms.

If the decision has been made to discontinue treatment, medication should be tapered, as rapidly as is feasible, but with recognition that discontinuation can be associated with certain symptoms [see DOSAGE AND ADMINISTRATION (2.8) and WARNINGS AND PRECAUTIONS (5.7)] for descriptions of the risks of discontinuation of duloxetine delayed-release capsules.

Families and caregivers of patients being treated with antidepressants for major depressive disorder or other indications, both psychiatric and nonpsychiatric, should be alerted about the need to monitor patients for the emergence of agitation, irritability, unusual changes in behavior, and the other symptoms described above, as well as the emergence of suicidality, and to report such symptoms immediately to health care providers. Such monitoring should include daily observation by families and caregivers. Prescriptions for duloxetine delayed-release capsules should be written for the smallest quantity of capsules consistent with good patient management, in order to reduce the risk of overdose.

Screening Patients for Bipolar Disorder

A major depressive episode may be the initial presentation of bipolar disorder. It is generally believed (though not established in controlled trials) that treating such an episode with an antidepressant alone may increase the likelihood of precipitation of a mixed/manic episode in patients at risk for bipolar disorder. Whether any of the symptoms described above represent such a conversion is unknown. However, prior to initiating treatment with an antidepressant, patients with depressive symptoms should be adequately screened to determine if they are at risk for bipolar disorder; such screening should include a detailed psychiatric history, including a family history of suicide, bipolar disorder, and depression. It should be noted that duloxetine delayed-release capsules are not approved for use in treating bipolar depression.

5.2 Hepatotoxicity

There have been reports of hepatic failure, sometimes fatal, in patients treated with duloxetine delayed-release capsules. These cases have presented as hepatitis with abdominal pain, hepatomegaly, and elevation of transaminase levels to more than twenty times the upper limit of normal (ULN) with or without jaundice, reflecting a mixed or hepatocellular pattern of liver injury. Duloxetine delayed-release capsules should be discontinued in patients who develop jaundice or other evidence of clinically significant liver dysfunction and should not be resumed unless another cause can be established.

Cases of cholestatic jaundice with minimal elevation of transaminase levels have also been reported. Other postmarketing reports indicate that elevated transaminases, bilirubin, and alkaline phosphatase have occurred in patients with chronic liver disease or cirrhosis.

Duloxetine delayed-release capsules increased the risk of elevation of serum transaminase levels in development program clinical trials. Liver transaminase elevations resulted in the discontinuation of 0.3% (92/34,756) of duloxetine delayed-release capsules-treated patients. In most patients, the median time to detection of the transaminase elevation was about two months. In adult placebo-controlled trials, for patients with normal and abnormal baseline ALT values, elevation of ALT >3 times the ULN occurred in 1.25% (144/11,496) of duloxetine delayed-release capsules-treated patients compared to 0.45% (39/8716) of placebo-treated patients. In adult placebo-controlled studies using a fixed dose design, there was evidence of a duloxetine delayed-release capsules dose response relationship for ALT and AST elevation of >3 times the ULN and >5 times the ULN, respectively.

Because it is possible that duloxetine delayed-release capsules and alcohol may interact to cause liver injury or that duloxetine delayed-release capsules may aggravate pre-existing liver disease, duloxetine delayed-release capsules should not be prescribed to patients with substantial alcohol use or evidence of chronic liver disease.

5.3 Orthostatic Hypotension, Falls and Syncope

Orthostatic hypotension, falls, and syncope have been reported in patients treated with the recommended duloxetine delayed-release capsules dosages. Syncope and orthostatic hypotension tend to occur within the first week of therapy but can occur at any time during duloxetine delayed-release capsules treatment, particularly after dose increases. The risk of falling appears to be related to the degree of orthostatic decrease in blood pressure (BP) as well as other factors that may increase the underlying risk of falls.

In an analysis of patients from all placebo-controlled trials, patients treated with duloxetine delayed-release capsules reported a higher rate of falls compared to patients treated with placebo. Risk appears to be related to the presence of orthostatic decrease in BP. The risk of BP decreases may be greater in patients taking concomitant medications that induce orthostatic hypotension (such as antihypertensives) or are potent CYP1A2 inhibitors [see WARNINGS AND PRECAUTIONS (5.12) and DRUG INTERACTIONS (7.1)] and in patients taking duloxetine delayed-release capsules at doses above 60 mg daily. Consideration should be given to dose reduction or discontinuation of duloxetine delayed-release capsules in patients who experience symptomatic orthostatic hypotension, falls and/or syncope during duloxetine delayed-release capsules therapy.

Risk of falling also appeared to be proportional to a patient's underlying risk for falls and appeared to increase steadily with age. As geriatric patients tend to have a higher underlying risk for falls due to a higher prevalence of risk factors such as use of multiple medications, medical comorbidities and gait disturbances, the impact of increasing age by itself is unclear. Falls with serious consequences including fractures and hospitalizations have been reported with duloxetine delayed-release capsules use [see ADVERSE REACTIONS (6.1)].

5.4 Serotonin Syndrome

The development of a potentially life-threatening serotonin syndrome has been reported with SNRIs and SSRIs, including duloxetine delayed-release capsules, alone but particularly with concomitant use of other serotonergic drugs (including triptans, tricyclic antidepressants, fentanyl, lithium, tramadol, tryptophan, buspirone, amphetamines, and St. John's Wort) and with drugs that impair metabolism of serotonin (in particular, MAOIs, both those intended to treat psychiatric disorders and also others, such as linezolid and intravenous methylene blue).

Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, delirium, and coma), autonomic instability (e.g., tachycardia, labile blood pressure, dizziness, diaphoresis, flushing, hyperthermia), neuromuscular symptoms (e.g., tremor, rigidity, myoclonus, hyperreflexia, incoordination), seizures, and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea). Patients should be monitored for the emergence of serotonin syndrome.

The concomitant use of duloxetine delayed-release capsules with MAOI antidepressants is contraindicated. Duloxetine delayed-release capsules should also not be started in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue. All reports with methylene blue that provided information on the route of administration involved intravenous administration in the dose range of 1 mg/kg to 8 mg/kg. No reports involved the administration of methylene blue by other routes (such as oral tablets or local tissue injection) or at lower doses. There may be circumstances when it is necessary to initiate treatment with an MAOI such as linezolid or intravenous methylene blue in a patient taking duloxetine delayed-release capsules. Duloxetine delayed-release capsules should be discontinued before initiating treatment with the MAOI [see DOSAGE AND ADMINISTRATION (2.9, 2.10) and CONTRAINDICATIONS (4)].

If concomitant use of duloxetine delayed-release capsules with other serotonergic drugs including triptans, tricyclic antidepressants, fentanyl, lithium, tramadol, buspirone, tryptophan, amphetamines, and St. John's Wort is clinically warranted, patients should be made aware of a potential increased risk for serotonin syndrome, particularly during treatment initiation and dose increases. Treatment with duloxetine delayed-release capsules and any concomitant serotonergic agents, should be discontinued immediately if the above events occur and supportive symptomatic treatment should be initiated.

5.5 Increased Risk of Bleeding

Drugs that interfere with serotonin reuptake inhibition, including duloxetine delayed-release capsules, may increase the risk of bleeding events. Case reports and epidemiological studies (case-control and cohort design) have demonstrated an association between use of drugs that interfere with serotonin reuptake and the occurrence of gastrointestinal bleeding. A post-marketing study showed a higher incidence of postpartum hemorrhage in mothers taking duloxetine delayed-release capsules. Other bleeding events related to SSRI and SNRI use have ranged from ecchymoses, hematomas, epistaxis, and petechiae to life-threatening hemorrhages. Concomitant use of aspirin, nonsteroidal anti-inflammatory drugs (NSAIDs), warfarin, and other anti-coagulants may add to this risk.

Inform patients about the risk of bleeding associated with the concomitant use of duloxetine delayed-release capsules and NSAIDs, aspirin, or other drugs that affect coagulation [see DRUG INTERACTIONS (7.4)].

5.6 Severe Skin Reactions

Severe skin reactions, including erythema multiforme and Stevens-Johnson syndrome (SJS), can occur with duloxetine delayed-release capsules. The reporting rate of SJS associated with duloxetine delayed-release capsules use exceeds the general population background incidence rate for this serious skin reaction (1 to 2 cases per million person years). The reporting rate is generally accepted to be an underestimate due to underreporting.

Duloxetine delayed-release capsules should be discontinued at the first appearance of blisters, peeling rash, mucosal erosions, or any other sign of hypersensitivity if no other etiology can be identified.

5.7 Discontinuation Syndrome

Discontinuation symptoms have been systematically evaluated in patients taking duloxetine delayed-release capsules. Following abrupt or tapered discontinuation in adult placebo-controlled clinical trials, the following symptoms occurred at 1% or greater and at a significantly higher rate in duloxetine delayed-release capsules-treated patients compared to those discontinuing from placebo: dizziness, headache, nausea, diarrhea, paresthesia, irritability, vomiting, insomnia, anxiety, hyperhidrosis, and fatigue.

During marketing of other SSRIs and SNRIs (serotonin and norepinephrine reuptake inhibitors), there have been spontaneous reports of adverse events occurring upon discontinuation of these drugs, particularly when abrupt, including the following: dysphoric mood, irritability, agitation, dizziness, sensory disturbances (e.g., paresthesias such as electric shock sensations), anxiety, confusion, headache, lethargy, emotional lability, insomnia, hypomania, tinnitus, and seizures. Although these events are generally self-limiting, some have been reported to be severe.

Patients should be monitored for these symptoms when discontinuing treatment with duloxetine delayed-release capsules. A gradual reduction in the dose rather than abrupt cessation is recommended whenever possible. If intolerable symptoms occur following a decrease in the dose or upon discontinuation of treatment, then resuming the previously prescribed dose may be considered. Subsequently, the healthcare provider may continue decreasing the dose but at a more gradual rate [see DOSAGE AND ADMINISTRATION (2.8)].

5.8 Activation of Mania/Hypomania

In adult placebo-controlled trials in patients with MDD, activation of mania or hypomania was reported in 0.1% (4/3779) of duloxetine delayed-release capsules-treated patients and 0.04% (1/2536) of placebo-treated patients. No activation of mania or hypomania was reported in DPNP, GAD, fibromyalgia, or chronic musculoskeletal pain placebo-controlled trials. Activation of mania or hypomania has been reported in a small proportion of patients with mood disorders who were treated with other marketed drugs effective in the treatment of major depressive disorder. As with these other agents, duloxetine delayed-release capsules should be used cautiously in patients with a history of mania.

5.9 Angle-Closure Glaucoma

The pupillary dilation that occurs following use of many antidepressant drugs including duloxetine delayed-release capsules may trigger an angle closure attack in a patient with anatomically narrow angles who does not have a patent iridectomy.

5.10 Seizures

Duloxetine delayed-release capsules have not been systematically evaluated in patients with a seizure disorder, and such patients were excluded from clinical studies. In adult placebo-controlled clinical trials, seizures/convulsions occurred in 0.02% (3/12,722) of patients treated with duloxetine delayed-release capsules and 0.01% (1/9513) of patients treated with placebo. Duloxetine delayed-release capsules should be prescribed with care in patients with a history of a seizure disorder.

5.11 Increases in Blood Pressure

In adult placebo-controlled clinical trials across the approved adult populations from baseline to endpoint, duloxetine delayed-release capsules treatment was associated with mean increases of 0.5 mm Hg in systolic blood pressure and 0.8 mm Hg in diastolic blood pressure compared to mean decreases of 0.6 mm Hg systolic and 0.3 mm Hg diastolic in placebo-treated patients. There was no significant difference in the frequency of sustained (3 consecutive visits) elevated blood pressure. In a clinical pharmacology study designed to evaluate the effects of duloxetine delayed-release capsules on various parameters, including blood pressure at supratherapeutic doses with an accelerated dose titration, there was evidence of increases in supine blood pressure at doses up to 200 mg twice daily (approximately 3.3 times the maximum recommended dosage). At the highest 200 mg twice daily dose, the increase in mean pulse rate was 5 to 6.8 beats and increases in mean blood pressure were 4.7 to 6.8 mm Hg (systolic) and 4.5 to 7 mm Hg (diastolic) up to 12 hours after dosing.

Blood pressure should be measured prior to initiating treatment and periodically measured throughout treatment [see ADVERSE REACTIONS (6.1)].

5.12 Clinically Important Drug Interactions

Both CYP1A2 and CYP2D6 are responsible for duloxetine delayed-release capsules metabolism.

Potential for Other Drugs to Affect Duloxetine Delayed-Release Capsules

CYP1A2 Inhibitors — Co-administration of duloxetine delayed-release capsules with potent CYP1A2 inhibitors should be avoided [see DRUG INTERACTIONS (7.1)].

CYP2D6 Inhibitors — Because CYP2D6 is involved in duloxetine delayed-release capsules metabolism, concomitant use of duloxetine delayed-release capsules with potent inhibitors of CYP2D6 would be expected to, and does, result in higher concentrations (on average of 60%) of duloxetine delayed-release capsules [see DRUG INTERACTIONS (7.2)].

Potential for Duloxetine Delayed-Release Capsules to Affect Other Drugs

Drugs Metabolized by CYP2D6 — Co-administration of duloxetine delayed-release capsules with drugs that are extensively metabolized by CYP2D6 and that have a narrow therapeutic index, including certain antidepressants (tricyclic antidepressants [TCAs], such as nortriptyline, amitriptyline, and imipramine), phenothiazines and Type 1C antiarrhythmics (e.g., propafenone, flecainide), should be approached with caution. Plasma TCA concentrations may need to be monitored and the dose of the TCA may need to be reduced if a TCA is co-administered with duloxetine delayed-release capsules.

Because of the risk of serious ventricular arrhythmias and sudden death potentially associated with elevated plasma levels of thioridazine, duloxetine delayed-release capsules and thioridazine should not be co-administered [see DRUG INTERACTIONS (7.9)].

Other Clinically Important Drug Interactions

Alcohol — Use of duloxetine delayed-release capsules concomitantly with heavy alcohol intake may be associated with severe liver injury. For this reason, duloxetine delayed-release capsules should not be prescribed for patients with substantial alcohol use [see WARNINGS AND PRECAUTIONS (5.2) and DRUG INTERACTIONS (7.15)].

CNS Acting Drugs — Given the primary CNS effects of duloxetine delayed-release capsules, it should be used with caution when it is taken in combination with or substituted for other centrally acting drugs, including those with a similar mechanism of action [see WARNINGS AND PRECAUTIONS (5.12) and DRUG INTERACTIONS (7.16)].

5.13 Hyponatremia

Hyponatremia may occur as a result of treatment with SSRIs and SNRIs, including duloxetine delayed-release capsules. In many cases, this hyponatremia appears to be the result of the syndrome of inappropriate antidiuretic hormone secretion (SIADH). Cases with serum sodium lower than 110 mmol/L have been reported with duloxetine delayed-release capsules use and appeared to be reversible when duloxetine delayed-release capsules were discontinued. Geriatric patients may be at greater risk of developing hyponatremia with SSRIs and SNRIs. Also, patients taking diuretics or who are otherwise volume depleted may be at greater risk [see USE IN SPECIFIC POPULATIONS (8.5)]. Discontinuation of duloxetine delayed-release capsules should be considered in patients with symptomatic hyponatremia and appropriate medical intervention should be instituted.

Signs and symptoms of hyponatremia include headache, difficulty concentrating, memory impairment, confusion, weakness, and unsteadiness, which may lead to falls. More severe and/or acute cases have been associated with hallucination, syncope, seizure, coma, respiratory arrest, and death.

5.14 Use in Patients with Concomitant Illness

Clinical experience with duloxetine delayed-release capsules in patients with concomitant systemic illnesses is limited. There is no information on the effect that alterations in gastric motility may have on the stability of duloxetine delayed-release capsule's enteric coating. In extremely acidic conditions, duloxetine delayed-release capsules, unprotected by the enteric coating, may undergo hydrolysis to form naphthol. Caution is advised in using duloxetine delayed-release capsules in patients with conditions that may slow gastric emptying (e.g., some diabetics).

Duloxetine delayed-release capsules have not been systematically evaluated in patients with a recent history of myocardial infarction or unstable coronary artery disease. Patients with these diagnoses were generally excluded from clinical studies during the product's premarketing testing.

Hepatic Impairment

Avoid use in patients with chronic liver disease or cirrhosis [see DOSAGE AND

ADMINISTRATION (2.7), WARNINGS AND PRECAUTIONS (5.2), and USE IN SPECIFIC POPULATIONS (8.9)].

Severe Renal Impairment

Avoid use in patients with severe renal impairment, GFR <30 mL/minute. Increased plasma concentration of duloxetine delayed-release capsules, and especially of its metabolites, occurred in patients with end-stage renal disease (requiring dialysis) [see DOSAGE AND ADMINISTRATION (2.7) and USE IN SPECIFIC POPULATIONS (8.10)].

Glycemic Control in Patients with Diabetes

As observed in DPNP trials, duloxetine delayed-release capsules treatment worsened glycemic control in some patients with diabetes. In three clinical trials of duloxetine delayed-release capsules for the management of neuropathic pain associated with diabetic peripheral neuropathy [see Clinical Studies (14.4)], the mean duration of diabetes was approximately 12 years, the mean baseline fasting blood glucose was 176 mg/dL, and the mean baseline hemoglobin A1c (HbA1c) was 7.8%. In the 12-week acute treatment phase of these studies, duloxetine delayed-release capsules were associated with a small increase in mean fasting blood glucose as compared to placebo. In the extension phase of these studies, which lasted up to 52 weeks, mean fasting blood glucose increased by 12 mg/dL in the duloxetine delayed-release capsules group and decreased by 11.5 mg/dL in the routine care group. HbA1c increased by 0.5% in the duloxetine delayed-release capsules group and by 0.2% in the routine care group.

5.15 Urinary Hesitation and Retention

Duloxetine delayed-release capsules are in a class of drugs known to affect urethral resistance. If symptoms of urinary hesitation develop during treatment with duloxetine delayed-release capsules, consideration should be given to the possibility that they might be drug-related.

In post marketing experience, cases of urinary retention have been observed. In some instances of urinary retention associated with duloxetine delayed-release capsules use, hospitalization and/or catheterization has been needed.

6. Adverse Reactions

The following serious adverse reactions are described below and elsewhere in the labeling:

- Suicidal Thoughts and Behaviors in Children, Adolescents, and Young Adults [see BOXED WARNING AND WARNINGS AND PRECAUTIONS (5.1)]
- Hepatotoxicity [see WARNINGS AND PRECAUTIONS (5.2)]
- Orthostatic Hypotension, Falls and Syncope [see WARNINGS AND PRECAUTIONS (5.3)]
- Serotonin Syndrome [see WARNINGS AND PRECAUTIONS (5.4)]
- Increased Risk of Bleeding [see WARNINGS AND PRECAUTIONS (5.5)]
- Severe Skin Reactions [see WARNINGS AND PRECAUTIONS (5.6)]
- Discontinuation Syndrome [see WARNINGS AND PRECAUTIONS (5.7)]
- Activation of Mania/Hypomania [see WARNINGS AND PRECAUTIONS (5.8)]
- Angle-Closure Glaucoma [see WARNINGS AND PRECAUTIONS (5.9)]

- Seizures [see WARNINGS AND PRECAUTIONS (5.10)]
- Increases in Blood Pressure [see WARNINGS AND PRECAUTIONS (5.11)]
- Clinically Important Drug Interactions [see WARNINGS AND PRECAUTIONS (5.12)]
- Hyponatremia [see WARNINGS AND PRECAUTIONS (5.13)]
- Urinary Hesitation and Retention [see WARNINGS AND PRECAUTIONS (5.15)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The stated frequencies of adverse reactions represent the proportion of patients who experienced, at least once, one treatment-emergent adverse reaction of the type listed. A reaction was considered treatment-emergent if it occurred for the first time or worsened while receiving therapy following baseline evaluation.

Adverse Reactions in Adults

Adult Clinical Trial Database

The data described below reflect exposure to duloxetine delayed-release capsules in placebo-controlled adult trials for MDD (N=3779), GAD (N=1018), OA (N=503), CLBP (N=600), DPNP (N=906), and FM (N=1294). The age range in this pooled population was 17 to 89 years of age. In this pooled population, 66%, 61%, 61%, 43%, and 94% of adult patients were female; and 82%, 73%, 85%, 74%, and 86% of adult patients were Caucasian in the MDD, GAD, OA and CLBP, DPNP, and FM populations, respectively. Most patients received duloxetine delayed-release capsules dosages of a total of 60 to 120 mg per day [see CLINICAL STUDIES (14)]. The data below do not include results of the trial that evaluated the efficacy of duloxetine delayed-release capsules for the treatment of GAD in patients ≥ 65 years old (Study GAD-5) [see CLINICAL STUDIES (14.3)]; however, the adverse reactions observed in this geriatric population were generally similar to adverse reactions in the overall adult population.

Adverse Reactions Leading to Treatment Discontinuation in Adult Placebo-Controlled Trials

Major Depressive Disorder

Approximately 8.4% (319/3779) of duloxetine delayed-release capsules-treated patients in placebo-controlled adult trials for MDD discontinued treatment due to an adverse reaction, compared with 4.6% (117/2536) of placebo-treated patients. Nausea (duloxetine delayed-release capsules 1.1%, placebo 0.4%) was the only adverse reaction reported as a reason for discontinuation and considered to be drug-related (i.e., discontinuation occurring in at least 1% of the duloxetine delayed-release capsules-treated patients and at a rate of at least twice that of placebo-treated patients).

Generalized Anxiety Disorder

Approximately 13.7% (139/1018) of the duloxetine delayed-release capsules-treated patients in placebo-controlled adult trials for GAD discontinued treatment due to an adverse reaction, compared with 5% (38/767) for placebo-treated patients. Common adverse reactions reported as a reason for discontinuation and considered to be drug-related (as defined above) included nausea (duloxetine delayed-release capsules 3.3%, placebo 0.4%), and dizziness (duloxetine delayed-release capsules 1.3%, placebo 0.4%).

Diabetic Peripheral Neuropathic Pain

Approximately 12.9% (117/906) of the duloxetine delayed-release capsules-treated patients in placebo-controlled adult trials for DPNP discontinued treatment due to an adverse reaction, compared with 5.1% (23/448) for placebo-treated patients. Common adverse reactions reported as a reason for discontinuation and considered to be drug-related (as defined above) included nausea (duloxetine delayed-release capsules 3.5%, placebo 0.7%), dizziness (duloxetine delayed-release capsules 1.2%, placebo 0.4%), and somnolence (duloxetine delayed-release capsules 1.1%, placebo 0%).

Fibromyalgia

Approximately 17.5% (227/1294) of the duloxetine delayed-release capsules-treated patients in 3- to 6-month placebo-controlled adult trials for FM discontinued treatment due to an adverse reaction, compared with 10.1% (96/955) for placebo-treated patients. Adverse reactions reported as a reason for discontinuation and considered to be drug-related (as defined above) included nausea (duloxetine delayed-release capsules 2.0%, placebo 0.5%), headache (duloxetine delayed-release capsules 1.2%, placebo 0.3%), somnolence (duloxetine delayed-release capsules 1.1%, placebo 0%), and fatigue (duloxetine delayed-release capsules 1.1%, placebo 0.1%).

Chronic Pain due to Osteoarthritis

Approximately 15.7% (79/503) of the duloxetine delayed-release capsules-treated patients in 13-week, placebo-controlled adult trials for chronic pain due to OA discontinued treatment due to an adverse reaction, compared with 7.3% (37/508) for placebo-treated patients. Adverse reactions reported as a reason for discontinuation and considered to be drug-related (as defined above) included nausea (duloxetine delayed-release capsules 2.2%, placebo 1%).

Chronic Low Back Pain

Approximately 16.5% (99/600) of the duloxetine delayed-release capsules-treated patients in 13-week, placebo-controlled adult trials for CLBP discontinued treatment due to an adverse reaction, compared with 6.3% (28/441) for placebo-treated patients. Adverse reactions reported as a reason for discontinuation and considered to be drug-related (as defined above) included nausea (duloxetine delayed-release capsules 3%, placebo 0.7%), and somnolence (duloxetine delayed-release capsules 1%, placebo 0%).

Most Common Adverse Reactions in Adult Trials

The most commonly observed adverse reactions in duloxetine delayed-release capsules-treated patients (as defined above) were:

Diabetic Peripheral Neuropathic Pain: nausea, somnolence, decreased appetite, constipation, hyperhidrosis, and dry mouth.

Fibromyalgia: nausea, dry mouth, constipation, somnolence, decreased appetite, hyperhidrosis, and agitation.

Chronic Pain due to Osteoarthritis: nausea, fatigue, constipation, dry mouth, insomnia, somnolence, and dizziness.

Chronic Low Back Pain: nausea, dry mouth, insomnia, somnolence, constipation, dizziness, and fatigue.

The most commonly observed adverse reactions in duloxetine delayed-release capsules-treated patients in all the pooled adult populations (i.e., MDD, GAD, DPNP, FM, OA, and

CLBP) (incidence of at least 5% and at least twice the incidence in placebo-treated patients) were nausea, dry mouth, somnolence, constipation, decreased appetite, and hyperhidrosis.

Table 2 displays the incidence of adverse reactions in placebo-controlled trials for approved adult populations (i.e., MDD, GAD, DPNP, FM, OA, and CLBP) that occurred in 5% or more of duloxetine delayed-release capsules-treated patients and with an incidence greater than placebo-treated patients.

Table 2: Adverse Reactions: Incidence of 5% or More and Greater than Placebo in Placebo-Controlled Trials of Approved Adult Populations^a

Adverse Reaction	Percentage of Patients Reporting Reaction	
	Duloxetine Delayed-Release Capsules (N=8100)	Placebo (N =5655)
Nausea ^c	23	8
Headache	14	12
Dry mouth	13	5
Somnolence ^e	10	3
Fatigue ^{b,c}	9	5
Insomnia ^d	9	5
Constipation ^c	9	4
Dizziness ^c	9	5
Diarrhea	9	6
Decreased appetite ^c	7	2
Hyperhidrosis ^c	6	1
Abdominal pain ^f	5	4

Table 2: Adverse Reactions: Incidence of 5% or More and Greater than Placebo in Placebo-Controlled Trials of Approved Adult Populations^a

^a Includes adults with MDD, GAD, DPNP, FM, and chronic musculoskeletal pain. The inclusion of an event in the table is determined based on the percentages before rounding; however, the percentages displayed in the table are rounded to the nearest integer.

^b Also includes asthenia.

^c Events for which there was a significant dose-dependent relationship in fixed-dose studies, excluding three MDD studies which did not have a placebo lead-in period or dose titration.

^d Also includes initial insomnia, middle insomnia, and early morning awakening.

^e Also includes hypersomnia and sedation.

^f Also includes abdominal discomfort, abdominal pain lower, abdominal pain upper, abdominal tenderness, and gastrointestinal pain.

Adverse Reactions in Pooled MDD and GAD Trials in Adults

Table 3 displays the incidence of adverse reactions in MDD and GAD placebo-controlled adult trials that occurred in 2% or more of duloxetine delayed-release capsules-treated patients and with an incidence greater than placebo-treated patients.

Table 3: Adverse Reactions: Incidence of 2% or More and Greater than Placebo in MDD and GAD Placebo-Controlled Trials in Adults^{a,b}

System Organ Class / Adverse Reaction	Percentage of Patients Reporting Reaction	
	Duloxetine Delayed-Release Capsules (N=3303)	Placebo (N=2352)
Gastrointestinal Disorders		
Nausea	23	7
Dry Mouth ^b	11	3
Constipation ^b	10	3
Diarrhea	9	5
Abdominal Pain ^c	5	4
Vomiting	3	2
Dyspepsia	2	1
General Disorders and Administration Site Conditions		
Fatigued	11	5
Infections and Infestations		
Nasopharyngitis	4	4
Upper Respiratory Tract Infection	3	3
Influenza	2	2
Metabolism and Nutrition Disorders		
Decreased Appetite ^b	8	1
Musculoskeletal and Connective Tissue Disorders		
Musculoskeletal Pain ^e	3	3
Muscle Spasms	2	2
Nervous System Disorders		
Headache	13	8
Somnolence ^{b,f}	11	3
Dizziness	9	5
Paraesthesia ^g	2	2
Tremor ^b	2	<1
Psychiatric Disorders		
Insomnia ^{b,h}	10	5
Agitation ⁱ	3	1
Reproductive System and Breast Disorders		
Erectile Dysfunction ^b	4	<1
Ejaculation Disorder ^j	2	<1
Respiratory, Thoracic, and Mediastinal Disorders		
Cough	2	2
Skin and Subcutaneous Tissue Disorders		
Hyperhidrosis	6	1
Vascular Disorders		
Flushing ^k	3	1
Blood pressure increased ^l	2	1

a The inclusion of an event in the table is determined based on the percentages before rounding; however, the percentages displayed in the table are rounded to the nearest integer.

b Incidence of 120 mg/day is significantly greater than the incidence for 60 mg/day.

c Includes abdominal discomfort, lower abdominal pain, upper abdominal pain, abdominal tenderness and gastrointestinal pain.

d Includes asthenia.

e Includes myalgia and neck pain.

f Includes hypersomnia and sedation.

g Includes hypoaesthesia, facial hypoaesthesia, genital hypoaesthesia and oral paraesthesia.

h Includes initial insomnia, middle insomnia, and early morning awakening.

i Includes feeling jittery, nervousness, restlessness, tension and psychomotor hyperactivity.

j Includes ejaculation failure.

k Includes hot flush.

l Includes increased diastolic blood pressure, increased systolic blood pressure, diastolic hypertension, essential hypertension, hypertension, hypertensive crisis, labile hypertension, orthostatic hypertension, secondary hypertension, and systolic hypertension.

Effects on Male and Female Sexual Function in Adults with MDD

Changes in sexual desire, sexual performance and sexual satisfaction often occur as manifestations of psychiatric disorders or diabetes, but they may also be a consequence of pharmacologic treatment. Because adverse sexual reactions are presumed to be voluntarily underreported, the Arizona Sexual Experience Scale (ASEX), a validated measure designed to identify sexual adverse reactions, was used prospectively in 4 MDD placebo-controlled adult trials (Studies MDD-1, MDD-2, MDD-3, and MDD-4) [see CLINICAL STUDIES (14.2)]. The ASEX scale includes five questions that pertain to the following aspects of sexual function: 1) sex drive, 2) ease of arousal, 3) ability to achieve erection (men) or lubrication (women), 4) ease of reaching orgasm, and 5) orgasm satisfaction. Positive numbers signify a worsening of sexual function from baseline. Negative numbers signify an improvement from a baseline level of dysfunction, which is commonly seen in depressed patients.

In these trials, duloxetine delayed-release capsules-treated male patients experienced significantly more sexual dysfunction, as measured by the total score on the ASEX and the ability to reach orgasm, than placebo-treated male patients (see Table 5). Duloxetine delayed-release capsules-treated female patients did not experience more sexual dysfunction than placebo-treated female patients as measured by ASEX total score. Healthcare providers should routinely inquire about possible sexual adverse reactions in duloxetine delayed-release capsules-treated patients.

Table 5: Mean Change in ASEX Scores by Gender in MDD Placebo-Controlled Adult Trials

Male Patients ^a		Female Patients ^a	
Duloxetine Delayed-Release Capsules	Placebo (n=87)	Duloxetine Delayed-Release Capsules	Placebo (n=126)

	(n=175)	(n=241)	(n=241)	(n=120)
ASEX Total (Items 1 to 5)	0.56b	-1.07	-1.15	1.07
Item 1 — Sex drive	-0.07	-0.12	-0.32	0.24
Item 2 — Arousal	0.01	-0.26	-0.21	0.18
Item 3 — Ability to achieve erection (men); Lubrication (women)	0.03	-0.25	-0.17	0.18
Item 4 — Ease of reaching orgasm	0.40c	-0.24	-0.09	0.13
Item 5 — Orgasm satisfaction	0.09	-0.13	-0.11	0.17

a n=Number of patients with non-missing change score for ASEX total.

b p=0.013 versus placebo.

c p<0.001 versus placebo

Vital Sign Changes in Adults

In placebo-controlled clinical trials across approved adult populations for change from baseline to endpoint, duloxetine delayed-release capsules-treated patients had mean increases of 0.23 mm Hg in systolic blood pressure (SBP) and 0.73 mm Hg in diastolic blood pressure (DBP) compared to mean decreases of 1.09 mm Hg in SBP and 0.55 mm Hg in DBP in placebo-treated patients. There was no significant difference in the frequency of sustained (3 consecutive visits) elevated blood pressure [see WARNINGS AND PRECAUTIONS (5.3, 5.11)].

Duloxetine delayed-release capsules treatment, for up to 26 weeks in placebo-controlled trials across approved adult populations, typically caused a small increase in heart rate for change from baseline to endpoint compared to placebo of up to 1.37 beats per minute (increase of 1.20 beats per minute in duloxetine delayed-release capsules-treated patients, decrease of 0.17 beats per minute in placebo-treated patients).

Laboratory Changes in Adults

Duloxetine delayed-release capsules treatment in placebo-controlled clinical trials across approved adult populations, was associated with small mean increases from baseline to endpoint in ALT, AST, CPK, and alkaline phosphatase; infrequent, modest, transient, abnormal values were observed for these analytes in duloxetine delayed-release capsules-treated patients when compared with placebo-treated patients [see WARNINGS AND PRECAUTIONS (5.2)]. High bicarbonate, cholesterol, and abnormal (high or low) potassium, were observed more frequently in duloxetine delayed-release capsules-treated patients compared to placebo-treated patients.

Other Adverse Reactions Observed During the Clinical Trial Evaluation of Duloxetine Delayed-Release Capsules in Adults

Following is a list of adverse reactions reported by patients treated with duloxetine delayed-release capsules in clinical adult trials. In clinical trials of all approved adult populations, 34,756 patients were treated with duloxetine delayed-release capsules. Of these, 27% (9337) took duloxetine delayed-release capsules for at least 6 months, and

12% (4317) took duloxetine delayed-release capsules for at least one year. The following listing is not intended to include reactions (1) already listed in previous tables or elsewhere in labeling, (2) for which a drug cause was remote, (3) which were so general as to be uninformative, (4) which were not considered to have significant clinical implications, or (5) which occurred at a rate equal to or less than placebo.

Reactions are categorized by body system according to the following definitions: frequent adverse reactions are those occurring in at least 1/100 patients; infrequent adverse reactions are those occurring in 1/100 to 1/1000 patients; rare reactions are those occurring in fewer than 1/1000 patients.

- Cardiac Disorders — Frequent: palpitations; Infrequent: myocardial infarction, tachycardia, and Takotsubo cardiomyopathy.
- Ear and Labyrinth Disorders — Frequent: vertigo; Infrequent: ear pain and tinnitus.
- Endocrine Disorders — Infrequent: hypothyroidism.
- Eye Disorders — Frequent: vision blurred; Infrequent: diplopia, dry eye, and visual impairment.
- Gastrointestinal Disorders — Frequent: flatulence; Infrequent: dysphagia, eructation, gastritis, gastrointestinal hemorrhage, halitosis, and stomatitis; Rare: gastric ulcer.
- General Disorders and Administration Site Conditions — Frequent: chills/rigors; Infrequent: falls, feeling abnormal, feeling hot and/or cold, malaise, and thirst; Rare: gait disturbance.
- Infections and Infestations — Infrequent: gastroenteritis and laryngitis.
- Investigations — Frequent: weight increased, weight decreased; Infrequent: blood cholesterol increased.
- Metabolism and Nutrition Disorders — Infrequent: dehydration and hyperlipidemia; Rare: dyslipidemia.
- Musculoskeletal and Connective Tissue Disorders — Frequent: musculoskeletal pain; Infrequent: muscle tightness and muscle twitching.
- Nervous System Disorders — Frequent: dysgeusia, lethargy, and paraesthesia/hypoesthesia; Infrequent: disturbance in attention, dyskinesia, myoclonus, and poor quality sleep; Rare: dysarthria.
- Psychiatric Disorders — Frequent: abnormal dreams and sleep disorder; Infrequent: apathy, bruxism, disorientation/confusional state, irritability, mood swings, and suicide attempt; Rare: completed suicide.
- Renal and Urinary Disorders — Frequent: urinary frequency; Infrequent: dysuria, micturition urgency, nocturia, polyuria, and urine odor abnormal.
- Reproductive System and Breast Disorders — Frequent: anorgasmia/orgasm abnormal; Infrequent: menopausal symptoms, sexual dysfunction, and testicular pain; Rare: menstrual disorder.
- Respiratory, Thoracic and Mediastinal Disorders — Frequent: yawning, oropharyngeal pain; Infrequent: throat tightness.
- Skin and Subcutaneous Tissue Disorders — Frequent: pruritus; Infrequent: cold sweat, dermatitis contact, erythema, increased tendency to bruise, night sweats, and photosensitivity reaction; Rare: ecchymosis.
- Vascular Disorders — Frequent: hot flush; Infrequent: flushing, orthostatic hypotension, and peripheral coldness.

The data described below reflect exposure to duloxetine delayed-release capsules (N=476) in pediatric patients aged 7 to 17 years of age from two 10-week, placebo-controlled trials in patients with MDD (N=341) (Studies MDD-6 and MDD-7) and one 10-week placebo-controlled trial in GAD (N=135) (Study GAD-6). Duloxetine delayed-release capsules are not approved for the treatment of MDD in pediatric patients [see USE IN SPECIFIC POPULATIONS (8.4)]. Of the duloxetine delayed-release capsules-treated patients in these studies, 42 were 7 to 11 years of age (58% were between 12 to 17 years old), 51% were female, and 69% were white. Patients received 30 to 120 mg of duloxetine delayed-release capsules per day during placebo-controlled acute treatment studies. In the pediatric MDD and GAD, trials up to 36 weeks long, there were 822 duloxetine delayed-release capsules-treated pediatric patients aged 7 to 17 years of age (most patients received 30 to 120 mg per day) – 42% were 7 to 11 years of age (58% were 12 to 17 years old) and 52% were female.

Most Common Adverse Reactions in Pediatric Trials

The most common adverse reactions ($\geq 5\%$ in duloxetine delayed-release capsules-treated patients and at least twice the incidence of placebo-treated patients) in all pooled pediatric populations (MDD, GAD, and another indication) were decreased weight, decreased appetite, nausea, vomiting, fatigue, and diarrhea.

Adverse Reactions in Pediatric Patients Aged 7 to 17 Years Old with MDD and GAD

The adverse reaction profile observed in clinical trials in pediatric patients aged 7 to 18 years old with MDD and GAD was consistent with the adverse reaction profile observed in adult clinical trials. The most common ($\geq 5\%$ and twice placebo) adverse reactions observed in these pediatric clinical trials included: nausea, diarrhea, decreased weight, and dizziness.

Table 6 provides the incidence of adverse reactions in MDD and GAD pediatric placebo-controlled trials that occurred in greater than 2% of patients treated with duloxetine delayed-release capsules and with an incidence greater than patients treated with placebo. Duloxetine delayed-release capsules are not approved in the treatment of MDD in pediatric patients [see USE IN SPECIFIC POPULATIONS (8.4)].

Table 6: Adverse Reactions: Incidence of 2% or More and Greater than Placebo in Three 10-week Pediatric Placebo-Controlled Trials in MDD and GADa

System Organ Class/Adverse Reaction	Percentage of Pediatric Patients Reporting Reaction	
	Duloxetine Delayed-Release Capsules (N=476)	Placebo (N=362)
Gastrointestinal Disorders		
Nausea	18	8
Abdominal Pain ^b	13	10
Vomiting	9	4
Diarrhea	6	3
Dry Mouth	2	1
General Disorders and Administration Site Conditions	7	5

Fatigue ^c		
Investigations	14	6
Decreased Weight ^d		
Metabolism and Nutrition Disorders	10	5
Decreased Appetite		
Nervous System Disorders		
Headache	18	13
Somnolence ^e	11	6
Dizziness	8	4
Psychiatric Disorders		
Insomnia ^f	7	4
Respiratory, Thoracic, and Mediastinal Disorders	4	2
Oropharyngeal Pain	3	1
Cough		

Table 6: Adverse Reactions: Incidence of 2% or More and Greater than Placebo in Three 10-week Pediatric Placebo-Controlled Trials in MDD and GADa

a Duloxetine delayed-release capsules are not approved for the treatment of pediatric MDD [see USE IN SPECIFIC POPULATIONS (8.4)]. The inclusion of an event in the table is determined based on the percentages before rounding; however, the percentages displayed in the table are rounded to the nearest integer.

b Also includes abdominal pain upper, abdominal pain lower, abdominal tenderness, abdominal discomfort, and gastrointestinal pain.

c Also includes asthenia.

d Frequency based on weight measurement meeting potentially clinically significant threshold of $\geq 3.5\%$ weight loss (N=467 duloxetine delayed-release capsules; N=354 Placebo).

e Also includes hypersomnia and sedation.

f Also includes initial insomnia, insomnia, middle insomnia, and terminal insomnia.

System Organ Class/Adverse Reaction

Percentage of Pediatric Patients

Reporting Reaction

Duloxetine Delayed-Release Capsules (N=476)

Placebo

(N=362)

Gastrointestinal Disorders

Nausea

Abdominal Pain^b

Vomiting

Diarrhea

Dry Mouth

18

13

9

6

2

8
10
4
3
1
General Disorders and Administration Site Conditions
Fatiguec
7
5
Investigations
Decreased Weightd
14
6
Metabolism and Nutrition Disorders Decreased Appetite
10
5
Nervous System Disorders
Headache
Somnolencee
Dizziness

18
11
8

13
6
4
Psychiatric Disorders
Insomniaf

7
4
Respiratory, Thoracic, and Mediastinal Disorders
Oropharyngeal Pain
Cough
4
3
2
1

Other adverse reactions that occurred at an incidence of less than 2% and were reported by more duloxetine delayed-release capsules-treated patients than placebo-treated patients in pediatric MDD and GAD clinical trials included: abnormal dreams (including nightmare), anxiety, flushing (including hot flush), hyperhidrosis, palpitations, pulse increased, and tremor (duloxetine delayed-release capsules are not approved to treat pediatric patients with MDD).

The most commonly reported symptoms following discontinuation of duloxetine delayed-release capsules in pediatric MDD and GAD clinical trials included headache, dizziness, insomnia, and abdominal pain [see WARNINGS AND PRECAUTIONS (5.7)].

Growth (Height and Weight) in Pediatric Patients 7 to 17 Years Old with GAD and MDD

Decreased appetite and weight loss have been observed in association with the use of SSRIs and SNRIs. Duloxetine delayed-release capsules-treated pediatric patients in clinical trials experienced a 0.1 kg mean decrease in weight at 10 weeks, compared with a mean weight gain of approximately 0.9 kg in placebo-treated pediatric patients. The proportion of patients who experienced a clinically significant decrease in weight ($\geq 3.5\%$) was greater in the duloxetine delayed-release capsules group than in the placebo group (16% and 6%, respectively). Subsequently, over the 4- to 6-month uncontrolled extension periods, duloxetine delayed-release capsules-treated patients on average trended toward recovery to their expected baseline weight percentile based on population data from age- and sex-matched peers.

In studies up to 9 months, duloxetine delayed-release capsules-treated pediatric patients experienced an increase in height of 1.7 cm on average (2.2 cm increase in patients 7 to 11 years of age and 1.3 cm increase in patients 12 to 17 years of age). While height increase was observed during these studies, a mean decrease of 1% in height percentile was observed (decrease of 2% in patients 7 to 11 years of age and increase of 0.3% in patients 12 to 17 years of age). Weight and height should be monitored regularly in pediatric patients treated with duloxetine delayed-release capsules [see USE IN SPECIFIC POPULATIONS (8.4)].

Additional pediatric use information is approved for Eli Lilly and Company, Inc.'s CYMBALTA (duloxetine) delayed-release capsules. However, due to Eli Lilly and Company Inc.'s marketing exclusivity rights, this drug product is not labeled with that pediatric information.

6.2 Postmarketing Experience

The following adverse reactions have been identified during post approval use of duloxetine delayed-release capsules. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Adverse reactions reported since market introduction that were temporally related to duloxetine delayed-release capsules therapy and not mentioned elsewhere in labeling include: acute pancreatitis, anaphylactic reaction, aggression and anger (particularly early in treatment or after treatment discontinuation), angioneurotic edema, angle-closure glaucoma, colitis (microscopic or unspecified), cutaneous vasculitis (sometimes associated with systemic involvement), extrapyramidal disorder, galactorrhea, gynecological bleeding, hallucinations, hyperglycemia, hyperprolactinemia, hypersensitivity, hypertensive crisis, muscle spasm, rash, restless legs syndrome, seizures upon treatment discontinuation, supraventricular arrhythmia, tinnitus (upon treatment discontinuation), trismus, and urticaria.

7. Drug Interactions

Both CYP1A2 and CYP2D6 are responsible for duloxetine metabolism.

7.1 Inhibitors of CYP1A2

When duloxetine delayed-release capsules 60 mg was co-administered with fluvoxamine 100 mg, a potent CYP1A2 inhibitor, to male subjects (n=14) duloxetine AUC was

increased approximately 6-fold, the C_{max} was increased about 2.5-fold, and duloxetine t_{1/2} was increased approximately 3-fold. Other drugs that inhibit CYP1A2 metabolism include cimetidine and quinolone antimicrobials such as ciprofloxacin and enoxacin [see WARNINGS AND PRECAUTIONS (5.12)].

7.2 Inhibitors of CYP2D6

Concomitant use of duloxetine delayed-release capsules (40 mg once daily) with paroxetine (20 mg once daily) increased the concentration of duloxetine AUC by about 60%, and greater degrees of inhibition are expected with higher doses of paroxetine. Similar effects would be expected with other potent CYP2D6 inhibitors (e.g., fluoxetine, quinidine) [see WARNINGS AND PRECAUTIONS (5.12)].

7.3 Dual Inhibition of CYP1A2 and CYP2D6

Concomitant administration of duloxetine delayed-release capsules 40 mg twice daily with fluvoxamine 100 mg, a potent CYP1A2 inhibitor, to CYP2D6 poor metabolizer subjects (n=14) resulted in a 6-fold increase in duloxetine AUC and C_{max}.

7.4 Drugs that Interfere with Hemostasis (e.g., NSAIDs, Aspirin, and Warfarin)

Serotonin release by platelets plays an important role in hemostasis. Epidemiological studies of the case-control and cohort design that have demonstrated an association between use of psychotropic drugs that interfere with serotonin reuptake and the occurrence of upper gastrointestinal bleeding have also shown that concurrent use of an NSAID or aspirin may potentiate this risk of bleeding. Altered anticoagulant effects, including increased bleeding, have been reported when SSRIs or SNRIs are co-administered with warfarin. Concomitant administration of warfarin (2 to 9 mg once daily) under steady state conditions with duloxetine delayed-release capsules 60 or 120 mg once daily for up to 14 days in healthy subjects (n=15) did not significantly change INR from baseline (mean INR changes ranged from 0.05 to +0.07). The total warfarin (protein bound plus free drug) pharmacokinetics (AUC_{τ,ss}, C_{max,ss} or t_{max,ss}) for both R- and S-warfarin were not altered by duloxetine. Because of the potential effect of duloxetine on platelets, patients receiving warfarin therapy should be carefully monitored when duloxetine delayed-release capsules are initiated or discontinued [see WARNINGS AND PRECAUTIONS (5.5)].

7.5 Lorazepam

Under steady-state conditions for duloxetine delayed-release capsules (60 mg Q 12 hours) and lorazepam (2 mg Q 12 hours), the pharmacokinetics of duloxetine were not affected by co-administration.

7.6 Temazepam

Under steady-state conditions for duloxetine delayed-release capsules (20 mg qhs) and temazepam (30 mg qhs), the pharmacokinetics of duloxetine were not affected by co-administration.

7.7 Drugs that Affect Gastric Acidity

Duloxetine delayed-release capsules have an enteric coating that resists dissolution until reaching a segment of the gastrointestinal tract where the pH exceeds 5.5. In extremely acidic conditions, duloxetine delayed-release capsules, unprotected by the enteric coating, may undergo hydrolysis to form naphthol. Caution is advised in using duloxetine delayed-release capsules in patients with conditions that may slow gastric

emptying (e.g., some diabetics). Drugs that raise the gastrointestinal pH may lead to an earlier release of duloxetine. However, co-administration of duloxetine delayed-release capsules with aluminum- and magnesium-containing antacids (51 mEq) or duloxetine delayed-release capsules with famotidine, had no significant effect on the rate or extent of duloxetine absorption after administration of a 40 mg oral dose. It is unknown whether the concomitant administration of proton pump inhibitors affects duloxetine absorption [see WARNINGS AND PRECAUTIONS (5.14)].

7.8 Drugs Metabolized by CYP1A2

In vitro drug interaction studies demonstrate that duloxetine does not induce CYP1A2 activity. Therefore, an increase in the metabolism of CYP1A2 substrates (e.g., theophylline, caffeine) resulting from induction is not anticipated, although clinical studies of induction have not been performed. Duloxetine is an inhibitor of the CYP1A2 isoform in in vitro studies, and in two clinical studies the average (90% confidence interval) increase in theophylline AUC was 7% (1% to 15%) and 20% (13% to 27%) when co-administered with duloxetine delayed-release capsules (60 mg twice daily).

7.9 Drugs Metabolized by CYP2D6

Duloxetine is a moderate inhibitor of CYP2D6. When duloxetine delayed-release capsules was administered (at a dose of 60 mg twice daily) in conjunction with a single 50 mg dose of desipramine, a CYP2D6 substrate, the AUC of desipramine increased 3-fold [see WARNINGS AND PRECAUTIONS (5.12)].

7.10 Drugs Metabolized by CYP2C9

Results of in vitro studies demonstrate that duloxetine does not inhibit activity. In a clinical study, the pharmacokinetics of S-warfarin, a CYP2C9 substrate, were not significantly affected by duloxetine [see DRUG INTERACTIONS (7.4)].

7.11 Drugs Metabolized by CYP3A

Results of in vitro studies demonstrate that duloxetine does not inhibit or induce CYP3A activity. Therefore, an increase or decrease in the metabolism of CYP3A substrates (e.g., oral contraceptives and other steroidal agents) resulting from induction or inhibition is not anticipated, although clinical studies have not been performed.

7.12 Drugs Metabolized by CYP2C19

Results of in vitro studies demonstrate that duloxetine does not inhibit CYP2C19 activity at therapeutic concentrations. Inhibition of the metabolism of CYP2C19 substrates is therefore not anticipated, although clinical studies have not been performed.

7.13 Monoamine Oxidase Inhibitors (MAOIs)

[See DOSAGE AND ADMINISTRATION (2.9, 2.10), CONTRAINDICATIONS (4), and WARNINGS AND PRECAUTIONS (5.4)].

7.14 Serotonergic Drugs

[See DOSAGE AND ADMINISTRATION (2.9, 2.10), CONTRAINDICATIONS (4), and WARNINGS AND PRECAUTIONS (5.4)].

7.15 Alcohol

When duloxetine delayed-release capsules and ethanol were administered several hours apart so that peak concentrations of each would coincide, duloxetine delayed-release

capsules did not increase the impairment of mental and motor skills caused by alcohol.

In the duloxetine delayed-release capsules clinical trials database, three duloxetine delayed-release capsules-treated patients had liver injury as manifested by ALT and total bilirubin elevations, with evidence of obstruction. Substantial intercurrent ethanol use was present in each of these cases, and this may have contributed to the abnormalities seen [see WARNINGS AND PRECAUTIONS (5.2, 5.12)].

7.16 CNS Drugs

[See WARNINGS AND PRECAUTIONS (5.12)].

7.17 Drugs Highly Bound to Plasma Protein

Because duloxetine is highly bound to plasma protein, administration of duloxetine delayed-release capsules to a patient taking another drug that is highly protein bound may cause increased free concentrations of the other drug, potentially resulting in adverse reactions. However, co-administration of duloxetine delayed-release capsules (60 or 120 mg) with warfarin (2 to 9 mg), a highly protein-bound drug, did not result in significant changes in INR and in the pharmacokinetics of either total S- or total R-warfarin (protein bound plus free drug) [see DRUG INTERACTIONS (7.4)].

8. Use in Specific Populations

8.1 Pregnancy

Risk Summary

Data from a postmarketing retrospective cohort study indicate that use of duloxetine in the month before delivery may be associated with an increased risk of postpartum hemorrhage. Data from published literature and from a postmarketing retrospective cohort study have not identified a clear drug-associated risk of major birth defects or other adverse developmental outcomes (see Data). There are risks associated with untreated depression and fibromyalgia in pregnancy, and with exposure to SNRIs and SSRIs, including duloxetine delayed-release capsules, during pregnancy (see Clinical Considerations).

In rats and rabbits treated with duloxetine during the period of organogenesis, fetal weights were decreased but there was no evidence of developmental effects at doses up to 3 and 6 times, respectively, the maximum recommended human dose (MRHD) of 120 mg/day given to adolescents on a mg/m² basis. When duloxetine was administered orally to pregnant rats throughout gestation and lactation, pup weights at birth and pup survival to 1 day postpartum were decreased at a dose 2 times the MRHD given to adolescents on a mg/m² basis. At this dose, pup behaviors consistent with increased reactivity, such as increased startle response to noise and decreased habituation of locomotor activity were observed. Post-weaning growth was not adversely affected.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Clinical Considerations

Disease-associated Maternal and/or Embryo/Fetal Risk

Women who discontinue antidepressants during pregnancy are more likely to experience a relapse of major depression than women who continue antidepressants. This finding is from a prospective, longitudinal study that followed 201 pregnant women with a history of major depressive disorder who were euthymic and taking antidepressants at the beginning of pregnancy. Consider the risk of untreated depression when discontinuing or changing treatment with antidepressant medication during pregnancy and postpartum.

Pregnant women with fibromyalgia are at increased risk for adverse maternal and infant outcomes including preterm premature rupture of membranes, preterm birth, small for gestational age, intrauterine growth restriction, placental disruption, and venous thrombosis. It is not known if these adverse maternal and fetal outcomes are a direct result of fibromyalgia or other comorbid factors.

Maternal Adverse Reactions

Use of duloxetine in the month before delivery may be associated with an increased risk of postpartum hemorrhage [see WARNINGS AND PRECAUTIONS (5.5)].

Fetal/Neonatal Adverse Reaction

Neonates exposed to duloxetine delayed-release capsules and other SNRIs or SSRIs late in the third trimester have developed complications requiring prolonged hospitalization, respiratory support, and tube feeding. Such complications can arise immediately upon delivery. Reported clinical findings have included respiratory distress, cyanosis, apnea, seizures, temperature instability, feeding difficulty, vomiting, hypoglycemia, hypotonia, hypertonia, hyperreflexia, tremor, jitteriness, irritability, and constant crying. These findings are consistent with either a direct toxic effect of the SNRIs or SSRIs, or possibly, a drug discontinuation syndrome. It should be noted that, in some cases, the clinical picture is consistent with serotonin syndrome [see WARNINGS AND PRECAUTIONS (5.4)].

Data

Human Data

Data from a postmarketing retrospective claims-based cohort study found an increased risk for postpartum hemorrhage among 955 pregnant women exposed to duloxetine in the last month of pregnancy compared to 4,128,460 unexposed pregnant women (adjusted relative risk: 1.53; 95% CI: 1.08 to 2.18). The same study did not find a clinically meaningful increase in the risk for major birth defects in the comparison of 2532 women exposed to duloxetine in the first trimester of pregnancy to 1,284,827 unexposed women after adjusting for several confounders. Methodologic limitations include possible residual confounding, misclassification of exposure and outcomes, lack of direct measures of disease severity, and lack of information about alcohol use, nutrition, and over-the-counter medication exposures.

Animal Data

In animal reproduction studies, duloxetine has been shown to have adverse effects on embryo/fetal and postnatal development.

When duloxetine was administered orally to pregnant rats and rabbits during the period

of organogenesis, there was no evidence of malformations or developmental variations at doses up to 45 mg/kg/day [3 and 6 times, respectively, the MRHD of 120 mg/day given to adolescents on a mg/m² basis]. However, fetal weights were decreased at this dose, with a no-effect dose of 10 mg/kg/day (approximately equal to the MRHD in rats and 2 times the MRHD in rabbits).

When duloxetine was administered orally to pregnant rats throughout gestation and lactation, the survival of pups to 1 day postpartum and pup body weights at birth and during the lactation period were decreased at a dose of 30 mg/kg/day (2 times the MRHD given to adolescents on a mg/m² basis); the no-effect dose was 10 mg/kg/day. Furthermore, behaviors consistent with increased reactivity, such as increased startle response to noise and decreased habituation of locomotor activity, were observed in pups following maternal exposure to 30 mg/kg/day. Post-weaning growth and reproductive performance of the progeny were not affected adversely by maternal duloxetine treatment.

8.2 Lactation

Risk Summary

Data from published literature report the presence of duloxetine in human milk (see Data). There are reports of sedation, poor feeding, and poor weight gain in infants exposed to duloxetine through breast milk (see Clinical Considerations). There are no data on the effect of duloxetine on milk production.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for duloxetine delayed-release capsules and any potential adverse effects on the breastfed child from duloxetine delayed-release capsules or from the underlying maternal condition.

Clinical Considerations

Infants exposed to duloxetine delayed-release capsules should be monitored for sedation, poor feeding and poor weight gain.

Data

Disposition of duloxetine delayed-release capsules was studied in 6 lactating women who were at least 12 weeks postpartum and had elected to wean their infants. The women were given 40 mg of duloxetine delayed-release capsules twice daily for 3.5 days. The peak concentration measured in breast milk occurred at a median of 3 hours after the dose. The amount of duloxetine delayed-release capsules in breast milk was approximately 7 mcg/day while on that dose; the estimated daily infant dose was approximately 2 mcg/kg/day, which is less than 1% of the maternal dose. The presence of duloxetine delayed-release capsules metabolites in breast milk was not examined.

8.4 Pediatric Use

The safety and effectiveness of duloxetine delayed-release capsules have been established for treatment of generalized anxiety disorder (GAD) in patients 7 to 17 years of age. The safety and effectiveness of duloxetine delayed-release capsules have not been established in pediatric patients with major depressive disorder (MDD), diabetic peripheral neuropathic pain, or chronic musculoskeletal pain.

Antidepressants increased the risk of suicidal thoughts and behavior in pediatric patients. Monitor all pediatric patients being treated with antidepressants for clinical

worsening and emergence of suicidal thoughts and behaviors, especially during the initial few months of treatment, or at times of dosage changes [see Warnings and Precautions (5.1)]. Perform regular monitoring of weight and growth in pediatric patients treated with duloxetine delayed-release capsules [see ADVERSE REACTIONS (6.1)].

Generalized Anxiety Disorder

Use of duloxetine delayed-release capsules for the treatment of GAD in patients 7 to 17 years of age is supported by one 10-week, placebo-controlled trial (GAD-6). The study included 272 pediatric patients with GAD of which 47% were 7 to 11 years of age (53% were 12 to 17 years of age). Duloxetine delayed-release capsules demonstrated superiority over placebo as measured by greater improvement in the Pediatric Anxiety Rating Scale (PARS) for GAD severity score [see CLINICAL STUDIES (14.3)].

The safety and effectiveness of duloxetine delayed-release capsules for the treatment of GAD in pediatric patients less than 7 years of age have not been established.

Fibromyalgia

The safety and effectiveness of duloxetine delayed-release capsules for the treatment of fibromyalgia in patients less than 13 years of age have not been established.

Major Depressive Disorder

The safety and effectiveness of duloxetine delayed-release capsules have not been established in pediatric patients for the treatment of MDD. Efficacy of duloxetine delayed-release capsules was not demonstrated in two 10-week, placebo-controlled trials with 800 pediatric patients aged 7 to 17 years old with MDD (MDD-6 and MDD-7). Neither duloxetine delayed-release capsules nor an active control (approved for treatment of pediatric MDD) was superior to placebo.

The most frequently observed adverse reactions in the MDD pediatric clinical trials included nausea, headache, decreased weight, and abdominal pain. Decreased appetite and weight loss have been observed in association with the use of SSRIs and SNRIs.

Juvenile Animal Toxicology Data

Duloxetine administration to young rats from post-natal day 21 (weaning) through post-natal day 90 (adult) resulted in decreased body weights that persisted into adulthood, but recovered when drug treatment was discontinued; slightly delayed (~1.5 days) sexual maturation in females, without any effect on fertility; and a delay in learning a complex task in adulthood, which was not observed after drug treatment was discontinued. These effects were observed at the high dose of 45 mg/kg/day (2 times the MRHD, for a child); the no-effect-level was 20 mg/kg/day (≈1 times the MRHD, for a child).

Additional pediatric use information is approved for Eli Lilly and Company, Inc.'s CYMBALTA (duloxetine) delayed-release capsules. However, due to Eli Lilly and Company Inc.'s marketing exclusivity rights, this drug product is not labeled with that pediatric information.

8.5 Geriatric Use

Geriatric Exposure in Premarketing Clinical Trials of Duloxetine Delayed-Release Capsules

- Of the 2,418 patients in MDD trials, 6% (143) were 65 years of age or over.
- Of the 1041 patients in CLBP trials, 21% (221) were 65 years of age or over.
- Of the 487 patients in OA trials, 41% (197) were 65 years of age or over.
- Of the 1,074 patients in the DPNP trials, 33% (357) were 65 years of age or over.
- Of the 1,761 patients in FM trials, 8% (140) were 65 years of age or over.

In the MDD, GAD, DPNP, FM, OA, and CLBP studies, no overall differences in safety or effectiveness were generally observed between these patients and younger adult patients, and other reported clinical experience has not identified differences in responses between these geriatric and younger adult patients, but greater sensitivity of some older patients cannot be ruled out.

SSRIs and SNRIs, including duloxetine delayed-release capsules have been associated with clinically significant hyponatremia in geriatric patients, who may be at greater risk for this adverse reaction [see WARNINGS AND PRECAUTIONS (5.13)].

In an analysis of data from all placebo-controlled-trials, duloxetine delayed-release capsules-treated patients reported a higher rate of falls compared to placebo-treated patients. The increased risk appears to be proportional to a patient's underlying risk for falls. Underlying risk appears to increase steadily with age. As geriatric patients tend to have a higher prevalence of risk factors for falls such as medications, medical comorbidities and gait disturbances, the impact of increasing age by itself on falls during duloxetine delayed-release capsules treatment is unclear. Falls with serious consequences including bone fractures and hospitalizations have been reported with duloxetine delayed-release capsules use [see WARNINGS AND PRECAUTIONS (5.3) and ADVERSE REACTIONS (6.1)].

The pharmacokinetics of duloxetine after a single dose of 40 mg were compared in healthy elderly females (65 to 77 years) and healthy middle-age females (32 to 50 years). There was no difference in the C_{max}, but the AUC of duloxetine was somewhat (about 25%) higher and the half-life about 4 hours longer in the elderly females. Population pharmacokinetic analyses suggest that the typical values for clearance decrease by approximately 1% for each year of age between 25 to 75 years of age; but age as a predictive factor only accounts for a small percentage of between-patient variability. Dosage adjustment based on the age of the adult patient is not necessary.

8.6 Gender

Duloxetine's half-life is similar in men and women. Dosage adjustment based on gender is not necessary.

8.7 Smoking Status

Duloxetine bioavailability (AUC) appears to be reduced by about one-third in smokers. Dosage modifications are not recommended for smokers.

8.8 Race

No specific pharmacokinetic study was conducted to investigate the effects of race.

8.9 Hepatic Impairment

Patients with clinically evident hepatic impairment have decreased duloxetine metabolism and elimination. After a single 20 mg dose of duloxetine delayed-release capsules, 6 cirrhotic patients with moderate liver impairment (Child-Pugh Class B) had a mean

plasma duloxetine clearance about 15% that of age- and gender-matched healthy subjects, with a 5-fold increase in mean exposure (AUC). Although C_{max} was similar to normals in the cirrhotic patients, the half-life was about 3 times longer [see DOSAGE AND ADMINISTRATION (2.7) and WARNINGS AND PRECAUTIONS (5.14)].

8.10 Severe Renal Impairment

Limited data are available on the effects of duloxetine delayed-release capsules in patients with end-stage renal disease (ESRD). After a single 60 mg dose of duloxetine delayed-release capsules, C_{max} and AUC values were approximately 100% greater in patients with ESRD receiving chronic intermittent hemodialysis than in subjects with normal renal function. The elimination half-life, however, was similar in both groups. The AUCs of the major circulating metabolites, 4-hydroxy duloxetine glucuronide and 5-hydroxy, 6-methoxy duloxetine sulfate, largely excreted in urine, were approximately 7- to 9-fold higher and would be expected to increase further with multiple dosing. Population PK analyses suggest that mild to moderate degrees of renal impairment (estimated CrCl 30 to 80 mL/min) have no significant effect on duloxetine apparent clearance [see DOSAGE AND ADMINISTRATION (2.7) and WARNINGS AND PRECAUTIONS (5.14)].

9. Drug Abuse and Dependence

9.2 Abuse

In animal studies, duloxetine did not demonstrate barbiturate-like (depressant) abuse potential.

While duloxetine delayed-release capsules have not been systematically studied in humans for its potential for abuse, there was no indication of drug-seeking behavior in the clinical trials. However, it is not possible to predict on the basis of premarketing experience the extent to which a CNS active drug will be misused, diverted, and/or abused once marketed. Consequently, physicians should carefully evaluate patients for a history of drug abuse and follow such patients closely, observing them for signs of misuse or abuse of duloxetine delayed-release capsules (e.g., development of tolerance, incrementation of dose, drug-seeking behavior).

9.3 Dependence

In drug dependence studies, duloxetine did not demonstrate dependence-producing potential in rats.

10. Overdosage

10.1 Signs and Symptoms

In postmarketing experience, fatal outcomes have been reported for acute duloxetine delayed-release capsules overdoses, primarily with mixed overdoses, but also with duloxetine delayed-release capsules only, including 1000 mg of duloxetine delayed-release capsules (approximately 8.3 times the maximum recommended dosage). Signs and symptoms of overdose (duloxetine delayed-release capsules alone or with mixed drugs) included somnolence, coma, serotonin syndrome, seizures, syncope, tachycardia, hypotension, hypertension, and vomiting.

10.2 Management of Overdose

There is no specific antidote to a duloxetine delayed-release capsules overdose, but if serotonin syndrome ensues, specific treatment (such as with cyproheptadine and/or temperature control) may be considered.

In case of acute overdose with duloxetine delayed-release capsules, treatment should consist of those general measures employed in the management of overdose with any drug, such as assuring an adequate airway, oxygenation, and ventilation and monitoring cardiac rhythm and vital signs. Gastric lavage with a large-bore orogastric tube with appropriate airway protection, if needed, may be indicated if performed soon after ingestion or in symptomatic patients. Induction of emesis is not recommended.

Activated charcoal may be useful in limiting absorption of duloxetine from the gastrointestinal tract. Administration of activated charcoal has been shown to decrease duloxetine AUC and C_{max} by an average of one-third, although some patients had a limited effect of activated charcoal. Due to the large volume of distribution of duloxetine, forced diuresis, dialysis, hemoperfusion, and exchange transfusion are unlikely to be beneficial.


In managing overdose, the possibility of multiple drug involvement should be considered. A specific caution involves patients who overdose with duloxetine delayed-release capsules and tricyclic antidepressants. In such a case, decreased clearance of the parent tricyclic and/or its active metabolite may increase the possibility of clinically significant sequelae and extend the time needed for close medical observation [see WARNINGS AND PRECAUTIONS (5.4) and DRUG INTERACTIONS (7)].

Consider contacting a poison control center (1-800-222-1222 or www.poisson.org) for additional information on the treatment of overdose.

11. Description

Duloxetine delayed-release capsules are selective serotonin and norepinephrine reuptake inhibitor (SNRI) for oral administration. Its chemical designation is (+)-(S)-N-methyl-γ-(1-naphthoxy)-2-thiophenethylamine hydrochloride. The molecular formula is C₁₈H₁₉NOS•HCl, which corresponds to a molecular weight of 333.88. The structural formula is:

Duloxetine hydrochloride USP is a white to brownish-white solid, which is slightly soluble in water.



Each capsule contains enteric-coated pellets of 20, 30, or 60 mg of duloxetine (equivalent to 22.4, 33.7, or 67.3 mg of duloxetine hydrochloride USP, respectively). These enteric-coated pellets are designed to prevent degradation of the drug in the

acidic environment of the stomach. Inactive ingredients include crospovidone, hydroxy propyl cellulose, hypromellose, hypromellose phthalate, sugar spheres, talc, titanium dioxide, and triethylcitrate. The empty hard gelatin capsule shells contain gelatin, titanium dioxide, and sodium lauryl sulphate. In addition, the 20 mg and 60 mg contain FD&C Blue No. 2 and iron oxide yellow and 30 mg contains FD&C Blue No. 2. The capsules are printed with edible ink containing black iron oxide, potassium hydroxide, propylene glycol, shellac, and strong ammonia solution.

12. Clinical Pharmacology

12.1 Mechanism of Action

Although the exact mechanisms of the antidepressant, central pain inhibitory and anxiolytic actions of duloxetine in humans are unknown, these actions are believed to be related to its potentiation of serotonergic and noradrenergic activity in the CNS.

12.2 Pharmacodynamics

Preclinical studies have shown that duloxetine is a potent inhibitor of neuronal serotonin and norepinephrine reuptake and a less potent inhibitor of dopamine reuptake. Duloxetine has no significant affinity for dopaminergic, adrenergic, cholinergic, histaminergic, opioid, glutamate, and GABA receptors in vitro. Duloxetine does not inhibit monoamine oxidase (MAO).

Duloxetine delayed-release capsules are in a class of drugs known to affect urethral resistance [see WARNINGS AND PRECAUTIONS (5.15)].

Cardiac Electrophysiology

The effect of duloxetine delayed-release capsules 160 mg and 200 mg administered twice daily (2.7 and 3.3 times the maximum recommended dosage, respectively) to steady state was evaluated in a randomized, double-blinded, two-way crossover study in 117 healthy female adult subjects. No QT interval prolongation was detected. Duloxetine delayed-release capsules appears to be associated with concentration-dependent but not clinically meaningful QT shortening.

12.3 Pharmacokinetics

Duloxetine has an elimination half-life of about 12 hours (range 8 to 17 hours) and its pharmacokinetics are dose proportional over the therapeutic range. Steady-state plasma concentrations are typically achieved after 3 days of dosing. Elimination of duloxetine is mainly through hepatic metabolism involving two P450 isozymes, CYP1A2 and CYP2D6.

Absorption

After oral duloxetine delayed-release capsules administration, duloxetine hydrochloride is well absorbed. There is a median 2 hour lag until absorption begins (Tlag), with maximal plasma concentrations (Cmax) of duloxetine occurring 6 hours post dose. There is a 3 hour delay in absorption and a one-third increase in apparent clearance of duloxetine after an evening dose as compared to a morning dose.

Effect of Food: Food does not affect the Cmax of duloxetine, but delays the time to reach peak concentration from 6 to 10 hours and it marginally decreases the extent of absorption (AUC) by about 10%.

Distribution

The apparent volume of distribution averages about 1640 L. Duloxetine is highly bound (>90%) to proteins in human plasma, binding primarily to albumin and α 1-acid glycoprotein. The interaction between duloxetine and other highly protein bound drugs has not been fully evaluated. Plasma protein binding of duloxetine is not affected by renal or hepatic impairment.

Elimination

Metabolism

Biotransformation and disposition of duloxetine in humans have been determined following oral administration of ¹⁴C-labeled duloxetine. Duloxetine comprises about 3% of the total radiolabeled material in the plasma, indicating that it undergoes extensive metabolism to numerous metabolites. The major biotransformation pathways for duloxetine involve oxidation of the naphthyl ring followed by conjugation and further oxidation. Both CYP1A2 and CYP2D6 catalyze the oxidation of the naphthyl ring in vitro. Metabolites found in plasma include 4-hydroxy duloxetine glucuronide and 5-hydroxy, 6-methoxy duloxetine sulfate.

Excretion

Many additional metabolites have been identified in urine, some representing only minor pathways of elimination. Only trace (<1% of the dose) amounts of unchanged duloxetine are present in the urine. Most (about 70%) of the duloxetine dose appears in the urine as metabolites of duloxetine; about 20% is excreted in the feces. Duloxetine undergoes extensive metabolism, but the major circulating metabolites have not been shown to contribute significantly to the pharmacologic activity of duloxetine.

Specific Populations

Pediatric Patients

Duloxetine steady-state plasma concentration was comparable in pediatric patients 7 to 17 years of age and adult patients. The average steady-state duloxetine concentration was approximately 30% lower in this pediatric population relative to adult patients. The model-predicted duloxetine steady state plasma concentrations in pediatric patients 7 to 17 years of age were mostly within the concentration range observed in adult patients and did not exceed the concentration range in adults.

13. Non Clinical Toxicology

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Duloxetine was administered in the diet to mice and rats for 2 years.

In female mice receiving duloxetine at 140 mg/kg/day (3 times the maximum recommended human dose (MRHD) of 120 mg/day given to children on a mg/m² basis), there was an increased incidence of hepatocellular adenomas and carcinomas. The no-

effect dose was 50 mg/kg/day (1 time the MRHD given to children). Tumor incidence was not increased in male mice receiving duloxetine at doses up to 100 mg/kg/day (2 times the MRHD given to children).

In rats, dietary doses of duloxetine up to 27 mg/kg/day in females (1 time the MRHD given to children) and up to 36 mg/kg/day in males (1.4 times the MRHD given to children) did not increase the incidence of tumors.

Mutagenesis

Duloxetine was not mutagenic in the in vitro bacterial reverse mutation assay (Ames test) and was not clastogenic in an in vivo chromosomal aberration test in mouse bone marrow cells. Additionally, duloxetine was not genotoxic in an in vitro mammalian forward gene mutation assay in mouse lymphoma cells or in an in vitro unscheduled DNA synthesis (UDS) assay in primary rat hepatocytes, and did not induce sister chromatid exchange in Chinese hamster bone marrow in vivo.

Impairment of Fertility

Duloxetine administered orally to either male or female rats prior to and throughout mating at doses up to 45 mg/kg/day (3 times the MRHD given to adolescents on a mg/m² basis) did not alter mating or fertility.

14. Clinical Studies

14.1 Overview of the Clinical Trials

The efficacy of duloxetine delayed-release capsules has been established in the following populations in adequate and well-controlled trials:

- Major Depressive Disorder (MDD): 4 short-term (Studies MDD-1, MDD-2, MDD-3, and MDD-4) and 1 maintenance trial (Study MDD-5) in adults [see CLINICAL STUDIES (14.2)].
- Generalized Anxiety Disorder (GAD): 3 short-term trials in adults (Studies GAD-1, GAD-2, and GAD-3), 1 maintenance trial in adults (Study GAD-4), 1 short-term trial in geriatric patients (Study GAD-5), and 1 short-term trial in pediatric patients 7 to 17 years of age (Study GAD-6) [see CLINICAL STUDIES (14.3)].
- Diabetic Peripheral Neuropathic Pain (DPNP): Two 12-week trials in adults (Studies DPNP-1 and DPNP-2) [see CLINICAL STUDIES (14.4)].
- Fibromyalgia (FM): Two trials in adults (one of 3 months duration and one of 6 months duration) (Studies FM-1 and FM-2) [see CLINICAL STUDIES (14.5)].
- Chronic Musculoskeletal Pain: Two 12- to 13-week trials in adult patients with chronic low back pain (CLBP) (Studies CLBP-1 and CLBP-3) and one 13-week trial in adult patients with chronic pain due to osteoarthritis (OA) (Study OA-1) [see CLINICAL STUDIES (14.6)].

Additionally, a summary of the following trials that did not demonstrate efficacy are presented below: Study FM-3 (a 16-week trial in adult patients with fibromyalgia), Study CLBP-2 (a 13-week trial in adult patients with CLBP), and Study OA-2 (a 13-week trial in adult patients with chronic pain due to OA).

Additional pediatric use information is approved for Eli Lilly and Company, Inc.'s

CYMBALTA (duloxetine) delayed-release capsules. However, due to Eli Lilly and Company Inc.'s marketing exclusivity rights, this drug product is not labeled with that pediatric information.

14.2 Major Depressive Disorder in Adults

The efficacy of duloxetine delayed-release capsules as a treatment for MDD in adults was established in 4 randomized, double-blind, placebo-controlled, fixed-dose trials in adult outpatients (18 to 83 years) meeting DSM-IV criteria for MDD:

- In Studies MDD-1 and MDD-2, patients were randomized to duloxetine delayed-release capsules 60 mg once daily (N=123 and N=128, respectively) or placebo (N=122 and N=139, respectively) for 9 weeks
- In Study MDD-3, patients were randomized to duloxetine delayed-release capsules 20 or 40 mg twice daily (N=86 and N=91, respectively) or placebo (N=89) for 8 weeks
- In Study MDD-4, patients were randomized to duloxetine delayed-release capsules 40 or 60 mg twice daily (N=95 and N=93, respectively) or placebo (N=93) for 8 weeks.

In all four trials, duloxetine delayed-release capsules demonstrated superiority over placebo as measured by improvement in the 17-item Hamilton Depression Rating Scale (HAM-D-17) total score (see Table 8). There is no evidence that doses greater than 60 mg/day confer additional benefits.

In all of these clinical trials, analyses of the relationship between treatment outcome and age, gender, and race did not suggest any differential responsiveness on the basis of these patient characteristics.

Table 8: Summary of the Primary Efficacy Results for Adult Trials in MDD

Study Number	Treatment Group	Primary Efficacy Measure: HAM-D-17		
		Mean Baseline Score (SD)	LS Mean Change from Baseline (SE)	Placebo-subtracted Difference ^a (95% CI)
Study MDD-1	Duloxetine delayed-release capsules (60 mg/day) ^b	21.5 (4.10)	-10.9 (0.70)	-4.9 (-6.8, -2.9)
	Placebo	21.1 (3.71)	-6.1 (0.69)	--
Study MDD-2	Duloxetine delayed-release capsules (60 mg/day) ^b	20.3 (3.32)	-10.5 (0.71)	-2.2 (-4.0, -0.3)
	Placebo	20.5 (3.42)	-8.3 (0.67)	--
Study MDD-3	Duloxetine delayed-release capsules (20 mg BID) ^b	18.6 (5.85)	-7.4 (0.80)	-2.4 (-4.7, -0.2)
	Duloxetine delayed-release capsules (40 mg BID) ^b	18.1 (4.52)	-8.6 (0.81)	-3.6 (-5.9, -1.4)
	Placebo	17.2 (5.11)	-5.0 (0.81)	--
Study MDD-4	Duloxetine delayed-release capsules (40 mg BID) ^b	19.9 (3.54)	-11.0 (0.49)	-2.2 (-3.6, -0.9)
	Duloxetine delayed-release capsules (60 mg BID) ^b	20.2 (3.41)	-12.1 (0.49)	-3.3 (-4.7, -1.9)
	Placebo	19.9 (3.58)	-8.8 (0.50)	--


SD: standard deviation; SE: standard error; LS Mean: least-squares mean; CI: confidence interval, not adjusted for multiplicity in trials where multiple dose groups were included.

- a Difference (drug minus placebo) in least-squares mean change from baseline.
- b Doses statistically significantly superior to placebo.

In Study MDD-5, 533 adult patients meeting DSM-IV criteria for MDD received duloxetine delayed-release capsules 60 mg once daily during an initial 12-week open-label treatment phase. Two hundred and seventy-eight patients who responded to open label treatment [defined as meeting the following criteria at weeks 10 and 12: a HAMD-17 total score ≤ 9 , Clinical Global Impressions of Severity (CGI-S) ≤ 2 , and not meeting the DSM-IV criteria for MDD] were randomly assigned to continuation of duloxetine delayed-release capsules at the same dosage (N=136) or to placebo (N=142) for 6 months.

In Study MDD-5, patients on duloxetine delayed-release capsules experienced a statistically significantly longer time to relapse of depression than did patients on placebo (see Figure 1). Relapse was defined as an increase in the CGI-S score of ≥ 2 points compared with that obtained at week 12, as well as meeting the DSM-IV criteria for MDD at 2 consecutive visits at least 2 weeks apart, where the 2-week temporal criterion had to be satisfied at only the second visit.

Figure 1: Cumulative Proportion of Adult Patients with MDD Relapse (Study MDD-5)



a Kaplan-Meier estimator method.

14.3 Generalized Anxiety Disorder

GAD Trials in Adults (Including Geriatric Patients)

The efficacy of duloxetine delayed-release capsules in the treatment of generalized anxiety disorder (GAD) was established in 1 fixed-dose randomized, double-blind, placebo-controlled trial and 2 flexible-dose randomized, double-blind, placebo-controlled trials in adult outpatients between 18 and 83 years of age meeting the DSM-IV criteria for GAD (Studies GAD-1, GAD-2, and GAD-3, respectively).

In Studies GAD-1 and GAD-2, the starting dose was 60 mg once daily (down titration to 30 mg once daily was allowed for tolerability reasons; the dosage could be increased to 60 mg once daily). Fifteen percent of patients were down titrated. Study GAD-3 had a starting dose of 30 mg once daily for 1 week before increasing it to 60 mg once daily.

Studies GAD-2 and GAD-3 involved dose titration with duloxetine delayed-release capsules doses ranging from 60 mg once daily to 120 mg once daily (N=168 and N=162) compared to placebo (N=159 and N=161) over a 10-week treatment period. The mean dosage for completers at endpoint in these trials was 104.8 mg/day. Study

GAD-1 evaluated duloxetine delayed-release capsules dosages of 60 mg once daily (N=168) and 120 mg once daily (N=170) compared to placebo (N=175) over a 9-week treatment period. While a 120 mg/day dose was shown to be effective, there is no evidence that doses greater than 60 mg/day confer additional benefit.

In all 3 trials, duloxetine delayed-release capsules demonstrated superiority over placebo as measured by greater improvement in the Hamilton Anxiety Scale (HAM-A) total score (see Table 8) and by the Sheehan Disability Scale (SDS) global functional impairment score. The SDS is a composite measurement of the extent emotional symptoms disrupt patient functioning in 3 life domains: work/school, social life/leisure activities, and family life/home responsibilities.

In Study GAD-4, 887 patients meeting DSM-IV-TR criteria for GAD received duloxetine delayed-release capsules 60 mg to 120 mg once daily during an initial 26-week open-label treatment phase. Four hundred and twenty-nine patients who responded to open-label treatment [defined as meeting the following criteria at weeks 24 and 26: a decrease from baseline HAM-A total score by at least 50% to a score no higher than 11, and a Clinical Global Impressions of Improvement (CGI-Improvement) score of 1 or 2] were randomly assigned to continuation of duloxetine delayed-release capsules at the same dosage (N=216) or to placebo (N=213) and were observed for relapse. Of the patients randomized, 73% had been in a responder status for at least 10 weeks. Relapse was defined as an increase in CGI-Severity score at least 2 points to a score ≥ 4 and a MINI (Mini-International Neuropsychiatric Interview) diagnosis of GAD (excluding duration), or discontinuation due to lack of efficacy. Patients taking duloxetine delayed-release capsules experienced a statistically significantly longer time to relapse of GAD than did patients taking placebo (see Figure 2).

Subgroup analyses did not indicate that there were any differences in treatment outcomes as a function of age or gender.

GAD Trial in Geriatric Patients

The efficacy of duloxetine delayed-release capsules in the treatment of patients ≥ 65 years of age with GAD was established in one 10-week flexible-dose, randomized, double-blind, placebo-controlled trial in adults ≥ 65 years of age meeting the DSM-IV criteria for GAD (Study GAD-5). In Study GAD-5, the starting dose was 30 mg once daily for 2 weeks before further dose increases in 30 mg increments at treatment weeks 2, 4, and 7 up to 120 mg once daily were allowed based on investigator judgment of clinical response and tolerability. The mean dosage for patients completing the 10-week acute treatment phase was 51 mg. Patients treated with duloxetine delayed-release capsules (N=151) demonstrated significantly greater improvement compared with placebo (N=140) on mean change from baseline to endpoint as measured by the HAM-A total score (see Table 8).

GAD Trial in Pediatric Patients 7 to 17 Years Old

The efficacy of duloxetine delayed-release capsules in the treatment of pediatric patients 7 to 17 years of age with GAD was established in 1 flexible-dose randomized, double-blind, placebo-controlled trial in pediatric outpatients with GAD (based on DSM-IV criteria) (Study GAD-6).


In Study GAD-6, the starting dosage was 30 mg once daily for 2 weeks. Further dosage increases in 30 mg increments up to 120 mg once daily were allowed based on investigator judgment of clinical response and tolerability. The mean dosage for patients completing the 10-week treatment phase was 57.6 mg/day. In this study, duloxetine delayed-release capsules (N=135) demonstrated superiority over placebo (N=137) from baseline to endpoint as measured by greater improvement in the Pediatric Anxiety Rating Scale (PARS) for GAD severity score (see Table 9).

Table 9: Summary of the Primary Efficacy Results for GAD Trials

Study Number (population) (measurement)	Treatment Group	Primary Efficacy Measure		
		Mean Baseline Score (SD)	LS Mean Change from Baseline (SE)	Placebo- subtracted Difference ^a (95% CI)
Study GAD-1 (Adult) (HAM-A)	Duloxetine delayed- release capsules (60 mg/day) ^b	25.1 (7.18)	-12.8 (0.68)	-4.4 (-6.2, -2.5)
	Duloxetine delayed- release capsules (120 mg/day) ^b	25.1 (7.24)	-12.5 (0.67)	-4.1 (-5.9, -2.3)
	Placebo	25.8 (7.66)	-8.4 (0.67)	--
Study GAD-2 (Adult) (HAM-A)	Duloxetine delayed- release capsules (60 to 120 mg/day) ^b	22.5 (7.44)	-8.1 (0.70)	-2.2 (-4.2, -0.3)
	Placebo	23.5 (7.91)	-5.9 (0.70)	--
Study GAD-3 (Adult) (HAM-A)	Duloxetine delayed- release capsules (60 to 120 mg/day) ^b	25.8 (5.66)	-11.8 (0.69)	-2.6 (-4.5, -0.7)
	Placebo	25.0 (5.82)	-9.2 (0.67)	--
Study GAD-5 (Geriatric) (HAM-A)	Duloxetine delayed- release capsules (60 to 120 mg/day) ^b	24.6 (6.21)	-15.9 (0.63)	-4.2 (-5.9, -2.5)
	Placebo	24.5 (7.05)	-11.7 (0.67)	--
Study GAD-6 (Pediatric) (PARS for GAD)	Duloxetine delayed- release capsules (30 to 120 mg/day) ^b	17.5 (1.98)	-9.7 (0.50)	-2.7 (-4.0, -1.3)
	Placebo	17.4 (2.24)	-7.1 (0.50)	--

SD: standard deviation; SE: standard error; LS Mean: least-squares mean; CI: confidence interval, not adjusted for multiplicity in trials where multiple dose groups were included.
^a Difference (drug minus placebo) in least squares mean change from baseline.
^b Dose statistically significantly superior to placebo.

Figure 2: Cumulative Proportion of Adult Patients with GAD Relapse (Study GAD-4)



a Kaplan-Meier estimator method.

14.4 Diabetic Peripheral Neuropathic Pain in Adults

The efficacy of duloxetine delayed-release capsules for the management of neuropathic pain associated with diabetic peripheral neuropathy in adults was established in 2 randomized, 12-week, double-blind, placebo-controlled, fixed-dose trials in adult patients having diabetic peripheral neuropathic pain (DPNP) for at least 6 months (Study DPNP-1 and Study DPNP-2). These trials enrolled a total of 791 patients of whom 592 (75%) completed the trials. Patients enrolled had Type I or II diabetes mellitus with a diagnosis of painful distal symmetrical sensorimotor polyneuropathy for at least 6 months. The patients had a baseline pain score of ≥ 4 on an 11-point scale ranging from 0 (no pain) to 10 (worst possible pain). Patients were permitted up to 4 grams of acetaminophen per day as needed for pain, in addition to duloxetine delayed-release capsules. Patients recorded their pain daily in a diary.

Both trials compared duloxetine delayed-release capsules 60 mg once daily or 60 mg twice daily with placebo. Study DPNP-1 additionally compared duloxetine delayed-release capsules 20 mg with placebo. A total of 457 patients (342 duloxetine delayed-release capsules, 115 placebo) were enrolled in Study DPNP-1 and a total of 334 patients (226 duloxetine delayed-release capsules, 108 placebo) were enrolled in Study DPNP-2.

Treatment with duloxetine delayed-release capsules 60 mg one or two times a day statistically significantly improved the endpoint mean pain scores from baseline and increased the proportion of patients with at least a 50% reduction in pain scores from baseline. For various degrees of improvement in pain from baseline to study endpoint, Figures 3 and 4 show the fraction of patients achieving that degree of improvement in Studies DPNP-1 and DPNP-2, respectively. The figures are cumulative, so that patients whose change from baseline is, for example, 50%, are also included at every level of improvement below 50%. Patients who did not complete the trial were assigned 0% improvement. Some patients experienced a decrease in pain as early as week 1, which persisted throughout the trial.

Figure 3: Percentage of DPNP Adult Patients Achieving Various Levels of Pain Relief as Measured by 24-Hour Average Pain Severity (Study DPNP-1)





Figure 4: Percentage of DPNP Adult Patients Achieving Various Levels of Pain Relief as Measured by 24-Hour Average Pain Severity (Study DPNP-2)



14.5 Fibromyalgia

Adult Trials in Fibromyalgia

The efficacy of duloxetine delayed-release capsules for the management of fibromyalgia in adults was established in two randomized, double-blind, placebo-controlled, fixed-dose trials in adult patients meeting the American College of Rheumatology criteria for fibromyalgia (a history of widespread pain for 3 months, and pain present at 11 or more of the 18 specific tender point sites). Study FM-1 was three months in duration and enrolled female patients only. Study FM-2 was six months in duration and enrolled male and female patients. Approximately 25% of participants had a comorbid diagnosis of MDD. Studies FM-1 and FM-2 enrolled a total of 874 patients of whom 541 (62%) completed the trials. A total of 354 patients (234 duloxetine delayed-release capsules, 120 placebo) were enrolled in Study FM-1 and a total of 520 patients (376 duloxetine delayed-release capsules, 144 placebo) were enrolled in Study FM-2 (5% male, 95% female). The patients had a baseline pain score of 6.5 on an 11-point scale ranging from 0 (no pain) to 10 (worst possible pain).

Studies FM-1 and FM-2 compared duloxetine delayed-release capsules 60 mg once daily or 120 mg daily (given in divided doses in Study FM-1 and as a single daily dose in Study

FM-2) with placebo. Study FM-2 additionally compared duloxetine delayed-release capsules 20 mg with placebo during the initial three months of a six-month trial.

Treatment with duloxetine delayed-release capsules 60 mg or 120 mg daily statistically significantly improved the endpoint mean pain scores from baseline and increased the proportion of patients with at least a 50% reduction in pain score from baseline. Pain reduction was observed in patients both with and without comorbid MDD. However, the degree of pain reduction may be greater in patients with comorbid MDD. For various degrees of improvement in pain from baseline to study endpoint, Figures 5 and 6 show the fraction of patients achieving that degree of improvement in Studies FM-1 and FM-2, respectively. The figures are cumulative so that patients whose change from baseline is, for example, 50%, are also included at every level of improvement below 50%. Patients who did not complete the trial were assigned 0% improvement. Some patients experienced a decrease in pain as early as week 1, which persisted throughout the trial. Improvement was also demonstrated on measures of function (Fibromyalgia Impact Questionnaires) and patient global impression of change (PGI). Neither trial demonstrated a benefit of 120 mg compared to 60 mg, and a higher dosage was associated with more adverse reactions and premature discontinuations of treatment.

Figure 5: Percentage of Adult Fibromyalgia Patients Achieving Various Levels of Pain Relief at Study Endpoint as Measured by 24-Hour Average Pain Severity (Study FM-1)





Figure 6: Percentage of Adult Fibromyalgia Patients Achieving Various Levels of Pain Relief at Study Endpoint as Measured by 24-Hour Average Pain Severity (Study FM-2)



Additionally, the benefit of up-titration in non-responders to duloxetine delayed-release capsules at 60 mg/day was evaluated in a separate trial (Study FM-3). Adult patients were initially treated with duloxetine delayed-release capsules 60 mg once daily for eight weeks in open-label fashion. Subsequently, completers of this phase were randomized to double-blind treatment with duloxetine delayed-release capsules at either 60 mg once daily or 120 mg once daily. Responders were defined as patients who had at least a 30% reduction in pain score from baseline at the end of the 8-week treatment. Patients who were non-responders at 8 weeks were no more likely to meet response criteria at the end of 60 weeks of treatment if blindly titrated to duloxetine delayed-release capsules 120 mg as compared to those who were blindly continued on duloxetine delayed-release capsules 60 mg.

Additional pediatric use information is approved for Eli Lilly and Company, Inc.'s CYMBALTA (duloxetine) delayed-release capsules. However, due to Eli Lilly and Company Inc.'s marketing exclusivity rights, this drug product is not labeled with that pediatric information.

14.6 Chronic Musculoskeletal Pain in Adults

Duloxetine delayed-release capsules are indicated for the treatment of chronic musculoskeletal pain in adults. This has been established in trials in adult patients with chronic low back pain and chronic pain due to osteoarthritis.

Trials in Chronic Low Back Pain in Adults

The efficacy of duloxetine delayed-release capsules in chronic low back pain (CLBP) in adults was assessed in two double-blind, placebo-controlled, randomized clinical trials of 13-weeks duration (Studies CLBP-1 and CLBP-2), and one of 12-weeks duration (CLBP-3). Studies CLBP-1 and CLBP-3 demonstrated efficacy of duloxetine delayed-release capsules in the treatment of CLBP. Patients in all trials had no signs of radiculopathy or spinal stenosis.

Study CLBP-1: Two hundred thirty-six adult patients (N=115 on duloxetine delayed-

release capsules, N=121 on placebo) enrolled and 182 (77%) completed 13-week treatment phase. After 7 weeks of treatment, duloxetine delayed-release capsules-treated patients with less than 30% reduction in average daily pain and who were able to tolerate 60 mg once daily had their duloxetine delayed-release capsules dosage, in a double-blinded fashion, increased to 120 mg once daily for the remainder of the trial. Patients had a mean baseline pain rating of 6 on a numerical rating scale ranging from 0 (no pain) to 10 (worst possible pain). After 13 weeks of treatment, patients taking duloxetine delayed-release capsules 60 to 120 mg daily had a significantly greater pain reduction compared to patients taking placebo. Randomization was stratified by the patients' baseline NSAIDs use status. Subgroup analyses did not indicate that there were differences in treatment outcomes as a function of NSAIDs use.

Study CLBP-2: Four hundred and four patients were randomized to receive fixed dosages of duloxetine delayed-release capsules daily or a matching placebo (N=59 on duloxetine delayed-release capsules 20 mg, N=116 on duloxetine delayed-release capsules 60 mg, N=112 on duloxetine delayed-release capsules 120 mg, N=117 on placebo) and 267 (66%) completed the entire 13-week trial. After 13 weeks of treatment, none of the three duloxetine delayed-release capsules dosages showed a statistically significant difference in pain reduction compared to placebo.

Study CLBP-3: Four hundred and one patients were randomized to receive fixed doses of duloxetine delayed-release capsules 60 mg daily or placebo (N=198 on duloxetine delayed-release capsules, N=203 on placebo), and 303 (76%) completed the trial. Patients had a mean baseline pain rating of 6 on a numerical rating scale ranging from 0 (no pain) to 10 (worst possible pain). After 12 weeks of treatment, patients taking duloxetine delayed-release capsules 60 mg daily had significantly greater pain reduction compared to patients taking placebo.

For various degrees of improvement in pain from baseline to study endpoint, Figures 8 and 9 show the fraction of patients in Studies CLBP-1 and CLBP-3 achieving that degree of improvement, respectively. The figures are cumulative, so that patients whose change from baseline is, for example, 50%, are also included at every level of improvement below 50%. Patients who did not complete the trial were assigned the value of 0% improvement.

Figure 8: Percentage of Adult Patients with CLBP Achieving Various Levels of Pain Relief as Measured by 24-Hour Average Pain Severity (Study CLBP-1)





Figure 9: Percentage of Adult Patients with CLBP Achieving Various Levels of Pain Relief as Measured by 24-Hour Average Pain Severity (Study CLBP-3)



Trials in Chronic Pain Due to Osteoarthritis in Adults

The efficacy of duloxetine delayed-release capsules in chronic pain due to osteoarthritis (OA) in adults was assessed in 2 double-blind, placebo-controlled, randomized clinical trials of 13-weeks duration (Study OA-1 and Study OA-2). All patients in both trials fulfilled the ACR clinical and radiographic criteria for classification of idiopathic OA of the

knee. Randomization was stratified by the patients' baseline NSAIDs-use status.


Patients assigned to duloxetine delayed-release capsules started treatment in both trials at a dose of 30 mg once daily for one week. After the first week, the dose of duloxetine delayed-release capsules was increased to 60 mg once daily. After 7 weeks of treatment with duloxetine delayed-release capsules 60 mg once daily, in Study OA-1 patients with sub-optimal response to treatment (<30% pain reduction) and tolerated duloxetine delayed-release capsules 60 mg once daily had their dose increased to 120 mg. However, in Study OA-2, all patients, regardless of their response to treatment after 7 weeks, were re-randomized to either continue receiving duloxetine delayed-release capsules 60 mg once daily or have their dosage increased to 120 mg once daily for the remainder of the trial. Patients in the placebo treatment groups in both trials received a matching placebo for the entire duration of trials. For both trials, efficacy analyses were conducted using 13-week data from the combined duloxetine delayed-release capsules 60 mg and 120 mg once daily treatment groups compared to the placebo group.

Study OA-1: Two hundred fifty-six patients (N=128 on duloxetine delayed-release capsules, N=128 on placebo) enrolled and 204 (80%) completed the trial. Patients had a mean baseline pain rating of 6 on a numerical rating scale ranging from 0 (no pain) to 10 (worst possible pain). After 13 weeks of treatment, patients taking duloxetine delayed-release capsules had significantly greater pain reduction than patients taking placebo. Subgroup analyses did not indicate that there were differences in treatment outcomes as a function of NSAIDs use.

Study OA-2: Two hundred thirty-one patients (N=111 on duloxetine delayed-release capsules, N=120 on placebo) enrolled and 173 (75%) completed the trial. Patients had a mean baseline pain of 6 on a numerical rating scale ranging from 0 (no pain) to 10 (worst possible pain). After 13 weeks of treatment, patients taking duloxetine delayed-release capsules did not show a significantly greater pain reduction than patients taking placebo.

In Study OA-1, for various degrees of improvement in pain from baseline to study endpoint, Figure 10 shows the fraction of patients achieving that degree of improvement. The figure is cumulative, so that patients whose change from baseline is, for example, 50%, are also included at every level of improvement below 50%. Patients who did not complete the trial were assigned the value of 0% improvement.

Figure 10: Percentage of Adult Patients with OA Achieving Various Levels of Pain Relief as Measured by 24-Hour Average Pain Severity (Study OA-1)



16. How Supplied/Storage and Handling

16.1 How Supplied

Duloxetine Delayed-Release Capsules USP, 30 mg are blue opaque/white opaque, size '3' hard gelatin capsule filled with white to off-white pellets and imprinted with 'X' on blue opaque cap and '02' on white opaque body with black ink.

Bottles of 30: NDC 80425-0014-01

Bottles of 60: NDC 80425-0014-02

16.2 Storage and Handling

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

17. Patient Counseling Information

Advise the patient to read the FDA-approved patient labeling (MEDICATION GUIDE).

-
- Suicidal Thoughts and Behaviors — Advise patients, their families, and their caregivers to look for the emergence of suicidal ideation and behavior, especially during treatment and when the dose is adjusted up or down and instruct them to report such symptoms to their healthcare provider [see BOXED WARNING and WARNINGS AND PRECAUTIONS (5.1)].
- Administration — Advise patients to swallow duloxetine delayed-release capsules whole and to not chew, crush, or open the capsule (do not sprinkle contents on food or mixed with liquids) because these actions might affect the enteric coating.

- Hepatotoxicity — Inform patients that severe liver problems, sometimes fatal, have been reported in patients treated with duloxetine delayed-release capsules. Instruct patients to talk to their healthcare provider if they develop itching, right upper belly pain, dark urine, or yellow skin/eyes while taking duloxetine delayed-release capsules, which may be signs of liver problems. Instruct patients to talk to their healthcare provider about their alcohol consumption. Use of duloxetine delayed-release capsules with heavy alcohol intake may be associated with severe liver injury [see WARNINGS AND PRECAUTIONS (5.2)].
- Alcohol — Although duloxetine delayed-release capsules does not increase the impairment of mental and motor skills caused by alcohol, use of duloxetine delayed-release capsules concomitantly with heavy alcohol intake may be associated with severe liver injury [see Warnings and Precautions (5.2) and Drug Interactions (7.15)].
- Orthostatic Hypotension, Falls and Syncope — Advise patients of the risk of orthostatic hypotension, falls and syncope, especially during the period of initial use and subsequent dose escalation, and in association with the use of concomitant drugs that might potentiate the orthostatic effect of duloxetine delayed-release capsules [see WARNINGS AND PRECAUTIONS (5.3)].
- Serotonin Syndrome — Caution patients about the risk of serotonin syndrome with the concomitant use of duloxetine delayed-release capsules and other serotonergic agents including triptans, tricyclic antidepressants, fentanyl, lithium, tramadol, buspirone, tryptophan, amphetamines, and St. John's Wort [see CONTRAINDICATIONS (4), WARNINGS AND PRECAUTIONS (5.4), and DRUG INTERACTIONS (7.14)]. Advise patients of the signs and symptoms associated with serotonin syndrome that may include mental status changes (e.g., agitation, hallucinations, delirium, and coma), autonomic instability (e.g., tachycardia, labile blood pressure, dizziness, diaphoresis, flushing, hyperthermia), neuromuscular changes (e.g., tremor, rigidity, myoclonus, hyperreflexia, incoordination), seizures, and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea). Caution patients to seek medical care immediately if they experience these symptoms.
- Increased Risk of Bleeding — Caution patients about the concomitant use of duloxetine delayed-release capsules and NSAIDs, aspirin, warfarin, or other drugs that affect coagulation since combined use of psychotropic drugs that interfere with serotonin reuptake and these agents has been associated with an increased risk of bleeding [see WARNINGS AND PRECAUTIONS (5.5) and USE IN SPECIFIC POPULATIONS (8.1)].
- Severe Skin Reactions — Caution patients that duloxetine delayed-release capsules may cause serious skin reactions. This may need to be treated in a hospital and may be life-threatening. Counsel patients to call their doctor right away or get emergency help if they have skin blisters, peeling rash, sores in their mouth, hives, or any other allergic reactions [see WARNINGS AND PRECAUTIONS (5.6)].
- Discontinuation of Treatment — Instruct patients that discontinuation of duloxetine delayed-release capsules may be associated with symptoms such as dizziness, headache, nausea, diarrhea, paresthesia, irritability, vomiting, insomnia, anxiety, hyperhidrosis, and fatigue, and should be advised not to alter their dosing regimen, or stop taking duloxetine delayed-release capsules without consulting their healthcare provider [see WARNINGS AND PRECAUTIONS (5.7)].
- Activation of Mania or Hypomania — Adequately screen patients with depressive symptoms for risk of bipolar disorder (e.g. family history of suicide, bipolar disorder, and depression) prior to initiating treatment with duloxetine delayed-release capsules. Advise patients to report any signs or symptoms of a manic reaction such as greatly

increased energy, severe trouble sleeping, racing thoughts, reckless behavior, talking more or faster than usual, unusually grand ideas, and excessive happiness or irritability [see WARNINGS AND PRECAUTIONS (5.8)].

- Angle-Closure Glaucoma — Advise patients that taking duloxetine delayed-release capsules can cause mild pupillary dilation, which in susceptible individuals, can lead to an episode of angle-closure glaucoma. Pre-existing glaucoma is almost always open-angle glaucoma because angle-closure glaucoma, when diagnosed, can be treated definitively with iridectomy. Open-angle glaucoma is not a risk factor for angle-closure glaucoma. Patients may wish to be examined to determine whether they are susceptible to angle-closure, and have a prophylactic procedure (e.g., iridectomy), if they are susceptible [see WARNINGS AND PRECAUTIONS (5.9)].
- Seizures — Advise patients to inform their healthcare provider if they have a history of seizure disorder [see WARNINGS AND PRECAUTIONS (5.10)].
- Effects on Blood Pressure — Caution patients that duloxetine delayed-release capsules may cause an increase in blood pressure [see WARNINGS AND PRECAUTIONS (5.11)].
- Concomitant Medications — Advise patients to inform their healthcare provider if they are taking, or plan to take, any prescription or over-the-counter medications, since there is a potential for interactions [see DOSAGE AND ADMINISTRATION (2.9, 2.10), CONTRAINDICATIONS (4), WARNINGS AND PRECAUTIONS (5.4, 5.12), and DRUG INTERACTIONS (7)].
- Hyponatremia — Advise patients that hyponatremia has been reported as a result of treatment with SNRIs and SSRIs, including duloxetine delayed-release capsules. Advise patients of the signs and symptoms of hyponatremia [see WARNINGS AND PRECAUTIONS (5.13)].
- Concomitant Illnesses — Advise patients to inform their healthcare provider about all of their medical conditions [see WARNINGS AND PRECAUTIONS (5.14)].
- Urinary Hesitation and Retention — Duloxetine delayed-release capsules are in a class of medicines that may affect urination. Instruct patients to consult with their healthcare provider if they develop any problems with urine flow [see WARNINGS AND PRECAUTIONS (5.15)].
- Pregnancy
 - Advise women to notify their healthcare provider if they become pregnant or intend to become pregnant during treatment with duloxetine delayed-release capsules.
 - Advise pregnant women or patients who intend to become pregnant that duloxetine delayed-release capsules use during the month before delivery may lead to an increased risk for postpartum hemorrhage and may increase the risk of neonatal complications requiring prolonged hospitalization, respiratory support, and tube feeding.
 - Advise pregnant women that there is a risk of relapse with discontinuation of antidepressants.
 - Advise patients that there is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to duloxetine during pregnancy [see USE IN SPECIFIC POPULATIONS (8.1)].
- Lactation — Advise breastfeeding women using duloxetine delayed-release capsules to monitor infants for sedation, poor feeding and poor weight gain and to seek medical care if they notice these signs [see USE IN SPECIFIC POPULATIONS (8.2)].
- Interference with Psychomotor Performance — Duloxetine delayed-release capsules

may be associated with sedation and dizziness. Therefore, caution patients about operating hazardous machinery including automobiles, until they are reasonably certain that duloxetine delayed-release capsules therapy does not affect their ability to engage in such activities.

Dispense with Medication Guide available at: <http://www.risingpharma.com/med-guides.html>

Distributed by:
Rising Health, LLC
Saddle Brook, NJ 07663

Made in India

Code: TS/DRUGS/19/1993

Revised: 11/2020

Medication Guide

MEDICATION GUIDE

Duloxetine Delayed-Release Capsules USP

(doo lox' e teen)

Read this Medication Guide before you start taking duloxetine delayed-release capsules and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or treatment.

Talk to your healthcare provider about:

all risks and benefits of treatment with antidepressant medicines
all treatment choices for depression or other serious mental illness

What is the most important information I should know about antidepressant medicines, depression, other serious mental illnesses, and suicidal thoughts or actions?

1. Duloxetine delayed-release capsules and other antidepressant medicines may increase suicidal thoughts or actions in some children, teenagers, or young adults within the first few months of treatment or when the dose is changed.

2. Depression and other serious mental illnesses are the most important causes of suicidal thoughts or actions. Some people may have a particularly high risk of having suicidal thoughts or actions. These include people who have (or have a family history of) bipolar illness (also called manic-depressive illness).

3. How can I watch for and try to prevent suicidal thoughts and actions?

Pay close attention to any changes in mood, behavior, actions, thoughts, or feelings, especially sudden changes. This is very important when an antidepressant medicine is started or when the dose is changed.

Call your healthcare provider right away to report new or sudden changes in mood,

behavior, thoughts, or feelings.

Keep all follow-up visits with your healthcare provider as scheduled. Call your healthcare provider between visits as needed, especially if you have concerns about symptoms.

Call your healthcare provider right away if you have any of the following symptoms or feelings, especially if they are new, worse, or worry you. In an emergency, call 911.

- attempts to commit suicide
- acting on dangerous impulses
- acting aggressive, being angry, or violent
- thoughts about suicide or dying
- new or worse depression
- new or worse anxiety
- panic attacks
- feeling very agitated or restless
- new or worse irritability
- trouble sleeping
- an extreme increase in activity or talking (mania)
- other unusual changes in behavior or mood

What else do I need to know about antidepressant medicines?

Never stop an antidepressant medicine without first talking to a healthcare provider.

Stopping an antidepressant medicine suddenly can cause other symptoms.

Antidepressants are medicines used to treat depression and other illnesses. It is important to discuss all the risks of treating depression and also the risks of not treating it. Patients should discuss all treatment choices with your healthcare provider, not just the use of antidepressants.

Antidepressant medicines have other side effects. Talk to your healthcare provider about the side effects of the medicine prescribed for you or your family member.

Antidepressant medicines can interact with other medicines. Know all of the medicines that you or your family member takes. Keep a list of all medicines to show your healthcare provider. Do not start new medicines without first checking with your healthcare provider.

What are duloxetine delayed-release capsules?

Duloxetine delayed-release capsules are a prescription medicine used to treat a certain type of depression called Major Depressive Disorder (MDD). Duloxetine delayed-release capsules belong to a class of medicines known as SNRIs (or serotonin-norepinephrine reuptake inhibitors).

Duloxetine delayed-release capsules are also used to treat or manage:

- Generalized Anxiety Disorder (GAD)
- Diabetic Peripheral Neuropathic Pain (DPNP)
- Fibromyalgia (FM)
- Chronic Musculoskeletal Pain

Who should not take duloxetine delayed-release capsules?

Do Not take duloxetine delayed-release capsules if you:

take a Monoamine Oxidase Inhibitor (MAOI). Ask your healthcare provider or pharmacist if you are not sure if you take an MAOI, including the antibiotic linezolid or intravenous

methylene blue.

Do not take an MAOI within 5 days of stopping duloxetine delayed-release capsules unless directed to do so by your healthcare provider.

Do not start duloxetine delayed-release capsules if you stopped taking an MAOI in the last 14 days unless directed to do so by your healthcare provider.

People who take duloxetine delayed-release capsules close in time to an MAOI may have a serious problem called Serotonin Syndrome (see “What are the possible side effects of duloxetine delayed-release capsules?”).

What should I tell my healthcare provider before taking duloxetine delayed-release capsules?

Before starting duloxetine delayed-release capsules, tell your healthcare provider if you:

have heart problems or high blood pressure

have diabetes (duloxetine delayed-release capsules treatment makes it harder for some people with diabetes to control their blood sugar)

have liver problems

have kidney problems

have glaucoma

have or had seizures or convulsions

have bipolar disorder or mania

have low sodium levels in your blood

have delayed stomach emptying

have or had bleeding problems

are pregnant or plan to become pregnant. Duloxetine delayed-release capsules may harm your unborn baby. Talk to your healthcare provider about the risk to your unborn baby if you take duloxetine delayed-release capsules during pregnancy.

Tell your healthcare provider right away if you become pregnant or think you are pregnant during treatment with duloxetine delayed-release capsules.

are breastfeeding or plan to breastfeed. Duloxetine passes into your breast milk and may harm your baby. Talk to your healthcare provider about the best way to feed your baby while taking duloxetine delayed-release capsules.

Tell your healthcare provider about all the medicines that you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Duloxetine delayed-release capsules and some medicines may interact with each other, may not work as well, or may cause serious side effects.

Especially tell your healthcare provider if you take:

triptans used to treat migraine headache

medicines used to treat mood, anxiety, psychotic or thought disorders, including tricyclics, lithium, buspirone, SSRIs, SNRIs or MAOIs

tramadol and fentanyl

amphetamines

cimetidine

the antibiotics ciprofloxacin, enoxacin

medicine to treat irregular heart rate (like propafenone, flecainide, quinidine)

theophylline

the blood thinner warfarin (Coumadin, Jantoven)
non-steroidal anti-inflammatory drug (NSAID) (like ibuprofen, naproxen or aspirin).
over-the-counter supplements such as tryptophan or St. John's Wort
thioridazine (Mellaril). Mellaril together with duloxetine delayed-release capsules can
cause serious heart rhythm problems or sudden death.

Ask your healthcare provider for a list of these medicines if you are not sure.

Do not take duloxetine delayed-release capsules with any other medicine that contain
duloxetine.

How should I take duloxetine delayed-release capsules?

Take duloxetine delayed-release capsules exactly as your healthcare provider tells you to
take it. Your healthcare provider may need to change the dose of duloxetine delayed-
release capsules until it is the right dose for you.

Swallow duloxetine delayed-release capsules whole. Do not chew or crush duloxetine
delayed-release capsules.

Do not open the capsule and sprinkle on food or mix with liquids. Opening the capsule
may affect how well duloxetine delayed-release capsules works.

Duloxetine delayed-release capsules may be taken with or without food.

If you miss a dose of duloxetine delayed-release capsules, take the missed dose as soon
as you remember. If it is almost time for the next dose, skip the missed dose and take
your next dose at the regular time. Do not take two doses of duloxetine delayed-release
capsules at the same time.

If you take too much duloxetine hydrochloride, call your healthcare provider or poison
control center at 1-800-222-1222 right away, or get emergency treatment.

When switching from another antidepressant to duloxetine delayed-release capsules
your healthcare provider may want to lower the dose of the initial antidepressant first to
potentially avoid side effects.

What should I avoid while taking duloxetine delayed-release capsules?

Duloxetine delayed-release capsules can cause sleepiness or may affect your ability to
make decisions, think clearly, or react quickly. You should not drive, operate heavy
machinery, or do other dangerous activities until you know how duloxetine delayed-
release capsules affect you.

Use of duloxetine delayed-release capsules concomitantly with heavy alcohol intake may
be associated with severe liver injury. Avoid heavy alcohol use while taking duloxetine
delayed-release capsules.

What are the possible side effects of duloxetine delayed-release capsules?

Duloxetine delayed-release capsules may cause serious side effects, including: See
"What is the most important information I should know about duloxetine delayed-release
capsules?"

Common possible side effects in people who take duloxetine delayed-release capsules
include:

1. liver damage. Symptoms may include:

itching
right upper abdominal pain
dark urine

yellow skin or eyes
enlarged liver
increased liver enzymes

2. changes in blood pressure and falls. Monitor your blood pressure before starting and throughout treatment. Duloxetine delayed-release capsules may:

increase your blood pressure.

decrease your blood pressure when standing and cause dizziness or fainting, mostly when first starting duloxetine delayed-release capsules or when increasing the dose.

increase risk of falls, especially in elderly.

3. Serotonin Syndrome: This condition can be life-threatening and symptoms may include:

agitation, hallucinations, coma or other changes in mental status

coordination problems or muscle twitching (overactive reflexes)

racing heartbeat, high or low blood pressure

sweating or fever

nausea, vomiting, or diarrhea

muscle rigidity

dizziness

flushing

tremor

seizures

4. abnormal bleeding: Duloxetine delayed-release capsules and other antidepressant medicines may increase your risk of bleeding or bruising, especially if you take the blood thinner warfarin (Coumadin, Jantoven), a non-steroidal anti-inflammatory drug (NSAIDs, like ibuprofen or naproxen), or aspirin.

5. severe skin reactions: Duloxetine delayed-release capsules may cause serious skin reactions that may require stopping its use. This may need to be treated in a hospital and may be life-threatening. Call your healthcare provider right away or get emergency help if you have skin blisters, peeling rash, sores in the mouth, hives or any other allergic reactions.

6. discontinuation symptoms: Do not stop duloxetine delayed-release capsules without first talking to your healthcare provider. Stopping duloxetine delayed-release capsules too quickly or changing from another antidepressant too quickly may result in serious symptoms including:

anxiety

irritability

feeling tired or problems sleeping

headache

sweating

dizziness

electric shock-like sensations

vomiting or nausea

diarrhea

7. manic episodes:

greatly increased energy

- severe trouble sleeping
- racing thoughts
- reckless behavior
- unusually grand ideas
- excessive happiness or irritability
- talking more or faster than usual

8. visual problems:

- eye pain
- changes in vision
- swelling or redness in or around the eye

Only some people are at risk for these problems. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are.

9. seizures or convulsions

10. low salt (sodium) levels in the blood. Elderly people may be at greater risk for this. Symptoms may include:

- headache
- weakness or feeling unsteady
- confusion, problems concentrating or thinking or memory problems

11. problems with urination. Symptoms may include:

- decreased urine flow
- unable to pass any urine

The most common side effects of duloxetine delayed-release capsules include:

- nausea
- dry mouth
- sleepiness
- fatigue
- constipation
- loss of appetite
- increased sweating
- dizziness

Common possible side effects in children and adolescents who take duloxetine delayed-release capsules include:

- nausea
- decreased weight
- dizziness

Side effects in adults may also occur in children and adolescents who take duloxetine delayed-release capsules. Children and adolescents should have height and weight monitored during treatment.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of duloxetine delayed-release capsules. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to 1-800-FDA-1088.

How should I store duloxetine delayed-release capsules?

Store duloxetine delayed-release capsules at room temperature between 20° to 25°C (68° to 77°F).

Keep duloxetine delayed-release capsules and all medicines out of the reach of children.

General information about the safe and effective use of duloxetine delayed-release capsules.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use duloxetine delayed-release capsules for a condition for which it was not prescribed. Do not give duloxetine delayed-release capsules to other people, even if they have the same symptoms that you have. It may harm them.

This Medication Guide summarizes the most important information about duloxetine delayed-release capsules. If you would like more information, talk with your healthcare provider. You may ask your healthcare provider or pharmacist for information about duloxetine delayed-release capsules that is written for healthcare professionals.

For more information, call Rising Health, LLC at 1-833-395-6928.

What are the ingredients in duloxetine delayed-release capsules?

Active ingredient: duloxetine hydrochloride

Inactive ingredients: crospovidone, hydroxy propyl cellulose, hypromellose, hypromellose phthalate, sugar spheres, talc, titanium dioxide, and triethylcitrate. The empty hard gelatin capsule shells contain gelatin, titanium dioxide, and sodium lauryl sulphate. In addition, the 20 mg and 60 mg contain FD&C Blue No. 2 and iron oxide yellow and 30 mg contains FD&C Blue No. 2. The capsules are printed with edible ink containing black iron oxide, potassium hydroxide, propylene glycol, shellac, and strong ammonia solution.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Coumadin, Jantoven, and Mellaril are trademarks of their respective owners and not trademarks of Aurobindo Pharma Limited.

Dispense with Medication Guide available at: <http://www.risingpharma.com/med-guides.html>

Distributed by:
Rising Health, LLC
Saddle Brook, NJ 07663

Made in India

Code: TS/DRUGS/19/1993

Revised: 04/2020

Principal Display Panel

Packed By: AdvancedRx NashvilleTN, 37217		Store at 20°-25°C (68°-77°F) Caution: Federal law PROHIBITS transfer of this drug to any person other than the patient for whom it was prescribed.	Packed By: AdvancedRx NashvilleTN, 37217		Store at 20°-25°C (68°-77°F) Caution: Federal law PROHIBITS transfer of this drug to any person other than the patient for whom it was prescribed.
DULOXETINE HCL 30MG CAPSULE DR			DULOXETINE HCL 30MG CAPSULE DR		
#30 Compare to CYMBALTA NDC: 80425-0014-01 Source NDC: 57237-0018-99 Lot: 30DUL301221 Expires: 1/1/2022			#60 Compare to CYMBALTA NDC: 80425-0014-02 Source NDC: 57237-0018-99 Lot: 60DUL301221 Expires: 1/1/2022		
					
RISING PHARM S/N: 000000116819		RISING PHARM S/N: 000000117035		RISING PHARM S/N: 000000117035	

DULOXETINE HCL DR

duloxetine hcl capsule, delayed release

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:80425-0014(NDC:57237-018)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DULOXETINE HYDROCHLORIDE (UNII: 9044SC542W) (DULOXETINE - UNII:O5TNM5N07U)	DULOXETINE	30 mg

Product Characteristics

Color	white, blue	Score	no score
Shape	CAPSULE	Size	16mm
Flavor		Imprint Code	X;02
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:80425-0014-2	60 in 1 BOTTLE; Type 0: Not a Combination Product	12/11/2022	
2	NDC:80425-0014-1	30 in 1 BOTTLE; Type 0: Not a Combination Product	12/11/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090778	12/11/2022	

Labeler - Advanced Rx Pharmacy of Tennessee, LLC (117023142)

Establishment

Name	Address	ID/FEI	Business Operations
Advanced Rx Pharmacy of Tennessee, LLC		117023142	repack(80425-0014)

Revised: 11/2022

Advanced Rx Pharmacy of Tennessee, LLC