

**CALCIUM GLUCONATE- calcium gluconate injection, solution**  
**Covetrus**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).*

-----

**CALCIUM GLUCONATE 23%**

**INDICATIONS:**

For use as an aid in the treatment of hypocalcemia (parturient paresis, milk fever) in cattle.

**DOSAGE AND ADMINISTRATION:**

Cattle: 250 to 500 mL

Administer intravenously or intraperitoneally. Solution should be warmed to body temperature and administered slowly. This product is sterile in unopened container. It contains no preservative. Use entire contents when first opened. It should be handled under strict aseptic conditions. If there is no apparent improvement in the animal's condition within 24 hours, consult a veterinarian.

**TAKE TIME OBSERVE LABEL DIRECTIONS**

NDC 11695-5411-5

Calcium gluconate 23%

Calcium borogluconate solution

Volume: 16.907 fl oz (500 mL)

Questions? (855) 724-3461

Reorder #069167

18-802

RMS 92-1174

AH-069167-01

REV: 0719

Lot No.

Exp. Date

Distributed by:

Covetrus North America

400 Metro Place North

Dublin, OH 43017

covetrus.com

Manufactured by:

Nova-Tech Inc.

Grand Island, NE 68801

Assembled in USA

## Composition:

Each 500 mL of sterile aqueous solution contains:

Calcium.....10.7 g

(as calcium borogluconate, equivalent to calcium gluconate, 23%)

## Milliequivalents per liter

### Cations

Calcium.....1069 mEq/L

### Anions

Gluconate.....1069 mEq/L

Store between 15°-30°C (59°-86°F).

**FOR ANIMAL USE ONLY**

**KEEP OUT OF REACH OF CHILDREN**



Composición: Cada 500 mL de solución acuosa estéril contienen:  
Calcio ..... 10.7 g  
(como borogluconato de calcio, equivalente a gluconato de calcio, 23%)  
Milliequivalentes por litro

Cationes  
Calcio ..... 1069 mEq/L

Aniones  
Gluconato ..... 1069 mEq/L

Indicaciones: Como ayuda en el tratamiento de hipocalcemia (paresis parturienta, fiebre de leche) en los bovinos.

#### Dosis y administración:

Bovinos: 250 a 500 mL

Administrar por vía intravenosa o intraperitoneal. La temperatura de la solución debe elevarse a la del cuerpo, y la administración debe ser lenta. Este producto es estéril en su envase sin abrir. No contiene agentes conservantes. Usar todo el contenido después de abrir. Debe manejarse bajo condiciones de asepsia estrictas. Si no hay mejoría aparente en las condiciones del animal dentro de las 24 horas, consultar a un veterinario.

Mantener en la temperatura ambiente controlada 15° y 30°C (59° y 86°F).

Tomarse el tiempo requerido para observar las direcciones de la etiqueta.

N° para nuevo pedido 069167



NDC: 11695-6411-5

# Calcium gluconate 23%

Calcium borogluconate solution  
Solución de borogluconato de calcio

Ⓢ Calcio gluconato 23%

**FOR ANIMAL USE ONLY  
KEEP OUT OF REACH OF CHILDREN**

**SOLAMENTE PARA USO ANIMAL  
MANTÉNGASE FUERA DEL ALCANCE DE  
LOS NIÑOS**

Volume: 16.907 fl oz (500 mL)

Composition: Each 500 mL of sterile aqueous solution contains:  
Calcium ..... 10.7 g  
(as calcium borogluconate, equivalent to calcium gluconate, 23%)  
Milliequivalents per liter

Cations  
Calcium ..... 1069 mEq/L

Anions  
Gluconate ..... 1069 mEq/L

Indications: For use as an aid in the treatment of hypocalcemia (parturient paresis, milk fever) in cattle.

#### Dosage and administration:

Cattle: 250 to 500 mL

Administer intravenously or intraperitoneally. Solution should be warmed to body temperature and administered slowly. This product is sterile in unopened container. It contains no preservative. Use entire contents when first opened. It should be handled under strict aseptic conditions. If there is no apparent improvement in the animal's condition within 24 hours, consult a veterinarian.

Store at controlled room temperature 15°-30°C (59°-86°F).

Questions? (855) 724-3461

Reorder #069167

18-802  
RMS 92-1174  
AH-069167-01  
REV: 0719  
Lot No.

Distributed by:  
Covetrus North America  
400 Metro Place North  
Dublin, OH 43017  
covetrus.com

Manufactured by:  
Nova-Tech, Inc.  
Grand Island, NE 68801

Exp. Date



Assembled  
in  
USA

Composición: Cada 500 mL de solución acuosa estéril contienen:  
 Calcio ..... 10.7 g  
 (como borogluconato de calcio, equivalente a gluconato de calcio, 23%)  
**Miliequivalentes por litro**

**Cationes**  
 Calcio ..... 1069 mEq/L  
**Aniones**  
 Gluconato ..... 1069 mEq/L

**Indicaciones:** Como ayuda en el tratamiento de hipocalcemia (paresis parturienta, fiebre de leche) en los bovinos.

**Dosis y administración:**

**Bovinos:** 250 a 500 mL  
 Administrar por vía intravenosa o intraperitoneal. La temperatura de la solución debe elevarse a la del cuerpo, y la administración debe ser lenta. Este producto es estéril en su envase sin abrir. No contiene agentes conservantes. Usar todo el contenido después de abrir. Debe manejarse bajo condiciones de asepsia estrictas. Si no hay mejoría aparente en las condiciones del animal dentro de las 24 horas, consultar a un veterinario.

Mantener en la temperatura ambiente controlada 15° y 30°C (59° y 86°F).

Tomarse el tiempo requerido para observar las direcciones de la etiqueta.

N° para nuevo pedido 069167



NDC: 11695-6411-5

# Calcium gluconate 23%

Calcium borogluconate solution  
 Solución de borogluconato de calcio

 **Calcio gluconato 23%**

**FOR ANIMAL USE ONLY  
 KEEP OUT OF REACH OF CHILDREN**

**SOLAMENTE PARA USO ANIMAL  
 MANTÉNGASE FUERA DEL ALCANCE DE  
 LOS NIÑOS**

**Volume: 16.907 fl oz (500 mL)**

Composition: Each 500 mL of sterile aqueous solution contains:  
 Calcium ..... 10.7 g  
 (as calcium borogluconate, equivalent to calcium gluconate, 23%)  
**Miliequivalents per liter**

**Cations**  
 Calcium ..... 1069 mEq/L  
**Anions**  
 Gluconate ..... 1069 mEq/L

**Indications:** For use as an aid in the treatment of hypocalcemia (parturient paresis, milk fever) in cattle.

**Dosage and administration:**

**Cattle:** 250 to 500 mL  
 Administer intravenously or intraperitoneally. Solution should be warmed to body temperature and administered slowly. This product is sterile in unopened container. It contains no preservative. Use entire contents when first opened. It should be handled under strict aseptic conditions. If there is no apparent improvement in the animal's condition within 24 hours, consult a veterinarian.

Store at controlled room temperature 15°-30°C (59°-86°F).

Questions? (855) 724-3461

**Reorder #069167**

18-802  
 RMS 92-1174  
 AH-069167-01  
 REV: 0719  
 Lot No.

Distributed by:  
 Covetrus North America  
 400 Metro Place North  
 Dublin, OH 43017  
 covetrus.com

Manufactured by:  
 Nova-Tech, Inc.  
 Grand Island, NE 68801



## CALCIUM GLUCONATE

calcium gluconate injection, solution

### Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:11695-6411
<b>Route of Administration</b>	INTRAVENOUS, INTRAPERITONEAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CALCIUM GLUCONATE (UNII: SQE6VB453K) (CALCIUM CATION - UNII:2M83C4R6ZB)	CALCIUM GLUCONATE	23 g in 100 mL

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11695-6411-5	500 mL in 1 BOTTLE, PLASTIC		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		03/16/2020	

**Labeler** - Covetrus (603750329)

**Registrant** - Nova-Tech, Inc. (196078976)

## Establishment

Name	Address	ID/FEI	Business Operations
Nova-Tech, Inc.		196078976	manufacture

Revised: 3/2020

Covetrus