

LIDOCAINE - lidocaine injection
Aspen Veterinary Resources, Ltd.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

LIDOCAINE INJECTION Local Anesthetic

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

For Animal Use Only

Keep Out of Reach of Children

INDICATIONS

Lidocaine is a potent local anesthetic producing epidural, nerve conduction and infiltration anesthesia.

CONTRAINDICATIONS

Lidocaine is contraindicated in animals with a known hypersensitivity to the drug.

PRECAUTIONS

Lidocaine is usually well tolerated. Nevertheless, as with all local anesthetics, untoward effects may occur due to hypersensitivity, faulty technique, overdosage and inadvertent intravascular or subarachnoid injection. In case of respiratory arrest, immediate resuscitation with oxygen is indicated.

DOSAGE AND ADMINISTRATION

Epidural

Cattle and Horses - 5 to 15 mL

Dogs and Cats - 1 mL per 10 pounds of body weight.

Nerve Block

Cattle and Horses - 5 to 20 mL

Infiltration

Dilute to 0.5% concentration (1 mL of 2% solution diluted with 3 mL of sterile water = 4 mL of a 0.5% solution)

COMPOSITION

Each mL of sterile aqueous solution contains:

Lidocaine Hydrochloride.....2.0%

Propylene Glycol.....5.2%

Sodium Chloride.....0.5%

Sodium Lactate.....0.5%

with Methylparaben 0.15%, Sodium Metabisulfite 0.10%,

Propylparaben 0.03%, and Disodium Edetate 0.001% as preservatives.

Store at controlled room temperature between 15°C and 30°C (59°F-86°F).

TAKE TIME OBSERVE LABEL DIRECTIONS

INDICATIONS: Lidocaine is a potent local anesthetic for producing epidural, nerve conduction and infiltration anesthesia.

CONTRAINDICATIONS: Lidocaine is contraindicated in animals with a known hypersensitivity to the drug.

PRECAUTIONS: Lidocaine is usually well tolerated. Nevertheless, as with all local anesthetics, untoward effects may occur due to hypersensitivity, faulty technique, overdosage and inadvertent intravascular or subarachnoid injection. In case of respiratory arrest, immediate resuscitation with oxygen is indicated.

DOSE AND ADMINISTRATION:

Epidural: Cattle and Horses – 5 to 15 mL

Dogs and Cats – 1 mL per 10 pounds of body weight

Nerve Block: Cattle and Horses – 5 to 20 mL

Infiltration: Dilute to 0.5% concentration (1 mL of 2% solution diluted with 3 mL of sterile water = 4 mL of a 0.5% solution).



Lidocaine Injectable

Local Anesthetic



**For Animal Use Only
Keep Out Of Reach Of Children**

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**NET CONTENTS:
3.38 fl oz (100 mL)**

COMPOSITION: Each mL of sterile aqueous solution contains:
Lidocaine Hydrochloride 2.0%
Propylene Glycol 5.2%
Sodium Chloride 0.5%
Sodium Lactate 0.5%

With Methylparaben 0.15%, Sodium Metabisulfite 0.10%, Propylparaben 0.03% and Disodium Edetate 0.001% as preservatives.

Store at controlled room temperature between 15° C and 30° C (59° F-86° F).



Manufactured for:
Aspen Veterinary Resources,® Ltd.
Liberty, MO 64068
www.aspenveterinaryresources.com

A2135
L-2000-04
Rev. 06/16

LOT NO.:

EXP. DATE:

LIDOCAINE

lidocaine injection

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:46066-222
Route of Administration	EPIDURAL, INFILTRATION		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	0.02 g in 1 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46066-222-04	100 mL in 1 BOTTLE		
2	NDC:46066-222-05	250 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		08/11/1995	

Labeler - Aspen Veterinary Resources, Ltd. (627265361)