

BASIC CARE CHILDRENS IBUPROFEN- ibuprofen tablet, chewable
Amazon.com Services LLC

Amazon Children's Ibuprofen Drug Facts

Active ingredient (in each chewable tablet)

Ibuprofen 100 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more

than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- mouth or throat burning may occur; give with food or water
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days

- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **chew or crush tablets completely before swallowing**
- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**

Dosing Chart

Weight (lb)	Age (yr)	Tablets
under 24	under 2	ask a doctor
24-35	2-3	1
36-47	4-5	1 ½
48-59	6-8	2
60-71	9-10	2 ½
72-95	11	3

Other information

- phenylketonurics: contains phenylalanine 6 mg per tablet
- store between 20-25°C (68-77°F)
- **do not use if printed seal under cap is broken or missing**

Inactive ingredients

acesulfame potassium, ammonium glycyrrhizin, aspartame, carnauba wax, croscarmellose sodium, hypromellose, magnesium stearate, mannitol, natural and artificial flavors, silicon dioxide, sodium lauryl sulfate, soybean oil, succinic acid

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

For Ages 2 to 11

Compare to Children's Motrin[®] active ingredient

Children's Ibuprofen

Chewable Tablets, 100 mg

Pain Reliever/Fever Reducer (NSAID)

Dye-Free

Orange Flavor

LASTS UP TO 8 HOURS


Chewables

actual size

Chew or crush tablets completely before swallowing

24 ORANGE-FLAVORED TABLETS

7W062 BN C1



KEEP CARTON FOR REFERENCE Gluten Free

Important: Read all product information before using. Keep this box for important information. This product is for children only.

Drug Facts

Active ingredient (in each chewable tablet) **Purposes**
 Ibuprofen (100 mg NSAID) Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses temporarily: ■ reduce fever
 ■ relieve minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings

Allergic alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ hives ■ facial swelling ■ asthma/wheezing
 ■ shock ■ skin redness ■ rash ■ hiccups

If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you child:
 ■ has had stomach ulcers or bleeding problems
 ■ takes a blood thinning (anticoagulant) or steroid drug
 ■ takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Some blood thinning, severe or persistent stomach or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Drug Facts (continued)

Do not use
 ■ if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
 ■ right before or after heart surgery

Ask a doctor before use if
 ■ stomach bleeding warning applies to your child
 ■ child has a history of stomach problems, such as heart burn
 ■ child has problems or serious side effects from taking pain relievers or fever reducers
 ■ child has not been drinking fluids
 ■ child has lost a lot of fluid due to vomiting or diarrhea
 ■ child has high blood pressure, heart disease, liver disease, kidney disease, or had a stroke
 ■ child has asthma ■ child is taking a diuretic

Ask a doctor or pharmacist before use if the child is
 ■ under a doctor's care for any serious condition
 ■ taking any other drug

When using this product

■ avoid hot liquid burning may occur, give with food or water
 ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if
 ■ child experiences any of the following signs of stomach bleeding:
 ■ dark stools ■ vomits blood ■ has blood or black stools
 ■ dark hair ■ vomits blood
 ■ child has symptoms of liver problems or stroke:
 ■ chest pain ■ trouble breathing
 ■ weakness in one part or side of body
 ■ slurred speech ■ leg swelling

Drug Facts (continued)

■ the child does not get any relief within first day (24 hours) of treatment
 ■ fever or pain/aches worsen or lasts more than 3 days
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222).

Directions

■ Chew or crush tablet completely before swallowing
 ■ This product does not contain directions or complete warnings for adult use
 ■ Do not give more than directed
 ■ Find right dose on chart below. If possible, use weight to dose; otherwise use age.
 ■ If needed, repeat dose every 6-8 hours
 ■ Do not use more than 4 times a day

Dosing Chart

Weight (lb)	Age (yr)	Tablets
under 24	under 2	ask a doctor
24-35	2-3	1
36-47	4-5	1½
48-59	6-8	2
60-71	9-10	2½
72-85	11	3



Drug Facts (continued)

Other information
 ■ Ibuprofen is a nonsteroidal anti-inflammatory drug (NSAID).
 ■ Do not use if the child has a known hypersensitivity to aspirin, ibuprofen, or other NSAIDs.
 ■ Do not use if the child has a history of stomach problems, such as heart burn or stomach bleeding.
 ■ Do not use if the child has a history of stomach problems, such as heart burn or stomach bleeding.
 ■ Do not use if the child has a history of stomach problems, such as heart burn or stomach bleeding.

Ingredients
 Active ingredients: acetaminophen, ibuprofen, polyethylene glycol, croscarmellose, hydroxypropyl methylcellulose, hypromellose, magnesium stearate, mannitol, natural and artificial flavors, silicon dioxide, xanthan gum, polyethylene glycol, croscarmellose, hypromellose, magnesium stearate, mannitol, natural and artificial flavors, silicon dioxide, xanthan gum.

Questions or comments? 1-800-719-9660

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Children's Motrin®.
 DISTRIBUTED BY: AMAZON.COM SERVICES LLC
 410 TERRY AVENUE N, SEATTLE, WA 98109
 © 2020 AMAZON.COM, INC. OR ITS AFFILIATES.
 ALL RIGHTS RESERVED.



For Ages 2 to 11 NDC 72288-763-62

Children's Ibuprofen

Chewable Tablets, 100 mg
 Pain Reliever/Fever Reducer (NSAID)

LASTS UP TO 8 HOURS
 Chewables
 Chew or crush tablets completely before swallowing

Dye-Free
Orange Flavor

actual size

24 ORANGE-FLAVORED TABLETS

BASIC CARE CHILDRENS IBUPROFEN
 ibuprofen tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72288-763
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ASPARTAME (UNII: Z0H242BBR1)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SOYBEAN OIL (UNII: 241ATL177A)	
SUCCINIC ACID (UNII: AB6MNQ6J6L)	

Product Characteristics

Color	WHITE (off white)	Score	2 pieces
Shape	ROUND	Size	13mm
Flavor	ORANGE	Imprint Code	L7W6
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72288-763-62	1 in 1 CARTON	08/11/2021	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076359	08/11/2021	

Revised: 8/2021

Amazon.com Services LLC