


**CHLORZOXAZONE- chlorzoxazone tablet**  
**Cameron Pharmaceuticals**

-----  
**Chlorzoxazone Tablets, USP 500 mg**

**DESCRIPTION**

Chlorzoxazone USP is a centrally acting skeletal muscle relaxant, available as tablets of 500 mg for oral administration. Its chemical name is 5-Chloro-2-benzoxazolinone, and its structural formula is:



$C_7H_4ClNO_2$   
MW 169.57

Chlorzoxazone USP is a white or practically white, practically odorless, crystalline powder. Chlorzoxazone is slightly soluble in water; sparingly soluble in alcohol, in isopropyl alcohol, and in methanol; soluble in solutions of alkali hydroxides and ammonia.

Chlorzoxazone tablets contain the inactive ingredients Pregelatinized Starch, Croscarmellose Sodium, Microcrystalline Cellulose, Docusate Sodium, Sodium Benzoate, Colloidal Silicon dioxide, Magnesium Stearate.

**CLINICAL PHARMACOLOGY**

Chlorzoxazone is a centrally-acting agent for painful musculoskeletal conditions. Data available from animal experiments as well as human study indicate that chlorzoxazone acts primarily at the level of the spinal cord and subcortical areas of the brain where it inhibits multisynaptic reflex arcs involved in producing and maintaining skeletal muscle spasm of varied etiology. The clinical result is a reduction of the skeletal muscle spasm with relief of pain and increased mobility of the involved muscles. Blood levels of chlorzoxazone can be detected in people during the first 30 minutes and peak levels may be reached, in the majority of the subjects, in about 1 to 2 hours after oral administration of chlorzoxazone. Chlorzoxazone is rapidly metabolized and is excreted in the urine, primarily in a conjugated form as the glucuronide. Less than one percent of a dose of chlorzoxazone is excreted unchanged in the urine in 24 hours.

**INDICATIONS**

Chlorzoxazone is indicated as an adjunct to rest, physical therapy, and other measures for the relief of discomfort associated with acute, painful musculoskeletal conditions. The mode of action of this drug has not been clearly identified, but may be related to its sedative properties. Chlorzoxazone does not directly relax tense skeletal muscles in man.

**CONTRAINDICATIONS**

Chlorzoxazone is contraindicated in patients with known intolerance to the drug.

## **WARNINGS**

Serious (including fatal) hepatocellular toxicity has been reported rarely in patients receiving chlorzoxazone. The mechanism is unknown but appears to be idiosyncratic and unpredictable. Factors predisposing patients to this rare event are not known. Patients should be instructed to report early signs and/or symptoms of hepatotoxicity such as fever, rash, anorexia, nausea, vomiting, fatigue, right upper quadrant pain, dark urine, or jaundice. Chlorzoxazone should be discontinued immediately and a physician consulted if any of these signs or symptoms develop. Chlorzoxazone use should also be discontinued if a patient develops abnormal liver enzymes (e.g., AST, ALT, alkaline phosphatase and bilirubin.)

The concomitant use of alcohol or other central nervous system depressants may have an additive effect.

### **Usage in Pregnancy**

The safe use of chlorzoxazone has not been established with respect to the possible adverse effects upon fetal development. Therefore, it should be used in women of childbearing potential only when, in the judgment of the physician, the potential benefits outweigh the possible risks.

## **PRECAUTIONS**

Chlorzoxazone should be used with caution in patients with known allergies or with a history of allergic reactions to drugs. If a sensitivity reaction occurs such as urticaria, redness, or itching of the skin, the drug should be stopped.

If any symptoms suggestive of liver dysfunction are observed, the drug should be discontinued.

## **ADVERSE REACTIONS**

Chlorzoxazone containing products are usually well tolerated. It is possible in rare instances that chlorzoxazone may have been associated with gastrointestinal bleeding. Drowsiness, dizziness, lightheadedness, malaise, or over-stimulation may be noted by an occasional patient. Rarely, allergic-type skin rashes, petechiae, or ecchymoses may develop during treatment. Angioneurotic edema or anaphylactic reactions are extremely rare. There is no evidence that the drug will cause renal damage. Rarely, a patient may note discoloration of the urine resulting from a phenolic metabolite of chlorzoxazone. This finding is of no known clinical significance.

To report SUSPECTED ADVERSE EVENTS, contact Novitium Pharma LLC at 1-855-204-1431 or FDA at 1-800- FDA-1088 or <http://www.fda.gov/> for voluntary reporting of adverse reactions.

## **OVERDOSAGE**

### **Symptoms**

Initially, gastrointestinal disturbances such as nausea, vomiting, or diarrhea together with drowsiness, dizziness, lightheadedness or headache may occur. Early in the course there may be malaise or sluggishness followed by marked loss of muscle tone, making voluntary movement impossible. The deep tendon reflexes may be decreased or absent. The sensorium remains intact, and there is no peripheral loss of sensation. Respiratory depression may occur with rapid, irregular respiration and intercostal and substernal retraction. The blood pressure is lowered, but shock has not been observed.

### **Treatment**

Gastric lavage or induction of emesis should be carried out, followed by administration of activated charcoal. Thereafter, treatment is entirely supportive. If respirations are depressed, oxygen and artificial respiration should be employed and a patent airway assured by use of an oropharyngeal airway or endotracheal tube. Hypotension may be counteracted by use of dextran, plasma, concentrated albumin or a vasopressor agent such as norepinephrine. Cholinergic drugs or analeptic drugs are of no value and should not be used.

## **DOSAGE AND ADMINISTRATION**

### **Usual Adult Dosage**

One tablet three or four times daily. If adequate response is not obtained with this dose, it may be increased to one and one-half tablets (750 mg) three or four times daily. As improvement occurs dosage can usually be reduced.

### **HOW SUPPLIED**

Chlorzoxazone Tablets, USP 500 mg for Oral administration are supplied as follows, White to off-white capsule shaped tablets debossed with N085 on the bisected side:

NDC 42494-426-10	100 Tablets
NDC 42494-427-50	500 Tablets

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense contents with a child-resistant closure (as required) and in a tight container as defined in the USP/NF.

#### **Manufactured for:**

Cameron Pharmaceuticals  
Louisville, KY 40245, USA

#### **Manufactured by:**

Novitium Pharma LLC,  
70 Lake Drive,  
East Windsor, New Jersey-08520

**Issued:** 08/2019

LB4157-02

### **PRINCIPAL DISPLAY PANEL - 100 Tablet Bottle Label**

NDC 42494-426-10  
100 Tablets

Chlorzoxazone  
Tablets, USP

500 mg

CAMERON  
PHARMACEUTICALS

Rx Only

NDC 42494-426-10 100 Tablets

# Chlorzoxazone Tablets, USP

**500 mg**



**Rx Only**

**Each Tablet Contains:**  
Chlorzoxazone, USP .....500 mg

This is a bulk package. Dispense contents with a child-resistant closure (as required) and in a tight container as defined in the USP/NF.

**Keep this and all medications out of the reach of children.**

**Usual Dosage:** See package insert for full prescribing information.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Keep tightly closed.

Manufactured for:  
**Cameron Pharmaceuticals**  
Louisville, KY 40245, USA

Manufactured by:  
**Novitium Pharma LLC**  
70 Lake Drive, East Windsor  
New Jersey 08520

**Revised:** 08/2019

LB4155-02



## PRINCIPAL DISPLAY PANEL - 500 Tablet Bottle Label

NDC 42494-427-50

500 Tablets

Chlorzoxazone  
Tablets, USP

500 mg

CAMERON  
PHARMACEUTICALS

Rx Only

NDC 42494-427-50 500 Tablets

# Chlorzoxazone Tablets, USP

**500 mg**



**Rx Only**

**Each Tablet Contains:**  
Chlorzoxazone, USP .....500 mg

This is a bulk package. Dispense contents with a child-resistant closure (as required) and in a tight container as defined in the USP/NF.

**Keep this and all medications out of the reach of children.**

**Usual Dosage:** See package insert for full prescribing information.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Keep tightly closed.

Manufactured for:  
**Cameron Pharmaceuticals**  
Louisville, KY 40245, USA

Manufactured by:  
**Novitium Pharma LLC**  
70 Lake Drive, East Windsor  
New Jersey 08520

**Revised:** 08/2019

LB4156-02



## CHLORZOXAZONE

chlorzoxazone tablet

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:42494-426
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Chlorzoxazone (UNII: H0DE420U8G) (Chlorzoxazone - UNII:H0DE420U8G)	Chlorzoxazone	500 mg

### Product Characteristics

Color	WHITE	Score	2 pieces
Shape	ROUND	Size	17mm
Flavor		Imprint Code	N85
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42494-426-10	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/27/2019	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA212254	11/27/2019	

## CHLORZOXAZONE

chlorzoxazone tablet

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:42494-427
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Chlorzoxazone (UNII: H0DE420U8G) (Chlorzoxazone - UNII:H0DE420U8G)	Chlorzoxazone	500 mg

**Product Characteristics**

<b>Color</b>	WHITE	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	N85
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42494-427-50	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/27/2019	03/24/2020

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA212254	11/27/2019	03/24/2020

**Labeler** - Cameron Pharmaceuticals (078371442)**Establishment**

Name	Address	ID/FEI	Business Operations
Novitium Pharma, LLC		080301870	MANUFACTURE(42494-426, 42494-427)

Revised: 3/2020

Cameron Pharmaceuticals