

**CORICIDIN HBP COUGH AND COLD- chlorpheniramine maleate,
dextromethorphan hydrochloride tablet
Bayer HealthCare LLC.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Coricidin HBP Cough and Cold UI 1614831

Drug Facts

Active ingredients (in each tablet) Purpose

Chlorpheniramine maleate 4 mg.....Antihistamine
Dextromethorphan hydrobromide 30 mg.....Cough suppressant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves runny nose and sneezing due to the common cold

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Do not use to sedate children.

Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
- cough that occurs with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if cough persists for more than 7 days, tends to recur, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. **Abuse of this product can lead to serious injury.**

Directions

- do not exceed recommended dose
- adults and children 12 years and over: take 1 tablet every 6 hours, not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

Other information

- store between 20° to 25°C (68° to 77°F)
- see end flap for expiration date and lot number

Inactive ingredients croscarmellose sodium, D&C red #27 aluminum lake, FD&C yellow #6 aluminum lake, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, silicon dioxide, stearic acid, talc, titanium dioxide

Questions or comments? 1-800-317-2165 (Mon-Fri 9AM - 5PM EST)

Carton 16

Coricidin[®] HBP
COLD RELIEF FOR PEOPLE WITH HIGH BLOOD PRESSURE*
COUGH & COLD 16 TABLETS
SAFETY SEALED: Do not use if blister is cut or broken.

Coricidin[®] HBP
COLD RELIEF FOR PEOPLE WITH HIGH BLOOD PRESSURE*
COUGH & COLD 16 TABLETS
Chlorpheniramine Maleate - Antihistamine
Dextromethorphan HBr - Cough Suppressant

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Drug Facts
Active ingredients (in each tablet)
Chlorpheniramine maleate 4 mgAntihistamine
Dextromethorphan hydrobromide 30 mg.....Cough suppressant
Purpose
temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
temporarily relieves runny nose and sneezing due to the common cold
Uses
temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
temporarily relieves runny nose and sneezing due to the common cold
Warnings
Do not use
if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
if you have ever had an allergic reaction to this product or any of its ingredients
in children under 12 years of age
Do not use to sedate children
Ask a doctor before use if you have
glaucoma
difficulty in urination due to enlargement of the prostate gland
cough that occurs with excessive phlegm (mucus)
a breathing problem such as emphysema or chronic bronchitis
persistent or chronic cough such as occurs with smoking, asthma, or emphysema
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

Drug Facts (continued)
When using this product
excitability may occur, especially in children
marked drowsiness may occur
avoid alcoholic beverages
alcohol, sedatives and tranquilizers may increase drowsiness
use caution when driving a motor vehicle or operating machinery
Stop use and ask a doctor if cough persists for more than 7 days, tends to recur, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
do not take more than recommended dose
adults and children 12 years and over: take 1 tablet with water every 6 hours. Do not exceed 4 tablets in 24 hours or as directed by a doctor.
children under 12 years of age: do not use

Other information
store between 20° to 25°C (68° to 77°F)
Inactive ingredients croscarmellose sodium, D&C red no. 27 aluminum lake, FD&C yellow no. 6 aluminum lake, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

Questions or comments?
1-800-317-2165 (Mon-Fri 9AM - 5PM EST)

Actual Size

LOT/EXP

Bayer
© 2020 Bayer
Whispery, NJ 07981
Pat. pending

Coricidin®

HBP

Decongestant-Free Cold

Symptom Relief for

People with **High Blood Pressure**

Cold and Cough

Dextromethorphan HBr-

Cough Suppressant

Chlorpheniramine Maleate-

Antihistamine

Relieves:

- Cough
- Runny nose
- Sneezing

16 TABLETS

CORICIDIN HBP COUGH AND COLD

chlorpheniramine maleate, dextromethorphan hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11523-0008
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	4 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	

MAGNESIUM STEARATE (UNII: 70097M6I30)	
TALC (UNII: 7SEV7J4R1U)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Product Characteristics

Color	red	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	CC
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11523-0008-1	2 in 1 CARTON	10/23/2020	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	10/23/2020	

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 5/2023

Bayer HealthCare LLC.