

CLOBETASOL PROPIONATE- clobetasol propionate ointment

Preferred Pharmaceuticals, Inc.

Reference Label Set Id: 48707910-7f65-465b-810e-238f985fd63e

Reference Label Set Id: d43c1251-806a-4cea-8efb-05d3af2ccd43

Clobetasol Propionate Gel, 0.05%

Clobetasol Propionate Cream USP, 0.05%

Clobetasol Propionate Ointment USP, 0.05%

Rx Only


FOR DERMATOLOGIC USE ONLY

NOT FOR OPHTHALMIC, ORAL OR INTRAVAGINAL USE

DESCRIPTION

Clobetasol propionate gel, cream and ointment contain the active compound clobetasol propionate, a synthetic corticosteroid, for topical dermatologic use. Clobetasol, an analog of prednisolone, has a high degree of glucocorticoid activity and a slight degree of mineralocorticoid activity.

Clobetasol propionate is a white to cream-colored crystalline powder insoluble in water. Chemically, it is 21-chloro-9-fluoro-11 β ,17-dihydroxy-16 β -methylpregna-1,4-diene-3,20-dione 17-propionate, and it has the following structural formula:



C₂₅H₃₂ClFO₅

Molecular Weight: 467

Each gram of the 0.05% gel contains 0.5 mg clobetasol propionate in a base of carbomer 934P, propylene glycol, purified water, and sodium hydroxide.

Each gram of the 0.05% cream contains clobetasol propionate 0.5 mg in a cream base of cetyl alcohol, chlorocresol, citric acid, glyceryl monostearate, glyceryl stearate/polyethylene glycol 100 stearate, propylene glycol, purified water, sodium citrate, stearyl alcohol, and white wax.

Each gram of the 0.05% ointment contains clobetasol propionate 0.5 mg in a base of propylene glycol, sorbitan sesquioleate, and white petrolatum.

CLINICAL PHARMACOLOGY

Like other topical corticosteroids, clobetasol propionate has anti-inflammatory, antipruritic, and vasoconstrictive properties. The mechanism of the anti-inflammatory activity of the topical steroids, in general, is unclear. However, corticosteroids are thought to act by the induction of phospholipase A₂ inhibitory proteins, collectively called lipocortins. It is postulated that these proteins control the biosynthesis of potent mediators of inflammation such as prostaglandins and leukotrienes by inhibiting the release of their common precursor, arachidonic acid. Arachidonic acid is released from membrane

phospholipids by phospholipase A₂.

Pharmacokinetics

The extent of percutaneous absorption of topical corticosteroids is determined by many factors, including the vehicle and the integrity of the epidermal barrier. Occlusive dressings with hydrocortisone for up to 24 hours has not been demonstrated to increase penetration; however, occlusion of hydrocortisone for 96 hours markedly enhances penetration. Topical corticosteroids can be absorbed from normal intact skin. Inflammation and/or other disease processes in the skin may increase percutaneous absorption. Greater absorption was observed for the clobetasol propionate gel formulation as compared to the cream formulation in *in vitro* human skin penetration studies. Studies performed with clobetasol propionate gel, cream and ointment indicate that they are in the super-high range of potency as compared with other topical corticosteroids.

INDICATIONS AND USAGE

Clobetasol propionate gel, cream and ointment are super-high potency corticosteroid formulations indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid responsive dermatoses. Treatment beyond 2 consecutive weeks is not recommended, and the total dosage should not exceed 50 g per week because of the potential for the drug to suppress the hypothalamic-pituitary-adrenal (HPA) axis. Use in pediatric patients under 12 years of age is not recommended. As with other highly active corticosteroids, therapy should be discontinued when control has been achieved. If no improvement is seen within 2 weeks, reassessment of the diagnosis may be necessary.

CONTRAINDICATIONS

Clobetasol propionate gel, cream and ointment are contraindicated in those patients with a history of hypersensitivity to any of the components of the preparations.

PRECAUTIONS

General

Clobetasol propionate is a highly potent topical corticosteroid that has been shown to suppress the HPA axis at doses as low as 2 g per day.

Systemic absorption of topical corticosteroids can produce reversible HPA axis suppression with the potential for glucocorticosteroid insufficiency after withdrawal from treatment. Manifestations of Cushing's syndrome, hyperglycemia, and glucosuria can also be produced in some patients by systemic absorption of topical corticosteroids while on therapy.

Patients applying a topical steroid to a large surface area or to areas under occlusion should be evaluated periodically for evidence of HPA axis suppression. This may be done by using the ACTH stimulation, A.M. plasma cortisol, and urinary free cortisol tests. Patients receiving superpotent corticosteroids should not be treated for more than 2 weeks at a time and only small areas should be treated at any one time due to the increased risk of HPA suppression. If HPA axis suppression is noted, an attempt should be made to withdraw the drug, to reduce the frequency of application, or to substitute a less potent corticosteroid. Recovery of HPA axis function is generally prompt upon discontinuation of topical corticosteroids. Infrequently, signs and symptoms of glucocorticosteroid insufficiency may occur requiring supplemental systemic corticosteroids. For information on systemic supplementation, see prescribing information for those products.

Pediatric patients may be more susceptible to systemic toxicity from equivalent doses due to their larger skin surface to body mass ratios (see **PRECAUTIONS: Pediatric Use**).

If irritation develops, clobetasol propionate should be discontinued and appropriate therapy instituted.

Allergic contact dermatitis with corticosteroids is usually diagnosed by observing *failure to heal* rather than noting a clinical exacerbation as with most topical products not containing corticosteroids. Such an observation should be corroborated with appropriate diagnostic patch testing. If concomitant skin infections are present or develop, an appropriate antifungal or antibacterial agent should be used. If a favorable response does not occur promptly, use of clobetasol propionate should be discontinued until the infection has been adequately controlled.

Clobetasol propionate gel, cream, and ointment should not be used in the treatment of rosacea or perioral dermatitis and it should not be used on the face, groin, or axillae.

Information for Patients

Patients using topical corticosteroids should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes.
2. This medication should not be used for any disorder other than that for which it was prescribed.
3. The treated skin area should not be bandaged, otherwise covered or wrapped, so as to be occlusive unless directed by the physician.
4. Patients should report any signs of local adverse reactions to the physician.
5. Patients should inform their physicians that they are using clobetasol propionate if surgery is contemplated.

Laboratory Tests

The following tests may be helpful in evaluating patients for HPA axis suppression: ACTH stimulation test, A.M. plasma cortisol test, Urinary free cortisol test.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term animal studies have not been performed to evaluate the carcinogenic potential of clobetasol propionate. Studies in the rat following subcutaneous administration at dosage levels up to 50 mcg/kg per day revealed that the females exhibited an increase in the number of resorbed embryos and a decrease in the number of living fetuses at the highest dose. Clobetasol propionate was non-mutagenic in three different test systems: the Ames test, the *Saccharomyces cerevisiae* gene conversion assay, and the *E. coli* B WP2 fluctuation test.

Pregnancy

Teratogenic Effects

Pregnancy Category C

Corticosteroids have been shown to be teratogenic in laboratory animals when administered systemically at relatively low dosage levels. Some corticosteroids have been shown to be teratogenic after dermal application to laboratory animals. Clobetasol propionate has not been tested for teratogenicity when applied topically; however, it is absorbed percutaneously, and when administered subcutaneously it was a significant teratogen in both the rabbit and mouse. Clobetasol propionate has greater teratogenic potential than steroids that are less potent. Teratogenicity studies in mice using the subcutaneous route resulted in fetotoxicity at the highest dose tested (1 mg/kg) and teratogenicity at all dose levels tested down to 0.03 mg/kg. These doses are approximately 1.4 and 0.04 times, respectively, the human topical dose of clobetasol propionate gel, cream and ointment. Abnormalities seen included cleft palate and skeletal abnormalities. In rabbits, clobetasol propionate was teratogenic at doses of 3 and 10 mcg/kg. These doses are approximately 0.02 and 0.05 times, respectively, the human topical dose of clobetasol propionate gel, cream and ointment. Abnormalities seen included cleft palate, cranioschisis, and other skeletal abnormalities. There are no adequate and well-controlled studies of the

teratogenic potential of clobetasol propionate in pregnant women. Clobetasol propionate gel, cream or ointment should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers

Systemically administered corticosteroids appear in human milk and could suppress growth, interfere with endogenous corticosteroid production, or cause other untoward effects. It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in human milk. Because many drugs are excreted in human milk, caution should be exercised when either clobetasol propionate gel, cream or ointment is administered to a nursing woman.

Pediatric Use

Safety and effectiveness of clobetasol propionate gel, cream, and ointment in pediatric patients have not been established. Use in children under 12 years of age is not recommended. Because of a higher ratio of skin surface area to body mass, pediatric patients are at a greater risk than adults of HPA axis suppression and Cushing's syndrome when they are treated with topical corticosteroids. They are therefore also at greater risk of adrenal insufficiency during or after withdrawal of treatment. Adverse effects including striae have been reported with inappropriate use of topical corticosteroids in infants and children (see PRECAUTIONS).

HPA axis suppression, Cushing's syndrome, linear growth retardation, delayed weight gain and intracranial hypertension have been reported in children receiving topical corticosteroids. Manifestations of adrenal suppression in children include low plasma cortisol levels, and an absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema.

Geriatric Use

Clinical studies of clobetasol propionate drug products in US clinical trials did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious.

ADVERSE REACTIONS

In a controlled clinical trial with clobetasol propionate gel, the only reported adverse reaction that was considered to be drug related was a report of burning sensation (1.8% of treated patients). In controlled clinical trials, the most frequent adverse reactions reported for clobetasol propionate cream were burning and stinging sensation in 1% of treated patients. Less frequent adverse reactions were itching, skin atrophy, and cracking and fissuring of the skin. In controlled clinical trials, the most frequent adverse events reported for clobetasol propionate ointment were burning sensation, irritation, and itching in 0.5% of treated patients. Less frequent adverse reactions were stinging, cracking, erythema, folliculitis, numbness of fingers, skin atrophy, and telangiectasia.

Cushing's syndrome has been reported in infants and adults as a result of prolonged use of topical clobetasol propionate formulations. The following additional local adverse reactions have been reported with topical corticosteroids, and they may occur more frequently with the use of occlusive dressings and higher potency corticosteroids. These reactions are listed in an approximately decreasing order of occurrence: dryness, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, secondary infection, irritation, striae, and miliaria.

OVERDOSAGE

Topically applied clobetasol propionate gel, cream or ointment can be absorbed in sufficient amounts to

produce systemic effects (see **PRECAUTIONS**).

DOSAGE AND ADMINISTRATION

Apply a thin layer of clobetasol propionate gel, cream or ointment to the affected skin areas twice daily and rub in gently and completely. (See **INDICATIONS AND USAGE**.)

Clobetasol propionate gel, cream and ointment are super-high potency topical corticosteroids; therefore, **treatment should be limited to 2 consecutive weeks, and amounts greater than 50 g per week should not be used.**

As with other highly active corticosteroids, therapy should be discontinued when control has been achieved. If no improvement is seen within 2 weeks, reassessment of diagnosis may be necessary.

Clobetasol propionate gel, cream or ointment should not be used with occlusive dressings.

HOW SUPPLIED

Clobetasol Propionate Ointment USP, 0.05% is supplied in tamper-evident tubes:

15g tubes – 68788-9802-1

30g tubes – 68788-9802-3

Store at 20°-25°C (68°-77°F) [see USP Controlled Room Temperature]. **DO NOT REFRIGERATE**

Mfd. by: Taro Pharmaceuticals Inc., Brampton, Ontario, Canada L6T 1C1

Dist by: **Taro Pharmaceuticals U.S.A., Inc.**, Hawthorne, NY 10532

Revised: April, 2014

PK-6540-1

352

Relabeled By: Preferred Pharmaceuticals Inc.

PRINCIPAL DISPLAY PANEL - 15 g Ointment Tube Carton

NDC 68788-9802-1

**Clobetasol Propionate
Ointment USP, 0.05%**

FOR EXTERNAL USE ONLY.
NOT FOR OPHTHALMIC USE.

Keep this and all medications out of the reach of children.

15 g

Rx only

TARO

Relabeled By: Preferred Pharmaceuticals Inc.

Clobetasol Propionate Ointment USP, 0.05%

Generic for Temovate

Each gram contains: Clobetasol propionate 0.5mg in a base of propylene glycol, sorbitan sesquioleate and white petrolatum.

Pkg Size: Exp Date:
Lot#: Batch#:

Ins:

Mfg: Taro Pharmaceuticals, Ontario, Canada
Prod#:

Warning

For external use only. Not for ophthalmic use. Rx Only. Keep this and all medications out of the reach of children. Store at 20°-25°C (68°-77°F). See USP Controlled Room Temperature. Do not refrigerate. See package insert for full prescribing information.

PREFERRED
Pharmaceuticals, Inc.
The Physician's Solution. Anaheim, Ca

CAUTION: Federal law PROHIBITS transfer of this drug to any person other than the patient for whom it was prescribed.

Clobetasol Propionate Ointment USP, 0.05%
Qty: Ins:
Lot#: Bat#:
Prod# (NDC):

Log

Clobetasol Propionate Ointment USP, 0.05%
Qty: Ins:
Lot#: Bat#:
Prod# (NDC):

Chart

Clobetasol Propionate Ointment USP, 0.05%
Qty: Ins:
Insurance NDC:
Lot#: Bat#:

Billing

Clobetasol Propionate Ointment USP, 0.05%
Qty: Ins:
Lot#: Bat#:
Prod# (NDC):

Patient



Directions English

Apply externally _____ times a day.
For external use only.
Use as directed by your doctor

Instrucciones Espanol:

Aplique externamente _____ veces al dia.
Para externamente aplique
Uso según lo dirigido por su doctor

CLOBETASOL PROPIONATE

clobetasol propionate ointment

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:68788-9802(NDC:51672-1259)
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Clobetasol Propionate (UNII: 779619577M) (Clobetasol - UNII:ADN79D536H)	Clobetasol Propionate	0.5 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
propylene glycol (UNII: 6DC9Q167V3)	
sorbitan sesquioleate (UNII: 0W8RRI5W5A)	
petrolatum (UNII: 4T6H12BN9U)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68788-9802-1	15 g in 1 TUBE; Type 0: Not a Combination Product	04/23/2012	
2	NDC:68788-9802-3	30 g in 1 TUBE; Type 0: Not a Combination Product	04/23/2012	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074248	04/23/2012	

Labeler - Preferred Pharmaceuticals, Inc. (791119022)

Registrant - Preferred Pharmaceuticals, Inc. (791119022)

Establishment

Name	Address	ID/FEI	Business Operations
Preferred Pharmaceuticals, Inc.		79 1119022	RELABEL(68788-9802)

Revised: 2/2017

Preferred Pharmaceuticals, Inc.