

**REGULAR STRENGTH ANTACID/ANTIGAS- aluminum hydroxide, magnesium hydroxide, dimethicone suspension**  
**RUGBY LABORATORIES**

-----  
**rug mlx mnt**

**Active ingredients (in each 10 mL dose)**

Aluminum hydroxide 400 mg (equivalent to dried gel, USP)

Magnesium hydroxide 400 mg

Simethicone 40mg

**Purposes**

Antacid

Antigas

**Uses**

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

**Warnings**

**Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet

**Ask a doctor or pharmacist before use if you are** taking a prescription drug.

Antacids may interact with certain prescription drugs.

**Stop use and ask a doctor if** symptoms last more than 2 weeks

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Directions**

- shake well before each use
- adults and children 12 years and older: take 10 mL to 20 mL (1-2 doses) four times a day or as directed by a doctor

- do not exceed 80 mL (8 doses) in a 24 hour period or use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor

### **Other information**

- each 10 mL dose contains: **magnesium 165 mg, sodium 6 mg**
- store at room temperature
- protect from freezing
- keep tightly closed

### **Inactive ingredients**

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

### **Questions or comments?**

**1-800-540-3765**

### **package Label**



NDC 0536-1317-83

Regular Strength

# Antacid/ Antigas Liquid

Alumina, Magnesia,  
and Simethicone  
Oral Suspension USP

Fast Relief of:  
Acid Indigestion  
Heartburn  
Sour Stomach  
Pressure and Bloating

Cool Mint Flavor

219-11212-3  
REV GC0820

12 FL OZ (355 mL)

## Drug Facts

TAMPER EVIDENT: DO NOT USE IF BREAKAWAY  
BAND ON BOTTLE CAP IS MISSING OR BROKEN.

### Active ingredients (in each 10 mL dose)

Aluminum hydroxide 400 mg (equivalent to dried gel USP)	Antacid
Magnesium hydroxide 400 mg	Antacid
Simethicone 40 mg	Antigas

### Purposes

**Uses** relieves • heartburn • sour stomach  
• acid indigestion • the symptoms referred to as gas

### Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks  
If pregnant or breast-feeding, ask a health professional before use.  
Keep out of reach of children.

### Directions

- shake well before each use
- adults and children 12 years and older: take 10 mL to 20 mL (1-2 doses) four times a day or as directed by a doctor
- do not exceed 80 mL (8 doses) in a 24 hour period or use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor

### Other information

- each 10 mL dose contains: magnesium 165 mg, sodium 6 mg
- store at room temperature • protect from freezing
- keep tightly closed

**Inactive ingredients** benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

Distributed by:  
RUGBY LABORATORIES  
17177 N Laurel Park Dr, Suite 233  
Livonia, MI 48152  
www.rugbylaboratories.com

Rev. 08/20 R-99 Re-order No. 371063



919-11212-3

## REGULAR STRENGTH ANTACID/ANTIGAS

aluminum hydroxide, magnesium hydroxide, dimethicone suspension

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0536-1317
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE -	ALUMINUM	400 mg

UNII:5QB0T2IUN0)	HYDROXIDE	in 10 mL
<b>MAGNESIUM HYDROXIDE</b> (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	400 mg in 10 mL
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	40 mg in 10 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>BUTYLPARABEN</b> (UNII: 3QPI1U3FV8)	
<b>CARBOXYMETHYLCELLULOSE SODIUM</b> (UNII: K679OBS311)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	

## Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	MINT	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0536-1317-83	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2020	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	08/01/2020	

**Labeler** - RUGBY LABORATORIES (079246066)

**Registrant** - GCP Laboratories (965480861)

## Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(0536-1317)

