

**STOMACH RELIEF MAXIMUM STRENGTH- bismuth subsalicylate suspension**  
**Demoulas Super Markets Inc**

-----  
**Stomach Relief Maximum Strength**

**Drug Facts**

**Active ingredient (In each 30 mL dose cup or 2 tablespoons)**

Bismuth subsalicylate 1050 mg

**Purposes**

Bismuth subsalicylate ..... Upset stomach reliever  
and antidiarrheal

**Uses**

relieves ■ travelers' diarrhea ■ diarrhea  
■ upset stomach due to overindulgence of food and  
drink including: ■ heart burn ■ indigestion  
■ nausea ■ gas ■ belching

**Warnings**

**Reye's syndrome:** Children and teenagers who have or  
are recovering from chicken pox or flu-like symptoms  
should not use this product. When using this product, if  
changes in behavior with nausea and vomiting occur,  
consult a doctor because these symptoms could be an  
early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

■ allergic to salicylates (including aspirin)  
■ taking other salicylate products

**Do not use** if you have ■ an ulcer  
■ a bleeding problem ■ black or bloody stool

**Ask a doctor before use if you have**

■ fever ■ mucus in stool

**Ask a doctor or pharmacist before use if you are** taking

any drug for ■ anticoagulation (thinning of the blood)  
■ diabetes ■ gout ■ arthritis

**When using this product** a temporary, but harmless,  
darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if**

■ diarrhea lasts more than 2 days  
■ symptoms get worse or last more than 2 days  
■ ringing in the ears or loss of hearing occurs

**If pregnant or breast feeding,** ask health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison  
Control Center right away.

- Directions.** ■ Shake well before use  
■ use dose cup or tablespoon (TBSP)  
■ adults and children 12 years and over: 1 dose (30 mL or 2 TBSP) every 1 hour as needed  
■ do not exceed 4 doses (120 mL or 8 TBSP) in 24 hours  
■ use until diarrhea stops but not more than 2 days  
■ children under 12 years: ask a doctor  
■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information**

- **each 30 mL dose cup contains:**  
■ sodium 13 mg, salicylate 455 mg  
■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C) ■ low sodium

**Inactive ingredients**

benzoic acid, D&C red # 22, D&C red # 28, flavor, purified water, saccharin sodium, salicylic acid, sodium salicylate, xanthan gum

**Questions?** 1-877-798-5944

**Principal Display Panel**

**MARKET**

**BASKET®**

**"MORE FOR YOUR DOLLAR"**

\*COMPARE TO

The active ingredient in  
PEPTO-BISMOL™ Maximum Strength

**Stomach Relief↓**

**MAXIMUM STRENGTH**

Bismuth Subsalicylate  
Antidiarrheal / Upset Stomach Reliever

**5 Symptom Relief of:**

- Nausea
- Upset Stomach
- Heartburn
- Diarrhea
- Indigestion

**8 FL OZ (237 mL)**

**TAMPER EVIDENT: Do not use if imprinted shrinkband is missing or broken**

\*This product is not manufactured or distributed by Procter & Gamble, Inc., the distributor of Pepto-Bismol™.

**DISTRIBUTED BY**

**Made in USA**

0 49705 73076 8

**PEEL**

**HERE**

↓ **Principal Display Label** ↓

<p><b>TAMPER EVIDENT:</b> Do not use if imprinted shrinkband is missing or broken</p>	 <p><b>Stomach Relief</b>  <b>MAXIMUM STRENGTH</b>                  Bismuth Subsalicylate                  Antidiarrheal / Upset Stomach Reliever                  5 Symptom Relief of:                  • Nausea • Upset stomach                  • Heartburn • Diarrhea                  • Indigestion</p> <p>8 FL OZ (237 mL)</p>	<p><b>Drug Facts (continued)</b></p> <p>Ask a doctor or pharmacist before use if you are taking any drug for</p> <ul style="list-style-type: none"> <li>■ anticoagulation (thinning of the blood)</li> <li>■ diabetes</li> <li>■ gout</li> <li>■ arthritis</li> </ul> <p>When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> <li>■ diarrhea lasts more than 2 days</li> <li>■ symptoms get worse or last more than 2 days</li> <li>■ ringing in the ears or loss of hearing occurs</li> </ul> <p>If pregnant or breast feeding, ask health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><small>*This product is not manufactured or distributed by Procter &amp; Gamble, Inc., the distributor of Pepto-Bismol™.</small></p> <p>DISTRIBUTED BY                  DEMOULAS SUPERMARKETS INC.                  875 EAST STREET                  TEWKSBURY, MA 01876</p> <p style="text-align: right;">Made in USA                  LR-089 REV01</p>
<p><b>Drug Facts</b></p> <p><b>Active ingredient</b> <span style="float: right;"><b>Purposes</b></span>                  (In each 30 mL dose cup or 2 tablespoons)                  Bismuth subsalicylate 1050 mg ...Upset stomach reliever and antidiarrheal</p> <p><b>Uses</b>                  relieves ■ travelers' diarrhea ■ diarrhea                  ■ upset stomach due to overindulgence of food and drink including: ■ heart burn ■ indigestion                  ■ nausea ■ gas ■ belching</p> <p><b>Warnings</b>  <b>Reye's syndrome:</b> Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  <b>Allergy alert: Contains salicylate. Do not take if you are</b>                  ■ allergic to salicylates (including aspirin)                  ■ taking other salicylate products</p> <p><b>Do not use</b> if you have ■ an ulcer                  ■ a bleeding problem ■ black or bloody stool</p> <p><b>Ask a doctor before use if you have</b>                  ■ fever ■ mucus in stool</p>		 0 49705 73076 8

<p><b>Drug Facts (continued)</b></p> <p><b>Directions</b> ■ Shake well before use</p> <ul style="list-style-type: none"> <li>■ use dose cup or tablespoon (TBSP)</li> <li>■ adults and children 12 years and over: 1 dose (30 mL or 2 TBSP) every 1 hour as needed</li> <li>■ do not exceed 4 doses (120 mL or 8 TBSP) in 24 hours</li> <li>■ use until diarrhea stops but not more than 2 days</li> <li>■ children under 12 years: ask a doctor</li> <li>■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ each 30 mL dose cup contains: sodium 13 mg, salicylate 455 mg</li> <li>■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C) ■ low sodium</li> </ul> <p><b>Inactive Ingredients</b>                  benzoic acid, D&amp;C red # 22, D&amp;C red # 28, flavor, purified water, saccharin sodium, salicylic acid, sodium salicylate, xanthan gum</p> <p><b>Questions?</b> 1-877-798-5944</p>
---

Stomach Relief, Maximum Strength by Market Basket®

res

**STOMACH RELIEF MAXIMUM STRENGTH**

bismuth subsalicylate suspension

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:53942-558
---------------------	----------------	---------------------------	---------------

<b>Route of Administration</b>	ORAL
--------------------------------	------

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	1050 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ)	
<b>SODIUM SALICYLATE</b> (UNII: WQ1H85SYP)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Product Characteristics

<b>Color</b>	pink	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	MINT	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53942-558-26	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2018	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	10/01/2018	

**Labeler** - Demoulas Super Markets Inc (007869647)

**Registrant** - AptaPharma Inc. (790523323)

### Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

AptaPharma Inc.

790523323

manufacture(53942-558)

Revised: 12/2023

Demoulas Super Markets Inc