

**ACETAMINOPHEN, CHLORPHENIRAMINE MALEATE, PHENYLEPHRINE HCL-**  
**acetaminophen, chlorpheniramine maleate, phenylephrine hcl tablet**  
**OPMX LLC**

-----  
**Coldtac Ultra tabs**

**(Acetaminophen 500mg, Chlorpheniramine Maleate 4mg, Phenylephrine HCl 10mg Tablet)**

***Drug Facts***

***Active ingredients***

***Purpose***

---

***Active ingredients (in each caplet)***

Acetaminophen 500 mg  
Chlorpheniramine Maleate 4 mg  
Phenylephrine HCl 10 mg

---

***Purpose***

Pain reliever  
Antihistamine  
Nasal decongestant

---

***Uses***

Temporarily relieves these symptoms of hay fever or other respiratory allergies:

- headache
- nasal congestion
- sinus congestion & pressure
- runny nose and sneezing
- minor aches & pain

Temporarily relieves these additional symptoms of hay fever:

- itching of the nose or throat
- itchy, watery eyes
- helps clear nasal passages
- helps decongest sinus opening and passages

***Warnings***

**Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if you take :

- more than 6 tablets in 24 hours, which is the maximum daily amount for this product
- with other drugs contains acetaminophen
- 3 or more alcoholic drinks every day while using this product.

**Allergy alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:**

if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult doctor promptly

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen,
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist  
If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI
- if you have ever had an allergic reaction to this product or any of its ingredients.
- to make a child sleepy

**Ask a doctor before use if you have:**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- glaucoma
- diabetes
- trouble urinating due to enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

**Ask a doctor or pharmacist if you are:**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product:**

- do not use more than directed
- excitability may occur, especially in children
- drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcohol drinks
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if:**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days

- fever gets worse or last more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- Do not take more than directed - See overdose warning
- Swallow whole; do not crush, chew or dissolve
- Do not exceed 6 tablets per 24 hours

Adults and children 12 years of age & over	1 tablet with water every 4-6 hours as needed
Children 4 to 12 years of age	Ask a doctor
Children under 4 years of age	Do not use

**Other information**

- Store at room temperature between 20-25°C (68-77°F)
- Tamper evident: Do not use if pouch is torn, broken or shows any sing of tampering

**Inactive ingredients**

magnesium stearate, povidone, silicon dioxide, sodium benzoate, sodium starch glycolate, starch, talc

**Questions or comments?**

Call: (619) 600-5632 (Mon-Fri 9am-5pm EST) or <https://www.opmx.us>

**Coldtac Ultra**

NDC 69729-122-06



Rápido Alivio a las Molestias de la Gripe y Resfriado

- Dolor de Cabeza
- Estornudos
- Congestión Nasal
- Fiebre
- Escurreniento Nasal
- Dolor de Garganta
- Cuerpo Cortado
- Ojos Llorosos

INSTRUCCIONES EN ESPAÑOL EN EL INTERIOR DE LA CAJA.

COLDTAC ULTRA

Tablets /Tabletas

NDC 69729-122-06

COLDTAC ULTRA



Acetaminophen 500 mg  
Chlorpheniramine Maleate 4 mg  
Phenylephrine HCl 10 mg

Effective and Quick Relief from Cold & Flu Symptoms

- Headache
- Fever
- Body Aches
- Sneezing
- Runny Nose
- Watery Eyes
- Nasal Congestion
- Sore Throat



6 Tablets  
3 Packs of 2 Tablets

COLDTAC ULTRA  
Tablets /Tabletas

READ AND KEEP OUTER CARTON FOR COMPLETE PRODUCT INFORMATION

Exclusively distributed by:

OPMX  
Chula Vista, CA 91910  
Phone: 619-600-5632



Made for:



San Diego, CA



8 10053 18125 6

REV. 08/2023 ITEM # RPK130

LOT

EXP. DATE

contains an NDA, ask a doctor or pharmacist before taking this product. ■ If you have ever had an allergic reaction to this product or any of its ingredients ■ to make a child sleep. ■ Ask a doctor before use if you have: ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis. ■ Ask a doctor or a pharmacist before use if you are: ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers. ■ When using this product: ■ do not use more than directed. ■ excitability may occur, especially in children ■ drowsiness may occur ■ alcohol, sedatives and tranquilizers may increase drowsiness. ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery. ■ Stop use and ask a doctor if: ■ nervousness, dizziness, or sleeplessness occur ■ pain or nasal congestion gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur. These could be signs of a serious condition. ■ If pregnant or breast-feeding: ask a health professional before use. ■ Keep out of reach of children. ■ Overdose warning: taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
■ Do not take more than directed  
■ See Overdose warning  
■ Swallow whole; do not crush, chew or dissolve  
■ Do not exceed 6 tablets per 24 hours

Children 4 to 12 years of age	Ask a doctor
Children under 4 years of age	Do not use
Adults and children 12 years of age & over	1 tablet with water every 4-6 hours as needed

**Other information**  
■ Store at room temperature between 20°-25°C (68°-77°F). ■ Avoid excessive heat and humidity. ■ Tamper Evident: do not use if package or blister unit is open or shows any sign of tampering.

**Inactive ingredients** magnesium stearate, povidone, silicon dioxide, sodium benzoate, sodium starch glycolate, starch, talc.

**Questions or comments?** Call: (619) 600-5632 (Mon-Fri 9am-5pm EST) or <https://www.opmx.us>







Inactive Ingredients	
Ingredient Name	Strength
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I3O)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	

Product Characteristics			
<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND (BICONVEX)	<b>Size</b>	13mm
<b>Flavor</b>		<b>Imprint Code</b>	S78
<b>Contains</b>			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69729-122-27	2 in 1 POUCH; Type 0: Not a Combination Product	08/26/2019	10/24/2022
2	NDC:69729-122-37	12 in 1 POUCH; Type 0: Not a Combination Product	08/26/2019	
3	NDC:69729-122-38	144 in 1 POUCH; Type 0: Not a Combination Product	08/26/2019	
4	NDC:69729-122-06	3 in 1 CARTON	09/04/2023	
4		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/26/2019	

**Labeler -** OPMX LLC (029918743)