

**ACETAMINOPHEN- acetaminophen tablet
WALGREEN CO.**

**Rapid Release Gelcaps
EXTRA STRENGTH
Acetaminophen Gelcaps USP, 500 mg
Pain reliever; Fever reducer
Aspirin free**

Active ingredient (in each Gelcap)

Acetaminophen, USP 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- backache
- minor pain of arthritis
- toothache
- the common cold
- premenstrual and menstrual cramps
- temporarily reduces fever

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

liver disease.

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of the reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Overdose warning

In case of accidental overdose, get medical help or contact a Poison Control Center right away (1-800- 222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed (see overdose warning)

adults and children 12 years and over

- take 2 gelcaps every 6 hours while symptoms last
- do not take more than 6 gelcaps in 24 hours, unless directed by a doctor
- do not take more than 10 days unless directed by a doctor

children under 12 years

- ask a doctor

Other information

- store at 20°-25°C (68°-77°F). See USP Controlled Room Temperature
- avoid high humidity
- see end panel for lot number and expiration date

Inactive Ingredients

ammonium hydroxide, black iron oxide, colloidal silicon dioxide, croscarmellose sodium, D&C red #33, FD&C blue #1, FD&C red #40, FD&C yellow #6, gelatin, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, hypromellose, iron oxide red, isopropyl alcohol, n-butyl alcohol, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glaze, stearic acid, titanium dioxide, yellow iron Oxide.

Questions or comments ?

call **1-877-770-3183** Mon-Fri 8:00 AM EST to 5:00 PM PST.

PDP

09160200002
01850000002

IMPORTANT: READ ALL DIRECTIONS CAREFULLY. KEEP THIS CARD IN THE BOX FOR INFORMATION.
Always follow the directions on the label and do not take more than the amount on the label.

Drug Facts

Active ingredient (in each tablet): acetaminophen USP 500 mg

Other ingredients: croscarmellose sodium, hydroxypropyl cellulose, polyethylene glycol, polyethylene glycol 400, polyethylene glycol 600, polyethylene glycol 800, polyethylene glycol 1000, polyethylene glycol 1500, polyethylene glycol 2000, polyethylene glycol 3000, polyethylene glycol 4000, polyethylene glycol 6000, polyethylene glycol 8000, polyethylene glycol 10000, polyethylene glycol 15000, polyethylene glycol 20000, polyethylene glycol 30000, polyethylene glycol 40000, polyethylene glycol 60000, polyethylene glycol 80000, polyethylene glycol 100000.

Warnings:

- Do not take if you have ever had an allergic reaction to acetaminophen, aspirin, salicylates, or any other drug containing acetaminophen.
- Do not take if you are taking any of the following drugs: aspirin, ibuprofen, naproxen, or any other drug containing acetaminophen.
- Do not take if you are taking any of the following drugs: aspirin, ibuprofen, naproxen, or any other drug containing acetaminophen.
- Do not take if you are taking any of the following drugs: aspirin, ibuprofen, naproxen, or any other drug containing acetaminophen.
- Do not take if you are taking any of the following drugs: aspirin, ibuprofen, naproxen, or any other drug containing acetaminophen.

200000003810
700000001140



Pain Reliever

ACETAMINOPHEN USP, 500 mg
PAIN RELIEVER / FEVER REDUCER



Pain Reliever

ACETAMINOPHEN USP, 500 mg
PAIN RELIEVER / FEVER REDUCER

Extra Strength

225
MAYO RELEASE
SEALING



Pain Reliever

ACETAMINOPHEN USP, 500 mg
PAIN RELIEVER / FEVER REDUCER

Extra Strength

225
MAYO RELEASE
SEALING



17000001157

COATING FREE AREA

700000004067
200000002502

Drug Facts (continued)

Other information

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-422-4225). Quick medical attention is critical for adults as well as for children when if you do not notice any signs or symptoms.

Directions

Do not take more than directed (see overdose warning) with or without food.

Take 2 tablets every 6 hours with or without food.

Do not take more than 6 tablets in 24 hours unless directed by a doctor.

Do not use for more than 14 days unless directed by a doctor.

Children under 12 years: ask a doctor.

COATING FREE AREA

700000004067
200000002502

Drug Facts (continued)

Other information

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-422-4225). Quick medical attention is critical for adults as well as for children when if you do not notice any signs or symptoms.

Directions

Do not take more than directed (see overdose warning) with or without food.

Take 2 tablets every 6 hours with or without food.

Do not take more than 6 tablets in 24 hours unless directed by a doctor.

Do not use for more than 14 days unless directed by a doctor.

Children under 12 years: ask a doctor.

ASPIRIN FREE

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
©2021 Walgreen Co. MADE IN INDIA



Walgreens

Pain Reliever

ACTAMINOPHEN 500 mg
PAIN RELIEVER / FEVER REDUCER

50
mg tablets

Walgreens

Compare the strength of the active ingredients. See Strengths and Other Information.

Walgreens

Pain Reliever

ACTAMINOPHEN 500 mg
PAIN RELIEVER / FEVER REDUCER

50
mg tablets

Walgreens

Compare the strength of the active ingredients. See Strengths and Other Information.



ITEM 846579 W10583-0722-L

3 11917421598 3

REV-8/22

Lot: _____
Exp. _____

COATING FREE AREA

IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING, AND STOP THE DRUGS FOR IMPORTANT DRUG INTERACTIONS.

DO NOT EXCEED THE MAXIMUM DOSE OF 4000 MG OF ACETAMINOPHEN IN 24 HOURS. EXCEEDING THIS DOSE CAN BE DANGEROUS OR FATAL.

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 500 mg, Pain reliever/fever reducer

Uses

- Temporarily relieves minor aches and pains due to:
 - headache
 - backache
 - muscle aches
 - minor pain of arthritis
 - the common cold
 - premenstrual and menstrual cramps
- Temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: more than 4000 mg of acetaminophen in 24 hours; with other drugs containing acetaminophen; or if more alcohol is drunk every day while using this product.

Other warnings: Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

Do not take

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

← 1 3/4 → ← 1 3/4 →

3

700000003697
200000002190

KEEP OUT OF REACH OF CHILDREN. ALWAYS USE AS DIRECTED.
DO NOT TAKE MORE THAN THE RECOMMENDED DOSE.
DO NOT TAKE THIS DRUG FOR MORE THAN 10 DAYS UNLESS YOUR PHYSICIAN INSTRUCTS YOU TO DO SO.

Drug Facts

Active ingredient (in each tablet)	Purpose
acetaminophen, USP 500 mg; <i>Paracetamol</i>	pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscle aches
 - backache
 - minor pain of arthritis
 - the common cold
 - teething
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Like all medicines, this product can cause side effects. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- more than 6,000 mg of acetaminophen
- or more than 10 tablets every day while using this product.

Some side effects include:

- nausea
- stomach pain
- loss of appetite
- indigestion
- diarrhea
- constipation
- or other symptoms.

Do not use with any other drug containing acetaminophen (paracetamol) or aspirin, unless your doctor instructs you to do so. Do not use with any other drug containing acetaminophen (paracetamol) or aspirin, unless your doctor instructs you to do so. Do not use with any other drug containing acetaminophen (paracetamol) or aspirin, unless your doctor instructs you to do so.

700000003697
200000002190

Drug Facts (cont'd)

Stop use and ask a doctor if:

- Worsening pain occurs or lasts more than 10 days
- Worsening fever occurs or lasts more than 3 days
- Worsening cough
- Worsening or swelling is present

These could be signs of a serious condition. If you experience chest pain, a heart attack, or other serious symptoms, stop use immediately and ask a doctor.

Directions

Do not take more than directed (see warnings). Adults and children 12 and over:

- Take 2 tablets every 4 to 6 hours while awake.
- Do not take more than 8 tablets in 24 hours.
- Do not take more than 4 tablets every day.

Children under 12 years:

- See label for directions.

Other information

- Store at 20° - 25°C (68° - 77°F). See USP Controlled Room Temperature.
- Protect from light and moisture.
- Keep out of reach of children.

Contains: Acetaminophen, USP 500 mg tablets. Each tablet contains 500 mg of acetaminophen, USP 500 mg. Each tablet contains 500 mg of acetaminophen, USP 500 mg.

100 TABLETS

Wetzelgreen's

Pain Reliever

ACETAMINOPHEN, USP 500 mg
PMN RELIEVER / FEVER REDUCER

100 TABLETS

ASPIRIN 83

Drug Facts (cont'd)

Warnings or cautions?

Call 1-877-378-2483 for more information on safety and effectiveness. This product is not intended to diagnose, cure, treat, or prevent any disease.

ASPIRIN 83

Wetzelgreen's

Pain Reliever

ACETAMINOPHEN, USP 500 mg
PMN RELIEVER / FEVER REDUCER

Extra Strength

100 TABLETS

ASPIRIN 83

Lot #	1191721597	Exp.	12/31/2018
Material #	615 081	Date	10/12/2018



ACETAMINOPHEN			
acetaminophen tablet			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-9798
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
Inactive Ingredients			
	Ingredient Name		Strength
	HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)		
	FERROSFERRIC OXIDE (UNII: XM0M87F357)		
	FD&C RED NO. 40 (UNII: WZB9127XOA)		
	FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		

GELATIN (UNII: 2G86QN327L)
HYDROXYMETHYL CELLULOSE (UNII: 273FM27VK1)
SHELLAC (UNII: 46N107B71O)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)
D&C RED NO. 33 (UNII: 9DBA0SBB0L)
HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K)
FERRIC OXIDE RED (UNII: 1K09F3G675)
BUTYL ALCOHOL (UNII: 8PJ61P6TS3)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)
AMMONIA (UNII: 5138Q19F1X)
POVIDONE (UNII: FZ989GH94E)
STARCH, CORN (UNII: O8232NY3SJ)
STEARIC ACID (UNII: 4ELV7Z65AP)
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)
ISOPROPYL ALCOHOL (UNII: ND2M416302)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)

Product Characteristics

Color	gray (Encapsulated with red opaque and blue gray opaque hard gelatin shells)	Score	score with uneven pieces
Shape	OVAL	Size	19mm
Flavor		Imprint Code	G1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9798-24	24 in 1 BOTTLE; Type 0: Not a Combination Product	09/24/2021	
2	NDC:0363-9798-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	09/24/2021	
3	NDC:0363-9798-15	150 in 1 BOTTLE; Type 0: Not a Combination Product	09/24/2022	
4	NDC:0363-9798-21	225 in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2022	
5	NDC:0363-9798-37	375 in 1 BOTTLE; Type 0: Not a Combination Product	06/25/2022	
6	NDC:0363-9798-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/24/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/24/2021	

Labeler - WALGREEN CO. (008965063)

Revised: 1/2024

WALGREEN CO.