

## **TUSSIN ADULT- dextromethorphan hbr, guaifenesin solution**

**Western Family Foods Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **Western Family Adult Tussin Drug Facts**

#### **Active ingredients (in each 10 mL)**

Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 200 mg

#### **Purposes**

Cough suppressant

Expectorant

#### **Uses**

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

#### **Warnings**

##### **Do not use**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

##### **Ask a doctor before use if you have**

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

##### **Stop use and ask a doctor if**

cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

##### **If pregnant or breast-feeding,**

ask a health professional before use.

##### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

#### **Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

| age                                   | dose                |
|---------------------------------------|---------------------|
| adults and children 12 years and over | 10 mL every 4 hours |
| children under 12 years               | do not use          |

### Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

### Inactive ingredients

anhydrous citric acid, FD&C red no. 40, flavor, glycerin, high fructose corn syrup, menthol, propylene glycol, purified water, sodium benzoate, sodium citrate, sucralose

### Principal Display Panel

ADULT

TUSSIN

COUGH & CHEST CONGESTION

DM

COUGH SUPPRESSANT (DEXTROMETHORPHAN HBr)

EXPECTORANT (GUAIFENESIN)

NON-DROWSY

Relieves:

COUGH

MUCUS

Alcohol Free

For Ages 12 & Over

Gluten Free

COMPARE TO ROBITUSSIN® COUGH + CHEST CONGESTION DM active ingredients

4 FL OZ (118 mL)



**ADULT**  
**TUSSIN**

**COUGH & CHEST CONGESTION**

**DM** COUGH SUPPRESSANT  
(DEXTROMETHORPHAN HBr)  
EXPECTORANT  
(GUAIFENESIN)

**NON-DROWSY**

Relieves:  
**COUGH**  
**MUCUS**

**Alcohol Free**

For Ages 12 & Over

**Gluten Free**



COMPARE TO ROBITUSSIN® COUGH + CHEST  
CONGESTION DM active ingredients\*

4 FL OZ  
(118 mL)



**ADULT**  
**TUSSIN**

**COUGH & CHEST CONGESTION**

**DM** COUGH SUPPRESSANT  
(DEXTROMETHORPHAN HBr)  
EXPECTORANT  
(GUAIFENESIN)

**NON-DROWSY**

Relieves:  
**COUGH**  
**MUCUS**

**Alcohol Free**

For Ages 12 & Over


**Gluten Free**



COMPARE TO ROBITUSSIN® COUGH + CHEST  
CONGESTION DM active ingredients\*

4 FL OZ  
(118 mL)





**DM**  
**COUGH & CHEST CONGESTION**  
**TUSSIN**  
**ADULT**

**WESTERN FAMILY**

**Drug Facts**

**Active ingredients**      **Purposes**  
**(in each 10 mL)**  
 Dextromethorphan HBr,  
 USP 20 mg.....Cough suppressant  
 Guaifenesin, USP 200 mg.....Expectorant

**Uses**

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

**Warnings**

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Stop use and ask a doctor** if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age ▶

**Drug Facts (continued)**

| age                                   | dose                |
|---------------------------------------|---------------------|
| adults and children 12 years and over | 10 mL every 4 hours |
| children under 12 years               | do not use          |

**Other information**

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** anhydrous citric acid, FD&C red no. 40, flavor, glycerin, high fructose corn syrup, menthol, propylene glycol, purified water, sodium benzoate, sodium citrate, sucralose

Proudly Distributed by  
 Western Family Foods, Inc.  
 P.O. Box 4057, Portland, OR 97208 U.S.A.  
 Copyright 2014 Marca Registrada  
[www.westernfamily.com](http://www.westernfamily.com)



**DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING**

\*This product is not manufactured or distributed by Pfizer, distributor of Robitussin® Cough + Chest Congestion DM.

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

02245-G-PER





T

LOT NO.

EXP.

: 35926 3M C10

## TUSSIN ADULT

dextromethorphan hbr, guaifenesin solution

### Product Information

|                         |                |                    |               |
|-------------------------|----------------|--------------------|---------------|
| Product Type            | HUMAN OTC DRUG | Item Code (Source) | NDC:55312-359 |
| Route of Administration | ORAL           |                    |               |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength                | Strength           |
|--|----------------------------------|--------------------|
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH)<br>(DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN<br>HYDROBROMIDE | 20 mg<br>in 10 mL  |
| GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)                           | GUAIFENESIN                      | 200 mg<br>in 10 mL |

### Inactive Ingredients

| Ingredient Name                             | Strength |
|---|----------|
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)    |          |
| FD&C RED NO. 40 (UNII: WZB9127XOA)          |          |
| GLYCERIN (UNII: PDC6A3C0OX)                 |          |
| HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S) |          |
| MENTHOL (UNII: L7T10EIP3A)                  |          |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3)         |          |
| WATER (UNII: 059QF0KO0R)                    |          |
| SODIUM BENZOATE (UNII: OJ245FE5EU)          |          |
| SODIUM CITRATE (UNII: 1Q73Q2JULR)           |          |
| SUCRALOSE (UNII: 96K6UQ3ZD4)                |          |

### Product Characteristics

|          |                  |              |  |
|----------|------------------|--------------|--|
| Color    | RED (Orange-Red) | Score        |  |
| Shape    |                  | Size         |  |
| Flavor   | CHERRY           | Imprint Code |  |
| Contains |                  |              |  |

### Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|-----------|---------------------|----------------------|--------------------|
|---|-----------|---------------------|----------------------|--------------------|

|                              |   |   |                             |                           |
|------------------------------|---|---|-----------------------------|---------------------------|
| 1                            | NDC:55312-359-26                                | 1 in 1 CARTON   |                             |                           |
| 1                            |   | 118 mL in 1 BOTTLE; Type 0: Not a Combination Product |                             |                           |
| 2                            | NDC:55312-359-34                                | 1 in 1 CARTON   |                             |                           |
| 2                            |   | 237 mL in 1 BOTTLE; Type 0: Not a Combination Product |                             |                           |
| <b>Marketing Information</b> |   |   |                             |                           |
| <b>Marketing Category</b>    | <b>Application Number or Monograph Citation</b> |   | <b>Marketing Start Date</b> | <b>Marketing End Date</b> |
| OTC monograph final          | part341   |   | 08/09/1991                  |                           |

**Labeler** - Western Family Foods Inc (192166072)

Revised: 9/2015

Western Family Foods Inc